

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 June 2003 (05.06.2003)

PCT

(10) International Publication Number
WO 03/045912 A1

(51) International Patent Classification⁷: C07D 207/16,
207/26, 213/64, 401/12, 401/14, 403/12, A61K 31/40,
A61P 7/02

(21) International Application Number: PCT/IB02/04757

(22) International Filing Date:
14 November 2002 (14.11.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/334,168 29 November 2001 (29.11.2001) US
60/384,895 31 May 2002 (31.05.2002) US

(71) Applicant (for all designated States except US):
WARNER-LAMBERT COMPANY LLC [US/US];
201 Tabor Road, Morris Plains, NJ 070950 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BIGGE, Christopher, Franklin** [US/US]; Pfizer Global Research and Development, Ann Arbor Laboratories, 2800 Plymouth Road, Ann Arbor, MI 48105 (US). **DUDLEY, Danette, Andrea** [US/US]; Pfizer Global Research and Development, Ann Arbor Laboratories, 2800 Plymouth Road, Ann Arbor, MI 48105 (US). **EDMUNDS, Jeremy, John** [US/US]; Pfizer Global Research and Development, Ann Arbor Laboratories, 2800 Plymouth Road, Ann Arbor, MI 48105 (US). **VAN HUIS, Chad, Alan** [US/US]; Pfizer Global Research and Development, Ann Arbor Laboratories, 2800 Plymouth Road; Ann Arbor, MI 48105 (US). **CASIMIRO-GARCIA, Agustin** [MX/US]; Pfizer Global Research and Development, Ann Arbor Laboratories, 2800 Plymouth Road, Ann Arbor, MI 48105 (US).

FILIPSKI, Kevin, James [US/US]; Pfizer Global Research and Development, Ann Arbor Laboratories, 2800 Plymouth Road, Ann Arbor, MI 48105 (US). **KOHR, Jeffrey, Thomas** [US/US]; Pfizer Global Research and Development, Ann Arbor Laboratories, 2800 Plymouth Road, Ann Arbor, MI 48105 (US).

(74) Agents: **LUMB, J., Trevor** et al.; Pfizer Inc., 201 Tabor Road, Morris Plains, NJ 07950 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: INHIBITORS OF FACTOR XA AND OTHER SERINE PROTEASES INVOLVED IN THE COAGULATION CASCADE



(57) Abstract: The present invention provides compounds of Formula (I), wherein A, B, C, G, and W¹ have any of the values defined in the specification, and pharmaceutically acceptable salt thereof, that are useful to treat thrombotic disorders. Also disclosed are pharmaceutical compositions comprising one or more compounds of Formula (I), processes for preparing compounds of Formula (I), and intermediates useful for preparing compounds of Formula (I).

WO 03/045912 A1

-1-

INHIBITORS OF FACTOR Xa AND OTHER SERINE PROTEASES INVOLVED IN THE COAGULATION CASCADE

FIELD OF THE INVENTION

5 The present invention relates to cyclic amino acid and proline derivatives which display inhibitory effects of the serine protease factor Xa. The invention also discloses pharmaceutically acceptable salts of the compounds, pharmaceutically acceptable compositions comprising the compounds or their salts, and methods of using them as therapeutic agents for treating or preventing
10 disease states in mammals characterized by abnormal thrombosis.

BACKGROUND OF THE INVENTION

In economically developed countries, cardiovascular disease represents a major cause of mortality. In particular, abnormal coagulation and inappropriate
15 thrombus formation within blood vessels precipitates many acute cardiovascular disease states. While it has long been recognized that a variety of plasma proteins such as fibrinogen, serine proteases, and cellular receptors are involved in hemostasis, it is abnormal regulation that has emerged as important contributing factors to cardiovascular disease.

20 Thrombin can be considered the key or principal regulatory enzyme in the coagulation cascade; it serves a pluralistic role as both a positive and negative feedback regulator in normal hemostasis. However, in some pathologic conditions, the positive feedback regulation is amplified through catalytic
25 activation of cofactors required for thrombin generation. Such cofactors include factor Xa, a serine protease which occupies a pivotal position in the coagulation cascade. Factor X is the zymogen of factor Xa. Factor X can be activated by either the intrinsic or extrinsic pathways of the coagulation system. Initiation of coagulation by either pathway in response to vascular injury activates factor X to
30 factor Xa. Factor Xa and its cofactor, factor Va, combine on a phospholipid membrane to form the "prothrombinase" complex, which activates prothrombin to thrombin. Thrombin cleaves fibrinogen to fibrin, activates platelets, and converts

-2-

factor XIII to XIIIa which is the principal enzyme involved in thrombus generation, growth, and stabilization. Accordingly, the location of the prothrombinase complex at the convergence of both the intrinsic and extrinsic coagulation pathways suggests that inhibition of factor Xa, and hence thrombin generation, may be a viable approach to limiting the procoagulant activity of thrombin.

Evidence exists for the role of factor Xa inhibitors as anticoagulants. Antistasin, a potent inhibitor of blood coagulation factor Xa from the Mexican leech, *Haementeria officinalis*, displays antithrombotic activity in various models of arterial and venous thrombosis (Lapatto et al., *Embo. J.*, 1997:5151-5161). Other protein or polypeptide factor Xa inhibitors include recombinant tick anticoagulant peptide (rTAP), which is known to accelerate the recombinant tissue plasminogen activator mediated clot lysis and prevent acute reocclusion in the dog, hence indicating factor Xa inhibitors may be useful as an adjunct to thrombolytic therapy (Mellott et al., *Fibrinolysis*, 1993: 195-202). Furthermore, in a canine coronary artery electrolytic lesion model, rTAP was demonstrated to reduce thrombus mass and time to occlusion in the absence of dramatic hemodynamic or hemostatic changes indicating the primary role for factor Xa in the process of arterial thrombosis (Lynch et al., *Thromb. Haemostasis*, 1995:640-645; Schaffer et al., *Circulation*, 1991: 1741- 1748). On the venous side, rTAP was also demonstrated to reduce fibrin deposition in a rabbit model of venous thrombosis while having little affect on systemic hemostatic parameters (Fioravanti et al., *Thromb. Res.*, 1993: 317-324). In addition to these relatively high molecular weight proteins that are not suitable as oral antithrombotic agents, there also exist examples of low molecular weight factor Xa inhibitors. In particular DX9065a, a low molecular weight synthetic factor Xa inhibitor, has also shown antithrombotic potential in various experimental thrombosis rat models. In both arteriovenous shunt and venous stasis models, inhibition of thrombus formation was achieved at doses that had little effect on APTT, indicating that DX9065a is effective in preventing thrombosis and hence has therapeutic antithrombotic potential (Wong et al., *Thromb. Res.*, 1996: 117-126).


-3-

Recently, it has been appreciated that factor Xa inhibition may provide sustained antithrombotic protection. Specifically, several animal studies show that inhibition of short term exposure to factor Xa produces a sustained antithrombotic. (Leadley, *Curr. Top. Med. Chem.*, 2001: v. 1, 151-159.) Finally, the article by Leadley observes that factor Xa inhibition potentially provides a large therapeutical window between antithrombotic efficacy and bleeding tendency. Consequently, there may exist a range in which factor Xa inhibition is achieved without an concurrent increase in a patients susceptibility to bleeding.

The majority of factor Xa inhibitors known to date have been summarized in two reviews (Edmunds et al., *Annual Reports in Medicinal Chemistry*, 1996:51 and Kunitada and Nagahara, *Curr. Pharm. Des.*, 1996:531-542). However, it is readily apparent that there still exists a need for more effective agents that regulate factor Xa proteolytic activity.

SUMMARY OF THE INVENTION

These and other needs are met by the present invention which is directed to a compound of Formula I




I

or a pharmaceutically acceptable salt thereof wherein:

A is aryl or substituted aryl or monocyclic heteroaryl or substituted monocyclic heteroaryl;

B is $\text{-NH-C(=O)-(C}_1\text{-C}_6\text{)alkyl}$, $\text{-NH-C(=O)-(C}_3\text{-C}_7\text{)cycloalkyl}$, -

-4-

$\text{NH}-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}(\text{C}_1-\text{C}_6 \text{ alkyl}),$ , $(\text{C}_3-\text{C}_7)\text{cycloalkyl}, (\text{C}_3-\text{C}_7)\text{heterocyclo}, (\text{C}_4-\text{C}_7)\text{cycloalkenyl},$ unsaturated $(\text{C}_4-\text{C}_7)\text{heterocyclo},$ aryl, or heteroaryl, any of which may be optionally substituted by halo, $(\text{C}_1-\text{C}_6)\text{alkyl},$ or halo $(\text{C}_1-\text{C}_6)\text{alkyl},$ O- $(\text{C}_1-\text{C}_6),$ -CN, haloalkyl, amino, alkylamino, amidino, amido, or sulfonamido;

5 C is phenyl or heteroaryl, wherein phenyl or heteroaryl is optionally substituted with one or more substituents selected from aryl, heteroaryl, halogen, hydroxy, $-\text{CO}_2\text{R}^2,$ $-\text{COR}^2,$ $-\text{CONR}^2\text{R}^{2'},$ alkoxy, alkyl, -CN, haloalkyl, amino, alkylamino, amidino, amido, or sulfonamido;

10 G is H, halo, $(\text{C}_1-\text{C}_6)\text{alkyl},$ halo $(\text{C}_1-\text{C}_6)\text{alkyl},$ hydroxy $(\text{C}_1-\text{C}_6)\text{alkyl},$ $-\text{CH}_2\text{O}-(\text{C}_1-\text{C}_6)\text{alkyl},$ $-\text{CH}_2-\text{CO}_2(\text{C}_1-\text{C}_6)\text{alkyl},$ $-\text{CH}_2-\text{NR}^2\text{R}^{2'},$ or $-\text{CH}_2-\text{CONH}(\text{C}_1-\text{C}_6)\text{alkyl};$

15 W^1 is a saturated or unsaturated, substituted or unsubstituted hydrocarbon chain or hydrocarbon-heteroatom chain having from 2 to 6 atoms, wherein W^1 connects the nitrogen atom at position 1 to the carbon atom at position 2 to form a four to eight membered ring;


R^1 is $(\text{C}_1-\text{C}_6)\text{alkoxy}, (\text{C}_3-\text{C}_7)\text{cycloalkyl}, (\text{C}_3-\text{C}_7)\text{heterocycloalkyl}, (\text{C}_4-\text{C}_7)\text{cycloalkenyl}, (\text{C}_4-\text{C}_7)\text{heterocycloalkenyl},$ aryl, monocyclic heteroaryl, or $-\text{NR}^3\text{R}^4;$

R^2 and $\text{R}^{2'}$ are each independently H or $(\text{C}_1-\text{C}_6)\text{alkyl};$ and

20 R^3 and R^4 are each independently H, $(\text{C}_1-\text{C}_6)\text{alkyl},$ aralkyl, aryl, monocyclic heteroaryl, alkoxycarbonyl, aralkoxycarbonyl, $-\text{SO}_2\text{alkyl},$ or joined together to form a saturated or unsaturated 3 to 7 membered ring.

The invention is also directed to a compound of formula II:

-5-



II

or a pharmaceutically acceptable salt thereof wherein

5 “---” is a bond or is absent;

X^{II} is CH_2 , CH , O , NH or N ;

R^{10} and R^{11} are each independently H , $-OH$, halo, alkyl, haloalkyl, $-NR^8R^9$, $-OR^2$, $-CN$, $-CH_2OH$, $-CH_2-NR^3R^4$, aryl, monocyclic heteroaryl, alkylaryl, $-CH=O$, $-CH_2OR^2$, $-COR^2$, $-CO_2R^2$, or $-CONR^3R^4$ or are taken

10 together to form $=O$, $=NOR^2$, $=C(C_1-C_6alkyl)_2$, or $=CR^2H$, with the proviso that when “---” is a bond, R^{11} is absent, and X^{II} is CH or N , and R^{10} is H , $-OH$, halo, alkyl, haloalkyl, $-NR^8R^9$, $-OR^2$, $-CN$, $-CH_2OH$, $-CH_2-NR^3R^4$, aryl, monocyclic heteroaryl, alkylaryl, $-CH=O$, $-CH_2OR^2$, $-COR^2$, $-CO_2R^2$, or $-CONR^3R^4$;

15 A is aryl and substituted aryl or monocyclic heteroaryl or substituted monocyclic heteroaryl;

B is $-NH-C(=O)-(C_1-C_6)alkyl$, $-NH-C(=O)-(C_3-C_7)cycloalkyl$, -

$NH-C(=O)-O(C_1-C_6 alkyl)$, $-C(=O)-R^1$, $(C_3-C_7)cycloalkyl$, $(C_3-C_7)heterocyclo$, $(C_4-C_7)cycloalkenyl$, unsaturated $(C_4-C_7)heterocyclo$, aryl, or heteroaryl, any of which may be optionally substituted by halo, $(C_1-C_6)alkyl$, or halo $(C_1-C_6)alkyl$,

20 $O-(C_1-C_6)$, $-CN$, haloalkyl, amino, alkylamino, amidino, amido, or sulfonamido;

C is phenyl or heteroaryl, wherein phenyl or heteroaryl is optionally substituted with one or more substituents selected from halogen, hydroxy, -

-6-

CO_2R^2 , $-\text{COR}^2$, $-\text{CONR}^2\text{R}^{2'}$, alkoxy, alkyl, $-\text{CN}$, haloalkyl, amino, alkylamino, amidino, amido, or sulfonamido;


G is H, halo, $(\text{C}_1\text{-C}_6)\text{alkyl}$, halo $(\text{C}_1\text{-C}_6)\text{alkyl}$, hydroxy $(\text{C}_1\text{-C}_6)\text{alkyl}$, $-\text{CH}_2\text{O}(\text{C}_1\text{-C}_6)\text{alkyl}$, $-\text{CH}_2\text{CO}_2(\text{C}_1\text{-C}_6)\text{alkyl}$, $-\text{CH}_2\text{NR}^2\text{R}^{2'}$, or $-\text{CH}_2\text{CONH}(\text{C}_1\text{-C}_6)\text{alkyl}$; or $-\text{CH}_2\text{CONH}(\text{C}_1\text{-C}_6)\text{alkyl}$;

R^1 is $(\text{C}_1\text{-C}_6)\text{alkoxy}$, $(\text{C}_3\text{-C}_7)\text{cycloalkyl}$, $(\text{C}_3\text{-C}_7)\text{heterocycloalkyl}$, $(\text{C}_4\text{-C}_7)\text{cycloalkenyl}$, $(\text{C}_4\text{-C}_7)\text{heterocycloalkenyl}$, aryl, monocyclic heteroaryl, or $-\text{NR}^3\text{R}^4$;

R^2 and $\text{R}^{2'}$ are each independently H or $(\text{C}_1\text{-C}_6)\text{alkyl}$; and

R^3 and R^4 are independently H, $(\text{C}_1\text{-C}_6)\text{alkyl}$, aralkyl, aryl, monocyclic heteroaryl, alkoxy carbonyl, aralkoxy carbonyl, $-\text{SO}_2\text{alkyl}$, or joined together to form a saturated or unsaturated 5 to 7 membered ring.

The invention is also directed to a compound of Formula III



or a pharmaceutically acceptable salt thereof wherein

"---" is a bond or is absent;

Z is C-H, C-halo, C- $(\text{C}_1\text{-C}_6)\text{alkyl}$, C-halo $(\text{C}_1\text{-C}_6)\text{alkyl}$, C- $(\text{C}_1\text{-C}_6)\text{alkoxy}$, or N;

X^{II} is CH_2 , CH, O, NH or N;

-7-

R^{10} and R^{11} are each independently H, -OH, halo, alkyl, haloalkyl, -
 NR^8R^9 , -OR², -CN, -CH₂OH, -CH₂-NR³R⁴, aryl, monocyclic heteroaryl,
 alkylaryl, -CH=O, -CH₂OR², -COR², -CO₂R², or -CONR³R⁴ or are taken
 together to form =O, =NOR², =C(C₁-C₆alkyl)₂, or =CR²H, with the proviso that
 5 when " --- " is a bond, R^{11} is absent, and X^{II} is CH or N, and R^{10} is H, -OH, halo,
 alkyl, haloalkyl, -NR⁸R⁹, -OR², -CN, -CH₂OH, -CH₂-NR³R⁴, aryl, monocyclic
 heteroaryl, alkylaryl, -CH=O, -CH₂OR², -COR², -CO₂R², or -CONR³R⁴;

A is aryl and substituted aryl or monocyclic heteroaryl or substituted
 monocyclic heteroaryl;

10 B is $-NH-\overset{\overset{O}{\parallel}}{C}-(C_1-C_6)alkyl$, $-NH-\overset{\overset{O}{\parallel}}{C}-(C_3-C_7)cycloalkyl$, -

$NH-\overset{\overset{O}{\parallel}}{C}-O(C_1-C_6 alkyl)$, $\overset{\overset{O}{\parallel}}{C}-R^1$, (C₃-C₇)cycloalkyl, (C₃-C₇)heterocyclo, (C₄-
 C₇)cycloalkenyl, unsaturated (C₄-C₇)heterocyclo, aryl, or heteroaryl, any of which
 may be optionally substituted by halo, (C₁-C₆)alkyl, or halo(C₁-C₆)alkyl,
 O-(C₁-C₆), -CN, haloalkyl, amino, alkylamino, amidino, amido, or sulfonamido;

15 G is H, halo, (C₁-C₆)alkyl, halo(C₁-C₆)alkyl, hydroxy(C₁-C₆)alkyl,
 -CH₂O-(C₁-C₆)alkyl, -CH₂-CO₂(C₁-C₆)alkyl, -CH₂-CONH₂, or -CH₂-CONH(C₁-
 C₆)alkyl;


R^2 and $R^{2'}$ are each independently H or (C₁-C₆)alkyl; and

R^3 and R^4 are independently H, (C₁-C₆)alkyl, aralkyl, aryl, monocyclic
 20 heteroaryl, alkoxycarbonyl, aralkoxycarbonyl, -SO₂alkyl, or joined together to
 form a saturated or unsaturated 5 to 7 membered ring; and

R^{12} and R^{13} are each independently H, halo, (C₁-C₆)alkyl, halo(C₁-
 C₆)alkyl, or (C₁-C₆)alkoxy.

25 What is also provided is a compound of Formula IV

-8-



IV

or a pharmaceutically acceptable salt thereof wherein

5 "----" is a bond or is absent;

Z is C-H, C-halo, C-(C₁-C₆)alkyl, C-halo(C₁-C₆)alkyl, C-(C₁-C₆)alkoxy, or

N;

X^{II} is CH₂, CH, O, NH or N;

R¹⁰ and R¹¹ are each independently H, -OH, halo, alkyl, haloalkyl, -

10 NR⁸R⁹, -OR², -CN, -CH₂OH, -CH₂-NR³R⁴, aryl, monocyclic heteroaryl,

alkylaryl, -CH=O, -CH₂OR², -COR², -CO₂R², or -CONR³R⁴ or are taken


together to form =O, =NOR², =C(C₁-C₆alkyl)₂, or =CR²H, with the proviso that

when "----" is a bond, R¹¹ is absent, and X^{II} is CH or N, and R¹⁰ is H, -OH, halo,

alkyl, haloalkyl, -NR⁸R⁹, -OR², -CN, -CH₂OH, -CH₂-NR³R⁴, aryl, monocyclic

15 heteroaryl, alkylaryl, -CH=O, -CH₂OR², -COR², -CO₂R², or -CONR³R⁴;

B is -NH-C(=O)-(C₁-C₆)alkyl, -NH-C(=O)-(C₃-C₇)cycloalkyl, -

NH-C(=O)-O(C₁-C₆ alkyl),  R¹, (C₃-C₇)cycloalkyl, (C₃-C₇)heterocyclo, (C₄-C₇)cycloalkenyl, unsaturated (C₄-C₇)heterocyclo, aryl, or heteroaryl, any of which may be optionally substituted by halo, (C₁-C₆)alkyl, or halo(C₁-C₆)alkyl,

-9-

O-(C₁-C₆), -CN, haloalkyl, amino, alkylamino, amidino, amido, or sulfonamido;

G is H, halo, (C₁-C₆)alkyl, halo(C₁-C₆)alkyl, hydroxy(C₁-C₆)alkyl, -CH₂O-(C₁-C₆)alkyl, -CH₂-CO₂(C₁-C₆)alkyl, -CH₂-CONH₂, or -CH₂-CONH(C₁-C₆)alkyl;

5 R² and R^{2'} are each independently H or (C₁-C₆)alkyl; and

R³ and R⁴ are independently H, (C₁-C₆)alkyl, aralkyl, aryl, monocyclic heteroaryl, alkoxy carbonyl, aralkoxy carbonyl, -SO₂alkyl, or joined together to form a saturated or unsaturated 5 to 7 membered ring;


10 R¹² and R¹³ are each independently H, halo, (C₁-C₆)alkyl, or halo(C₁-C₆)alkyl, or (C₁-C₆)alkoxy;

R¹⁴ and R¹⁵ are each independently H, halo, alkyl, or haloalkyl, or NR⁸R⁹ wherein R⁸ and R⁹ are as defined for R³ and R⁴;

X and Y are each C, or one of X and Y is C and the other is N, provided that when one of X or Y is N, R¹⁴ or R¹⁵ is absent at that position.

15

What is also provided is a compound of Formula V



V

20

or a pharmaceutically acceptable salt thereof wherein

"---" is a bond or is absent;

Z is C-H, C-halo, C-(C₁-C₆)alkyl, C-halo(C₁-C₆)alkyl, C-(C₁-C₆)alkoxy, or

-10-

N;

G is H, halo, (C₁-C₆)alkyl, halo(C₁-C₆)alkyl, hydroxy(C₁-C₆)alkyl, -CH₂O-(C₁-C₆)alkyl, -CH₂-CO₂(C₁-C₆)alkyl, -CH₂-CONH₂, or -CH₂-CONH(C₁-C₆)alkyl;

5 X^{II} is CH₂, CH, O, NH or N;

R¹⁰ and R¹¹ are each independently H, -OH, halo, alkyl, haloalkyl, -NR⁸R⁹, -OR², -CN, -CH₂OH, -CH₂-NR³R⁴, aryl, monocyclic heteroaryl, alkylaryl, -CH=O, -CH₂OR², -COR², -CO₂R², or -CONR³R⁴ or are taken together to form =O, =NOR², =C(C₁-C₆alkyl)₂, or =CR²H, with the proviso that
 10 when "----" is a bond, R¹¹ is absent, and X^{II} is CH or N, and R¹⁰ is H, -OH, halo, alkyl, haloalkyl, -NR⁸R⁹, -OR², -CN, -CH₂OH, -CH₂-NR³R⁴, aryl, monocyclic heteroaryl, alkylaryl, -CH=O, -CH₂OR², -COR², -CO₂R², or -CONR³R⁴;

R² is H or (C₁-C₆)alkyl; and

R³ and R⁴ are independently H, (C₁-C₆)alkyl, aralkyl, aryl, monocyclic
 15 heteroaryl, alkoxycarbonyl, aralkoxycarbonyl, -SO₂alkyl, or joined together to form a saturated or unsaturated 5 to 7 membered ring;

R¹² and R¹³ are each independently H, halo, (C₁-C₆)alkyl, or halo(C₁-C₆)alkyl, or (C₁-C₆)alkoxy;

R¹⁴ and R¹⁵ are each independently H, halo, alkyl, or haloalkyl, or NR⁸R⁹
 20 wherein R⁸ and R⁹ are as defined for R³ and R⁴;

X and Y are each independently C or N, provided that when one of X or Y is N, R¹⁴ or R¹⁵ is absent at that position.

X' and Y' are each independently CH or N;



is absent or is a fused heterocyclic or fused heteroaryl ring,




25 provided that when is present, Y' is C; and

R¹⁶ and R¹⁷ are each independently H, halo, cyano, (C₁-C₆)alkoxy

-11-

carbonyl, aminomethyl, t-butyl, H_2NSO_2 —, Me_2NSO_2 —, $(\text{C}_1\text{-C}_6)\text{alkoxy}$,
 MeSO_2 —, MeSO —, MeS —, NR^8R^9 , or hydroxy

What is also provided is a compound of Formula VI



VI

or a pharmaceutically acceptable salt thereof wherein

“---” is a bond or is absent;

10 Z is C-H, C-halo, C-($\text{C}_1\text{-C}_6$)alkyl, C-halo($\text{C}_1\text{-C}_6$)alkyl, C-($\text{C}_1\text{-C}_6$)alkoxy, or N;

G is H, halo, ($\text{C}_1\text{-C}_6$)alkyl, halo($\text{C}_1\text{-C}_6$)alkyl, hydroxy($\text{C}_1\text{-C}_6$)alkyl, - CH_2O -($\text{C}_1\text{-C}_6$)alkyl, - CH_2CO_2 ($\text{C}_1\text{-C}_6$)alkyl, - CH_2CONH_2 , or - CH_2CONH ($\text{C}_1\text{-C}_6$)alkyl;

15 Q is NH or is absent;

X^{II} is CH_2 , CH, O, NH or N;

R^{10} and R^{11} are each independently H, -OH, halo, alkyl, haloalkyl, - NR^8R^9 , - OR^2 , -CN, - CH_2OH , - $\text{CH}_2\text{-NR}^3\text{R}^4$, aryl, monocyclic heteroaryl, alkylaryl, -CH=O, - CH_2OR^2 , - COR^2 , - CO_2R^2 , or - CONR^3R^4 or are taken

20 together to form =O, = NOR^2 , = $\text{C}(\text{C}_1\text{-C}_6\text{alkyl})_2$, or = CR^2H , with the proviso that when “---” is a bond, R^{11} is absent, and X^{II} is CH or N, and R^{10} is H, -OH, halo,

-12-

alkyl, haloalkyl, $-NR^8R^9$, $-OR^2$, $-CN$, $-CH_2OH$, $-CH_2-NR^3R^4$, aryl, monocyclic heteroaryl, alkylaryl, $-CH=O$, $-CH_2OR^2$, $-COR^2$, $-CO_2R^2$, or $-CONR^3R^4$;

R^1 is OH, $-O-(C_1-C_6)alkyl$, $(C_3-C_7)cycloalkyl$, $(C_3-C_7)heterocycloalkyl$, $(C_4-C_7)cycloalkenyl$, $(C_4-C_7)heterocycloalkenyl$, aryl, monocyclic heteroaryl, or –

5 NR^3R^4 , wherein R^3 and R^4 are independently H, $(C_1-C_6)alkyl$, aralkyl, aryl, monocyclic heteroaryl, alkoxycarbonyl, aralkoxycarbonyl, $-SO_2alkyl$, or joined together to form a saturated or unsaturated 5 to 7 membered ring;

R^2 is H or $(C_1-C_6)alkyl$; and

R^3 and R^4 are independently H, $(C_1-C_6)alkyl$, aralkyl, aryl, monocyclic
10 heteroaryl, alkoxycarbonyl, aralkoxycarbonyl, $-SO_2alkyl$, or joined together to form a 5 to 7 membered ring;

R^{12} and R^{13} are each independently H, halo, $(C_1-C_6)alkyl$, or halo $(C_1-C_6)alkyl$, or $(C_1-C_6)alkoxy$;


R^{14} and R^{15} are each independently H, halo, alkyl, or haloalkyl, or NR^8R^9
15 wherein R^8 and R^9 are as defined for R^3 and R^4 ;

X and Y are each independently C or N, provided that when one of X or Y is N, R^{14} or

R^{15} is absent at that position.

20 What is also provided is a compound of Formula VII

-13-



VII

or a pharmaceutically acceptable salt thereof wherein

- 5 G is H, halo, (C₁-C₆)alkyl, halo(C₁-C₆)alkyl, hydroxy(C₁-C₆)alkyl, -CH₂O-(C₁-C₆)alkyl, -CH₂-CO₂(C₁-C₆)alkyl, -CH₂-CONH₂, or -CH₂-CONH(C₁-C₆)alkyl;

X^{II} is CH₂, CH, O, NH or N;

- R^{10} and R^{11} are each independently H, -OH, halo, alkyl, haloalkyl, -
 10 NR⁸R⁹, -OR², -CN, -CH₂OH, -CH₂-NR³R⁴, aryl, monocyclic heteroaryl, alkylaryl, -CH=O, -CH₂OR², -COR², -CO₂R², or -CONR³R⁴ or are taken together to form =O, =NOR², =C(C₁-C₆alkyl)₂, or =CR²H, with the proviso that when "----" is a bond, R^{11} is absent, and X^{II} is CH or N, and R^{10} is H, -OH, halo, alkyl, haloalkyl, -NR⁸R⁹, -OR², -CN, -CH₂OH, -CH₂-NR³R⁴, aryl, monocyclic
 15 heteroaryl, alkylaryl, -CH=O, -CH₂OR², -COR², -CO₂R², or -CONR³R⁴;

R^2 and $R^{2'}$ are each independently H or (C₁-C₆)alkyl; and

R^3 and R^4 are each independently H, (C₁-C₆)alkyl, aralkyl, aryl, monocyclic heteroaryl, alkoxycarbonyl, aralkoxycarbonyl, -SO₂alkyl, or joined together to form a saturated or unsaturated 3 to 7 membered ring;


- 20 R^{12} and R^{13} are each independently H, halo, (C₁-C₆)alkyl, or halo(C₁-C₆)alkyl, or (C₁-C₆)alkoxy;

-14-

R^{14} and R^{15} are each independently H, halo, alkyl, or haloalkyl, or NR^8R^9 wherein R^8 and R^9 are as defined for R^3 and R^4 ;

X and Y are each independently C or N, provided that when one of X or Y is N, R¹⁴ or

5 R¹⁵ is absent at that position; and



is an optionally substituted 4, 5, 6 or 7-membered


ring;

wherein R^x and R^y are H, halo, hydroxymethyl, (C₁-C₆)alkyl, (C₁-

C₆)alkoxy, or taken together are O=;

10 X^{III} is CH_2 , O, S, NH, or $N(C_1-C_6)alkyl$, provided that when R^x and R^y taken together are $O=$, X^{III} is CH_2 .

What is also provided is a compound of Formula VIII



VIII

or a pharmaceutically acceptable salt or tautomer thereof wherein

"---" is a bond or is absent;

Z is C-H, C-halo, C-(C₁-C₆)alkyl, C-halo(C₁-C₆)alkyl, C-(C₁-C₆)alkoxy, or

20 N;

-15-

G is H, halo, (C₁-C₆)alkyl, halo(C₁-C₆)alkyl, hydroxy(C₁-C₆)alkyl, -CH₂O-(C₁-C₆)alkyl, -CH₂-CO₂(C₁-C₆)alkyl, -CH₂-CONH₂, or -CH₂-CONH(C₁-C₆)alkyl;

X^{II} is CH₂, CH, O, NH or N;

5 R¹⁰ and R¹¹ are each independently H, -OH, halo, alkyl, haloalkyl, -NR⁸R⁹, -OR², -CN, -CH₂OH, -CH₂-NR³R⁴, aryl, monocyclic heteroaryl, alkylaryl, -CH=O, -CH₂OR², -COR², -CO₂R², or -CONR³R⁴ or are taken together to form =O, =NOR², =C(C₁-C₆alkyl)₂, or =CR²H, with the proviso that when "----" is a bond, R¹¹ is absent, and X^{II} is CH or N, and R¹⁰ is H, -OH, halo,
10 alkyl, haloalkyl, -NR⁸R⁹, -OR², -CN, -CH₂OH, -CH₂-NR³R⁴, aryl, monocyclic heteroaryl, alkylaryl, -CH=O, -CH₂OR², -COR², -CO₂R², or -CONR³R⁴;

R² and R^{2'} are each independently H or (C₁-C₆)alkyl; and

R³ and R⁴ are each independently H, (C₁-C₆)alkyl, aralkyl, aryl, monocyclic heteroaryl, alkoxy carbonyl, aralkoxy carbonyl, -SO₂alkyl, or joined
15 together to form a saturated or unsaturated 3 to 7 membered ring;

R¹² and R¹³ are each independently H, halo, (C₁-C₆)alkyl, or halo(C₁-C₆)alkyl, or (C₁-C₆)alkoxy;

R¹⁴ and R¹⁵ are each independently H, halo, alkyl, or haloalkyl, or NR⁸R⁹ wherein R⁸ and R⁹ are as defined for R³ and R⁴;

20 X and Y are each independently C or N, provided that when one of X or Y is N, R¹⁴ or

R¹⁵ is absent at that position; and


Y^{II} is CH₂, CH(C₁-C₆alkyl), NH, or N(C₁-C₆alkyl); and

R¹⁸ is H, (C₁-C₆)alkyl, hydroxymethyl, CH₂O-(C₁-C₆)alkyl, or NR³R⁴.

25

What is also provided is a compound of Formula IX

-16-



IX

or a pharmaceutically acceptable salt thereof wherein

5 “---” is a bond or is absent;

Z is C-H, C-halo, C-(C₁-C₆)alkyl, C-halo(C₁-C₆)alkyl, C-(C₁-C₆)alkoxy, or N;

G is H, halo, (C₁-C₆)alkyl, halo(C₁-C₆)alkyl, hydroxy(C₁-C₆)alkyl, -CH₂O-(C₁-C₆)alkyl, -CH₂-CO₂(C₁-C₆)alkyl, -CH₂-CONH₂, or -CH₂-CONH(C₁-C₆)alkyl;

10 X^{II} is CH_2 , CH , NH or N ;

R^{10} and R^{11} are each independently H, -OH, halo, alkyl, haloalkyl, - NR^8R^9 , -OR², -CN, -CH₂OH, -CH₂-NR³R⁴, aryl, monocyclic heteroaryl, alkylaryl, -CH=O, -CH₂OR², -COR², -CO₂R², or -CONR³R⁴ or are taken together to form =O, =NOR², =C(C₁-C₆alkyl)₂, or =CR²H, with the proviso that when "---" is a bond, R¹¹ is absent, and X^{II} is CH or N, and R¹⁰ is H, -OH, halo, alkyl, haloalkyl, -NR⁸R⁹, -OR², -CN, -CH₂OH, -CH₂-NR³R⁴, aryl, monocyclic heteroaryl, alkylaryl, -CH=O, -CH₂OR², -COR², -CO₂R², or -CONR³R⁴;

R² and R^{2'} are each independently H or (C₁-C₆)alkyl; and

R³ and R⁴ are each independently H, (C₁-C₆)alkyl, aralkyl, aryl,
20 monocyclic heteroaryl, alkoxycarbonyl, aralkoxycarbonyl, -SO₂alkyl, or joined
together to form a saturated or unsaturated 3 to 7 membered ring;

-17-

R^{12} and R^{13} are each independently H, halo, (C_1-C_6) alkyl, or halo (C_1-C_6) alkyl, or (C_1-C_6) alkoxy;

R^{14} and R^{15} are each independently H, halo, alkyl, or haloalkyl, or NR^8R^9 wherein R^8 and R^9 are as defined for R^3 and R^4 ;


5 X and Y are each independently C or N, provided that when one of X or Y is N, R^{14} or R^{15} is absent at that position;

X^a , Y^a , and Z^a are each independently CH, CR^{18} or N; and

R^{18} is H, (C_1-C_6) alkyl, hydroxymethyl, $CH_2O-(C_1-C_6)$ alkyl, CH^2NR^2 , or NR^3R^4 .

10

The invention also provides a compound of Formula X



X

15 or a pharmaceutically acceptable salt thereof wherein

"---" is a bond or is absent;

Z is C-H, C-halo, C- (C_1-C_6) alkyl, C-halo (C_1-C_6) alkyl, C- (C_1-C_6) alkoxy, or N;

20 G is H, halo, (C_1-C_6) alkyl, halo (C_1-C_6) alkyl, hydroxy (C_1-C_6) alkyl, $-CH_2O-(C_1-C_6)$ alkyl, $-CH_2CO_2(C_1-C_6)$ alkyl, $-CH_2CONH_2$, or $-CH_2CONH(C_1-C_6)$ alkyl;

X^{II} is CH_2 , CH, O, NH or N;

-18-

R^{10} and R^{11} are each independently H, -OH, halo, alkyl, haloalkyl, -
 NR^8R^9 , -OR², -CN, -CH₂OH, -CH₂-NR³R⁴, aryl, monocyclic heteroaryl,
 alkylaryl, -CH=O, -CH₂OR², -COR², -CO₂R², or -CONR³R⁴ or are taken
 together to form =O, =NOR², =C(C₁-C₆alkyl)₂, or =CR²H, with the proviso that
 5 when " --- " is a bond, R^{11} is absent, and X^{II} is CH or N, and R^{10} is H, -OH, halo,
 alkyl, haloalkyl, -NR⁸R⁹, -OR², -CN, -CH₂OH, -CH₂-NR³R⁴, aryl, monocyclic
 heteroaryl, alkylaryl, -CH=O, -CH₂OR², -COR², -CO₂R², or -CONR³R⁴;

R^2 and $R^{2'}$ are each independently H or (C₁-C₆)alkyl; and

R^3 and R^4 are each independently H, (C₁-C₆)alkyl, aralkyl, aryl,
 10 monocyclic heteroaryl, alkoxycarbonyl, aralkoxycarbonyl, -SO₂alkyl, or joined
 together to form a saturated or unsaturated 3 to 7 membered ring;

R^{12} and R^{13} are each independently H, halo, (C₁-C₆)alkyl, or halo(C₁-
 C₆)alkyl, or (C₁-C₆)alkoxy;

R^{14} and R^{15} are each independently H, halo, alkyl, or haloalkyl, or NR⁸R⁹
 15 wherein R^8 and R^9 are as defined for R^3 and R^4 ;

X and Y are each independently C or N, provided that when one of X or Y
 is N, R^{14} or R^{15} is absent at that position;

J₁, J₂, J₃, and J₄ are C or one of J₁, J₂, J₃, and J₄ is N;

R^{19} , R^{20} , R^{21} , and R^{22} are each independently H, halo, hydroxy, NH₂,
 20 NR²³R²⁴, NO₂, SH, (C₁-C₆)alkyl, halo(C₁-C₆)alkyl, (C₁-C₆)alkanoyl, (C₁-
 C₆)alkoxycarbonyl, or (C₁-C₆)alkoxy, wherein R^{23} and R^{24} are each
 independently H, (C₁-C₆)alkyl, aralkyl, aryl, monocyclic heteroaryl,
 alkoxycarbonyl, aralkoxycarbonyl, -SO₂alkyl, or joined together to form a
 saturated or unsaturated 3 to 7 membered ring, or R^{19} and R^{20} , R^{20} and R^{21} , or R^{21}
 25 and R^{22} , together with the carbons to which they are attached, form a 5, 6, or 7
 membered saturated or unsaturated cycloalkyl or heterocycloalkyl ring, or an aryl
 or heteroaryl ring;

provided that when any of J₁, J₂, J₃, or J₄ is N, R^{19} , R^{20} , R^{21} , or R^{22} is
 absent at that position.

The invention is also directed to a compound which is:

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

5 Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

10 Pyrrolidine-1,2-dicarboxylic acid 1-[(2,4-difluoro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-p-tolylamide;

Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-[(4-methoxy-phenyl)-amide];

15 Pyrrolidine-1,2-dicarboxylic acid 1-[(4-bromo-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-[(4-isopropyl-phenyl)-amide];

20 Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-phenylamide;

Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-[(4-trifluoromethyl-phenyl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-[(3-methoxy-phenyl)-amide];

25 Pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

30 Pyrrolidine-1,2-dicarboxylic acid 1-[(4-ethyl-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(3,4-difluoro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(3-fluoro-4-methyl-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[5-(2-sulfamoyl-phenyl)-pyridin-2-yl]-amide];

5 Pyrrolidine-1,2-dicarboxylic acid 2-[(3-chloro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

10 Pyrrolidine-1,2-dicarboxylic acid 2-[(3-chloro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[5-(2-sulfamoyl-phenyl)-pyridin-2-yl]-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(1H-tetrazol-5-yl)-phenyl]-amide];

15 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

20 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Oxo-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Methylene-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

25 4-Methyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4,4-Difluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

30 4-Amino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Hydroxymethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

1-(4-Chloro-phenylcarbamoyl)-5-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-3-carboxylic acid;

1-(4-Chloro-phenylcarbamoyl)-5-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-3-carboxylic acid methyl ester;

5 Pyrrolidine-1,2,4-tricarboxylic acid 4-amide 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Aminomethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Methoxymethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Methylamino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

15 4-Acetylamino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Methanesulfonylamino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

20 5-Oxo-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

Piperidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

Morpholine-3,4-dicarboxylic acid 4-[(4-chloro-phenyl)-amide] 3-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

25 Piperazine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

2,5-Dihydro-pyrrole-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

30 Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3-methyl-biphenyl-4-yl)-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3-methyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-biphenyl-4-yl)-amide];

5 Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[5-(2-methanesulfonyl-phenyl)-pyrimidin-2-yl]-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[5-(2-methanesulfonyl-phenyl)-pyrimidin-2-yl]-amide];

10 Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[5-(2-methanesulfonyl-phenyl)-3-methyl-pyridin-2-yl]-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[5-(2-methanesulfonyl-phenyl)-3-methyl-pyridin-2-yl]-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3-trifluoromethyl-biphenyl-4-yl)-amide];

15 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3-trifluoromethyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[5-(2-methanesulfonyl-phenyl)-6-methyl-pyridin-2-yl]-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[5-(2-methanesulfonyl-phenyl)-6-methyl-pyridin-2-yl]-amide];

20 Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-ethyl-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-ethyl-2'-methanesulfonyl-biphenyl-4-yl)-amide];

25 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3,5-difluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3,5-dichloro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

30 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3,5-dichloro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3,5-dimethyl-biphenyl-4-yl)-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[(2'-methanesulfonyl-3,5-dimethyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 2-[(3-chloro-2'-methanesulfonyl-5-methyl-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide];

5 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(3-chloro-2'-methanesulfonyl-5-methyl-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 2-[(2-chloro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide];

10 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(2-chloro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-2-methyl-biphenyl-4-yl)-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[(2'-methanesulfonyl-2-methyl-biphenyl-4-yl)-amide];

15 Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[(2-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

20 4-[[1-(4-Chloro-phenylcarbonyl)-pyrrolidine-2-carbonyl]-amino]-2'-methanesulfonyl-biphenyl-3-carboxylic acid methyl ester;

4-[[1-(4-Chloro-phenylcarbonyl)-4-hydroxy-pyrrolidine-2-carbonyl]-amino]-2'-methanesulfonyl-biphenyl-3-carboxylic acid methyl ester;

Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-[(4-trifluoromethyl-phenyl)-amide];

25 Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-m-tolylamide;

Pyrrolidine-1,2-dicarboxylic acid 1-[(3-acetyl-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

30 Pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-2-methyl-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(2-fluoro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

-24-

Pyrrolidine-1,2-dicarboxylic acid 1-[(3-chloro-4-fluoro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(3-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

5 Pyrrolidine-1,2-dicarboxylic acid 1-[(4-cyano-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-2-methyl-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

10 Pyrrolidine-1,2-dicarboxylic acid 1-[(3,4-dichloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-[(3-trifluoromethyl-phenyl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-dimethylamino-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

15 (2S)-Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-cyano-phenyl)-amide];

20 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(4-fluoro-phenyl)-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

25 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-p-tolylamide;

4-Oxo-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]; and

30 4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]; or

a pharmaceutically acceptable salt thereof.

-25-

A compound which is:

(R)-Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

(S)-Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

(2R,4S)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(4-fluoro-phenyl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-p-tolylamide;

(2R)-4-Oxo-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]; and

(2R,4S)-4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]; or a pharmaceutically acceptable salt thereof.

The invention is also directed to a compound which is:

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 5 (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Propoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Propoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 10 (2R,4R)-4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-azetidin-1-yl)-phenyl]-amide];
- 15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-azetidin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-pyrrolidin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-pyrrolidin-1-yl)-phenyl]-amide];
- 20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-azepan-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-azepan-1-yl)-phenyl]-amide];
- 25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- 30 (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

-27-

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-phenyl}-amide};

5 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{4-(2,3-dimethyl-5-oxo-2,5-dihydro-pyrazol-1-yl)-phenyl}-amide};

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{4-(2-hydroxymethyl-pyrrolidin-1-yl)-phenyl}-amide};

10 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-pyrrolidin-1-yl-phenyl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-pyrazol-1-yl-phenyl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-pyrazol-1-yl-phenyl)-amide];

15 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-[1,2,4]triazol-1-yl-phenyl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-[1,2,3]triazol-2-yl-phenyl)-amide];

20 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-[1,2,3]triazol-1-yl-phenyl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(4-acetyl-amino-phenyl)-amide] 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{4-(cyclopentanecarbonyl-amino)-phenyl}-amide}; and

25 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-pyrimidin-5-yl-phenyl)-amide]; or

a pharmaceutically acceptable salt thereof.

A compound which is:

30 1-(4-Chloro-phenylcarbamoyl)-5-[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenylcarbamoyl]-pyrrolidine-3-carboxylic acid;

-28-

1-(4-Chloro-phenylcarbamoyl)-5-[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenylcarbamoyl]-pyrrolidine-3-carboxylic acid methyl ester;

1-(4-Chloro-phenylcarbamoyl)-5-[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenylcarbamoyl]-pyrrolidine-3-carboxylic acid ethyl ester;

5 Pyrrolidine-1,2,4-tricarboxylic acid 4-amide 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

4-Cyano-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

4-(1H-Tetrazol-5-yl)-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

4-Aminomethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

4-Methylaminomethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

15 4-Dimethylaminomethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

4-Acetyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

20 Pyrrolidine-1,2,4-tricarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide} 4-methylamide;

Pyrrolidine-1,2,4-tricarboxylic acid 1-[(4-chloro-phenyl)-amide] 4-dimethylamide 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

4-Trifluoromethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

25 4-Methyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

4-Isopropoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

30 4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

4,4-Difluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

- 1-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-phenyl)-pyrrolidine-2-carboxylic acid;
- 1-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-hydroxy-pyrrolidine-2-carbonyl]-amino}-phenyl)-pyrrolidine-2-carboxylic acid;
- 5 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(5-(2-oxo-piperidin-1-yl)-pyrimidin-2-yl)-amide];
- 10 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(2-(2-oxo-piperidin-1-yl)-pyrimidin-5-yl)-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(5-(2-oxo-piperidin-1-yl)-pyrimidin-2-yl)-amide];
- 15 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-(2-oxo-piperidin-1-yl)-pyrimidin-5-yl)-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide} 1-[(5-fluoro-pyridin-2-yl)-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-fluoro-pyridin-2-yl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 20 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-fluoro-pyridin-2-yl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)-amide];
- 4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 25 4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(5-fluoro-pyridin-2-yl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)-amide];
- 30 4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)-amide];

-30-

- 4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(5-fluoro-pyridin-2-yl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-phenyl]-amide];
- 5 1-(4-[[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-5-oxo-pyrrolidine-2-carboxylic acid;
- 1-(4-[[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-5-oxo-pyrrolidine-2-carboxylic acid;
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(4,4-dimethyl-2,5-dioxo-imidazolidin-1-yl)-2-fluoro-phenyl]-amide];
- 10 [1-(4-[[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-2,5-dioxo-imidazolidin-4-yl]-acetic acid;
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(5,5-dimethyl-2,4-dioxo-oxazolidin-3-yl)-2-fluoro-phenyl]-amide];
- 15 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,5-dioxo-pyrrolidin-1-yl)-2-fluoro-phenyl]-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-oxazolidin-3-yl)-phenyl]-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,5-dioxo-oxazolidin-3-yl)-2-fluoro-phenyl]-amide];
- 20 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-imidazolidin-1-yl)-phenyl]-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,5-dioxo-imidazolidin-1-yl)-2-fluoro-phenyl]-amide];
- 25 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-methyl-2,5-dioxo-imidazolidin-1-yl)-phenyl]-amide];
- 3-(4-[[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-2-oxo-imidazolidine-4-carboxylic acid;
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2,4,5-trioxo-imidazolidin-1-yl)-phenyl]-amide];
- 30 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,5-dioxo-2,5-dihydro-pyrrol-1-yl)-2-fluoro-phenyl]-amide];

- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{[4-(2-aminomethyl-5-oxo-pyrrolidin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{[4-(2-aminomethyl-5-oxo-pyrrolidin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];
- 5 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{[4-(2-carbamoyl-5-oxo-pyrrolidin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{[4-(2-carbamoyl-5-oxo-pyrrolidin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{[4-(2,2-bis-
10 hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];
- 3-(4-{{[1-(4-Chloro-phenyl)carbamoyl]-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-2-oxo-imidazolidine-4-carboxylic acid;
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide].
- 15 2-{{[4-(3,3-dimethyl-2,5-dioxo-pyrrolidin-1-yl)-2-fluoro-phenyl]-amide};
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(3,3,4-trimethyl-2,5-dioxo-pyrrolidin-1-yl)-phenyl]-amide};
- 1-(4-{{[1-(4-Chloro-phenyl)carbamoyl]-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-5-oxo-pyrrolidine-2,2-dicarboxylic acid;
- 20 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(2-methyl-5-oxo-pyrrolidin-1-yl)-phenyl]-amide};
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(2-methyl-6-oxo-piperidin-1-yl)-phenyl]-amide};
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
- 25 2-{{[2-fluoro-4-(3-hydroxy-2-oxo-piperidin-1-yl)-phenyl]-amide};
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(5-methyl-2,6-dioxo-tetrahydro-pyrimidin-1-yl)-phenyl]-amide};
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(2-oxo-tetrahydro-pyrimidin-1-yl)-phenyl]-amide};
- 30 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(3-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[[4-(2,5-dioxo-piperazin-1-yl)-2-fluoro-phenyl]-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[[2-fluoro-4-(3-hydroxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- 5 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[[2-fluoro-4-(3-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[[2-fluoro-4-(6-methyl-2-oxo-4-trifluoromethyl-2H-pyridin-1-yl)-phenyl]-
amide];
- 10 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[[4-(2,6-dioxo-piperidin-1-yl)-2-fluoro-phenyl]-amide];
- 1-(4-[[1-(4-Chloro-phenyl)carbamoyl]-4-methoxy-pyrrolidine-2-carbonyl]-
amino)-3-fluoro-phenyl)-2-oxo-piperidine-3-carboxylic acid;
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
15 2-[[4-(5,5-dihydroxy-2,4,6-trioxo-tetrahydro-pyrimidin-1-yl)-2-fluoro-phenyl]-
amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[[2-fluoro-4-(2,4,6-trioxo-tetrahydro-pyrimidin-1-yl)-phenyl]-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
20 2-[[4-(4,4-dimethyl-2,6-dioxo-piperidin-1-yl)-2-fluoro-phenyl]-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[[2-fluoro-4-(2-hydroxy-4-oxo-azetidin-1-yl)-phenyl]-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(acetyl-methyl-amino)-
2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];
- 25 4-(4-[[1-(4-Chloro-phenyl)carbamoyl]-4-methoxy-pyrrolidine-2-carbonyl]-
amino)-3-fluoro-phenyl)-1H-pyrrole-3-carboxylic acid;
- 2-(4-[[1-(4-Chloro-phenyl)carbamoyl]-4-methoxy-pyrrolidine-2-carbonyl]-
amino)-3-fluoro-phenyl)-2H-pyrazole-3-carboxylic acid;
- 1-(4-[[1-(4-Chloro-phenyl)carbamoyl]-4-methoxy-pyrrolidine-2-carbonyl]-
30 amino)-3-fluoro-phenyl)-1H-imidazole-2-carboxylic acid;
- 1-(4-[[1-(4-Chloro-phenyl)carbamoyl]-4-methoxy-pyrrolidine-2-carbonyl]-
amino)-3-fluoro-phenyl)-1H-pyrrole-2-carboxylic acid;

- 3-(4-{[1-(4-Chloro-phenyl)carbamoyl]-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-1H-pyrrole-2-carboxylic acid;
- 3-(4-{[1-(4-Chloro-phenyl)carbamoyl]-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-furan-2-carboxylic acid;
- 5 3-(4-{[1-(4-Chloro-phenyl)carbamoyl]-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-thiophene-2-carboxylic acid;
- 4-(4-{[1-(4-Chloro-phenyl)carbamoyl]-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-furan-3-carboxylic acid;
- 10 4-(4-{[1-(4-Chloro-phenyl)carbamoyl]-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-thiophene-3-carboxylic acid;
- 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- 3-Oxo-pyrazolidine-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
- 15 3-Oxo-pyrazolidine-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[[4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 5-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 20 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 5-[[4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 25 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 5-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3,5'-difluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(5'-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
- 30 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3,5'-difluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

-34-

- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[(3,5'-difluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[(5'-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- 5 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[(3,5'-difluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-
amide] 2-[(3,5'-difluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
10 2-[(3,5'-difluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
- 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[(3,5'-difluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide], or
a pharmaceutically acceptable salt thereof.
- 15 The invention is also directed to a compound which is:
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[(4-pyrazol-1-yl-phenyl)-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[(2-fluoro-4-pyrazol-1-yl-phenyl)-amide];
- 20 (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[(2-fluoro-4-pyrazol-1-yl-phenyl)-amide];
- (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[(4-pyrazol-1-yl-phenyl)-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
25 phenyl)-amide] 2-[[2-fluoro-4-(3-methyl-pyrazol-1-yl)-phenyl]-amide];
- (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[[2-fluoro-4-(5-methyl-pyrazol-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[[4-(5-methyl-pyrazol-1-yl)-phenyl]-amide];
- 30 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[[4-(3,5-dimethyl-pyrazol-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3,5-dimethyl-pyrazol-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3,5-dimethyl-pyrazol-1-yl)-2-fluoro-phenyl]-amide];

5 (2R,4S)-4-Amino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

(2R,4R)-4-Amino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

10 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

(2R)-4-Hydroxyimino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methylsulfamoyl-biphenyl-4-yl)-amide];

15 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-dimethylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

25 (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

(2R,4S)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

30 (2R,4R)-4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

(2R)-4,4-Difluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(4-fluoro-phenyl)-amide];
- (2R,4R)-4-Acetylamino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- 5 (2R,4R)-4-Methanesulfonylamino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- (2R,4S)-4-(1H-Tetrazol-5-yl)-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- (2R,4S)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 10 (2R,4R)-4-Cyano-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- (2R,4R)-4-(1H-Tetrazol-5-yl)-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 15 (2R,4R)-4-Trifluoromethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-cyano-3-fluoro-biphenyl-4-yl)-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[(2'-aminomethyl-3-fluoro-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide];
- 20 (2R,4R)-4'-[[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3'-fluoro-biphenyl-2-carboxylic acid methyl ester;
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,5-dihydro-pyrrole-1-carbonyl)-phenyl]-amide];
- 25 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(dimethylcarbamoyl-2-fluoro-phenyl)-amide];
- (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-phenyl]-amide];
- Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-phenyl]-amide];
- 30 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2-methyl-pyrrolidine-1-carbonyl)-phenyl]-amide];

-37-

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(ethyl-methyl-carbamoyl)-phenyl]-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-phenyl)-amide];

5 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2R-methyl-pyrrolidine-1-carbonyl)-phenyl]-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2S-methyl-pyrrolidine-1-carbonyl)-phenyl]-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(pyrrolidine-1-carbonyl)-phenyl]-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-2-pyrrolidin-1-yl-phenyl]-amide];

(2R,4R)-4-[[1-(4-Chloro-phenylcarbamoyl)-4-hydroxy-pyrrolidine-2-carbonyl]-amino]-3-pyrrolidin-1-yl-benzoic acid methyl ester;

15 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(azetidine-1-carbonyl)-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(pyrrolidine-1-carbonyl)-phenyl]-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-phenyl]-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(4-dimethylcarbamoyl-phenyl)-amide];

(2R,4R)-4-[[1-(4-Chloro-phenylcarbamoyl)-4-hydroxy-pyrrolidine-2-carbonyl]-amino]-3-dimethylamino-benzoic acid methyl ester;

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-phenyl)-amide];

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-phenyl]-amide];

30 (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(pyrrolidine-1-carbonyl)-phenyl]-amide];

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(pyrrolidine-1-carbonyl)-phenyl]-amide];

5 (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-2-fluoro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-methyl-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

10 (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-methyl-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-quinolin-8-yl-phenyl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3,5-difluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

15 Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-2-methyl-biphenyl-4-yl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-biphenyl-4-yl)-amide];

20 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-biphenyl-4-yl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-methyl-2'-methylsulfonyl-biphenyl-4-yl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

25 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3-methyl-biphenyl-4-yl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3-trifluoromethyl-biphenyl-4-yl)-amide];

30 5-Methyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

3-Hydroxymethyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
5-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

(R) 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 5-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

5 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
5-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
5-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

1-(4-Chloro-phenylcarbamoyl)-5-(3-fluoro-2'-methanesulfonyl-biphenyl-
10 4-ylcarbamoyl)-4,5-dihydro-1H-pyrazole-3-carboxylic acid ethyl ester;

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[(2'-methoxy-biphenyl-4-yl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[(2'-hydroxy-biphenyl-4-yl)-amide];

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[(2-fluoro-4-iodo-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[[2-fluoro-4-(3-hydroxy-2-oxo-piperidin-1-yl)-phenyl]-amide];

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-tetrahydro-pyrimidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-imidazolidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-oxazolidin-3-yl)-phenyl]-amide];

25 (2R,4R)-1-(4-[[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-
carbonyl]-amino]-3-fluoro-phenyl)-2-oxo-piperidine-3-carboxylic acid ethyl
ester;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[[2-fluoro-4-(3-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl]-

30 amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(3-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(3-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-morpholin-4-yl-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2-methyl-5-oxo-pyrrolidin-1-yl)-phenyl]-amide];

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(2-methyl-5-oxo-pyrrolidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(4-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(4-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(5-chloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

20 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(4-tert-butyl-phenyl)-amide] 1-[(4-chloro-phenyl)-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2[(3,5'-difluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

25 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)-amide];

4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide];

4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(2,3-dimethyl-5-oxo-2,5-dihydro-pyrazol-1-yl)-phenyl]-amide];

30 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,3-dimethyl-5-oxo-2,5-dihydro-pyrazol-1-yl)-phenyl]-amide];

-41-

- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide]-TFA Salt;
- 5 4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide]; or
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 10 2-{[2-fluoro-4-(6-oxo-6H-pyridazin-1-yl)-phenyl]-amide};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(6-methyl-5-oxo-5H-[1,2,4]triazin-4-yl)-phenyl]-amide};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(3-chloro-5-methyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];
- 15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(3-methyl-5-oxo-4,5-dihydro-pyrazol-1-yl)-phenyl]-amide};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(5-chloro-3-methyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];
- 20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(3,5-dimethoxy-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(5-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};
- 25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-oxo-3-pyrrol-1-yl-2H-pyridin-1-yl)-phenyl]-amide};
- 30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-oxo-3-phenyl-2H-pyridin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-methyl-2-oxo-pyrrolidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-methyl-2-oxo-imidazolidin-1-yl)-phenyl]-

5 amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3,5-dimethyl-2-oxo-imidazolidin-1-yl)-2-fluoro-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3,5,5-trimethyl-2-oxo-imidazolidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-isopropyl-5,5-dimethyl-2-oxo-imidazolidin-1-yl)-phenyl]-amide];

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-methyl-2-oxo-imidazolidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(5,5-dimethyl-2-oxo-imidazolidin-1-yl)-2-fluoro-phenyl]-

20 amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3,4-dimethyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(4-chloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(3-methyl-2-oxo-pyrrolidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(3-methyl-2-oxo-imidazolidin-1-yl)-phenyl]-amide];

30

-43-

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(3,5-dimethyl-2-oxo-imidazolidin-1-yl)-2-fluorophenyl]-amide];

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(3,5,5-trimethyl-2-oxo-imidazolidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(3-isopropyl-5,5-dimethyl-2-oxo-imidazolidin-1-yl)-phenyl]-amide];

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(5-methyl-2-oxo-imidazolidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(5,5-dimethyl-2-oxo-imidazolidin-1-yl)-2-fluorophenyl]-amide];

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(3,4-dimethyl-2-oxo-2H-pyridin-1-yl)-2-fluorophenyl]-amide];

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(4-chloro-2-oxo-2H-pyridin-1-yl)-2-fluorophenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];
or a pharmaceutically acceptable salt thereof.

The invention is also directed to a compound which is:

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chlorophenyl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-quinolin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chlorophenyl)-amide] 2-[[2-fluoro-4-(3-oxo-3H-isoquinolin-2-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chlorophenyl)-amide] 2-[[2-fluoro-4-(1-oxo-1H-isoquinolin-2-yl)-phenyl]-amide];

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chlorophenyl)-amide] 2-[[2-fluoro-4-(5-oxo-5H-[1,2,4]triazin-4-yl)-phenyl]-amide];

- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(6-oxo-6H-pyrimidin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyrazin-1-yl)-phenyl]-amide];
- 5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyrimidin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-[1,3,5]triazin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(6-methyl-2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(1-methyl-1H-imidazol-2-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(1-methyl-1H-pyrrol-2-yl)-phenyl]-amide];
- 15 (2R,4R)-5-Aza-spiro[2.4]heptane-5,6-dicarboxylic acid 5-[(4-chloro-phenyl)-amide] 6-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyridine-1-carbonyl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-methyl-3H-imidazol-4-yl)-phenyl]-amide];
- 20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-methyl-2H-pyrazol-3-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[5-(2,5-dihydro-pyrrole-1-carbonyl)-pyridin-2-yl]-amide];
- 25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-methyl-5-oxo-2,5-dihydro-pyrazol-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-methoxymethyl-3-methyl-pyrazol-1-yl)-phenyl]-amide];
- 30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-methoxymethyl-5-methyl-pyrazol-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-ethyl-3-methyl-pyrazol-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-isobutyl-3-methyl-pyrazol-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-isopropyl-pyrazol-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-ethyl-pyrazol-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-isobutyl-pyrazol-1-yl)-phenyl]-amide];

(2R,4R)-2-(4-[[1-(4-Chloro-phenylcarbonyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-2H-pyrazole-3-carboxylic acid methyl ester;

(2R,4R)-2-(4-[[1-(4-Chloro-phenylcarbonyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-2H-pyrazole-3-carboxylic acid;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-methoxymethyl-pyrazol-1-yl)-phenyl]-amide];

(2R,4R)-5-(4-[[1-(4-Chloro-phenylcarbonyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-oxazole-4-carboxylic acid ethyl ester;

(2R,4R)-5-(4-[[1-(4-Chloro-phenylcarbonyl)-4-hydroxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-oxazole-4-carboxylic acid ethyl ester;

(2R,4R)-5-(4-[[1-(4-Chloro-phenylcarbonyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-oxazole-4-carboxylic acid;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(4-methoxymethyl-oxazol-5-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(4-ethyl-oxazol-5-yl)-2-fluoro-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(4-isopropyl-oxazol-5-yl)-phenyl]-amide];

(2R,4R)-5-(6-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-pyridin-3-yl)-oxazole-4-carboxylic acid ethyl ester;

(2R,4R)-5-(6-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-pyridin-3-yl)-oxazole-4-carboxylic acid;

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[5-(4-methoxymethyl-oxazol-5-yl)-pyridin-2-yl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[5-(4-ethyl-oxazol-5-yl)-pyridin-2-yl]-amide};

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[5-(4-vinyl-oxazol-5-yl)-pyridin-2-yl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[5-(4-isobutyl-oxazol-5-yl)-pyridin-2-yl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(4-methyl-isoxazol-3-yl)-phenyl]-amide};

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(4-methyl-isoxazol-3-yl)-phenyl]-amide};

(2R,4R)-3-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-isoxazole-4-carboxylic acid methyl ester;

20 (2R,4R)-3-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-isoxazole-4-carboxylic acid;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(4-methoxymethyl-isoxazol-3-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(4-isopropyl-isoxazol-3-yl)-phenyl]-amide};

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(4-cyclopropyl-isoxazol-3-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(4-cyclopropyl-isoxazol-3-yl)-phenyl]-amide};

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-methyl-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-methyl-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-methyl-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[(4-chloro-phenyl)-amide] 1-[[2,6-difluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[(5-chloro-pyridin-2-yl)-amide] 1-[[2,6-difluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 2-[(4-chloro-phenyl)-amide] 1-[[2,6-difluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

10 (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 2-[(5-chloro-pyridin-2-yl)-amide] 1-[[2,6-difluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(5'-methyl-2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(5'-methyl-2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];

15 (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(5'-methyl-2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(5'-methyl-2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methyl-2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(2'-methyl-2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methyl-2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];

25 (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(2'-methyl-2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(ethyl-isopropyl-carbamoyl)-phenyl]-amide];

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(ethyl-isopropyl-carbamoyl)-2-fluoro-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3-ethyl-pyrrolidine-1-carbonyl)-phenyl]-amide];

- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(3-ethyl-pyrrolidine-1-carbonyl)-2-fluoro-phenyl]-amide};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(3R-amino-pyrrolidine-1-carbonyl)-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];
- 5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(3R-amino-pyrrolidine-1-carbonyl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(3S-amino-pyrrolidine-1-carbonyl)-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(3S-amino-pyrrolidine-1-carbonyl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];
- 10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(3R-methylamino-pyrrolidine-1-carbonyl)-phenyl]-amide};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(3R-methylamino-pyrrolidine-1-carbonyl)-phenyl]-amide};
- 15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(3S-methylamino-pyrrolidine-1-carbonyl)-phenyl]-amide};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(3S-methylamino-pyrrolidine-1-carbonyl)-phenyl]-amide};
- 20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(3R-ethylamino-pyrrolidine-1-carbonyl)-phenyl]-amide};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(3R-ethylamino-pyrrolidine-1-carbonyl)-phenyl]-amide};
- 25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(3S-ethylamino-pyrrolidine-1-carbonyl)-phenyl]-amide};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(3S-ethylamino-pyrrolidine-1-carbonyl)-phenyl]-amide};
- 30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[4-(2,5-dihydro-pyrrole-1-carbonyl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,5-dihydro-pyrrole-1-carbonyl)-2-fluoro-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(2,5-dihydro-pyrrole-1-carbonyl)-2-fluoro-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(pyrrole-1-carbonyl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(pyrrole-1-carbonyl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(pyrrole-1-carbonyl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(pyrrole-1-carbonyl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-fluoro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-1-(4-[[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(6-hydroxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-1-(4-[[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid;

(2R,4R)-1-(4-[[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-6-oxo-1,6-dihydro-pyridine-2-carboxylic acid methyl ester;

(2R,4R)-[1-(4-[[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-phenyl)-2,5-dioxo-imidazolidin-4-yl]-acetic acid;

(2R,4R)-1-(4-[[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid methyl ester;

-50-

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-3-trifluoromethyl-2H-pyridin-1-yl)-phenyl]-amide];

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(4-hydroxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-1-(4-[[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-6-oxo-1,6-dihydro-pyridine-2-carboxylic acid;

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-5-trifluoromethyl-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(6-cyclopropyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide];

15 (2R,4R)-1-(4-[[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid ethyl ester;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3,3-dimethyl-2-oxo-azetidin-1-yl)-phenyl]-amide];

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(7-oxo-7H-thieno[2,3-c]pyridin-6-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-1-aza-spiro[4.5]dec-1-yl)-phenyl]-amide];

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-benzoyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-acetylsulfamoyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-thiazolidin-3-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-methyl-2-oxo-piperidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-3-sulfamoyl-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,5-dioxo-2,5-dihydro-pyrrol-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-amino-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-3-phenoxy-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,2-dimethyl-5-oxo-pyrrolidin-1-yl)-2-fluoro-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-nitro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-1-(4-[[1-(4-Chloro-phenylcarbonyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid tert-butyl ester;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,5-dioxo-pyrrolidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-benzothiazol-2-yl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-amino-4,6-dimethyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3-diethylaminomethyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(3-aminomethyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide} 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(3-methyl-5-nitro-2-oxo-2H-pyridin-1-yl)-phenyl}-amide};

(2R,4R)-Acetic acid 1-(4-{{1-(4-chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl}-amino}-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridin-3-yl ester;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{4-[3-(cyclopentanecarbonyl-amino)-2-oxo-2H-pyridin-1-yl]-2-fluoro-phenyl}-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(5-amino-3-methyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide} 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-[2-oxo-3-(pyridin-4-ylcarbamoyl)-2H-pyridin-1-yl]-phenyl}-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{4-(3,5-dichloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(3-amino-5-bromo-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide} 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{4-(3-cyano-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{4-(3-cyano-2-oxo-5-phenyl-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{4-(3-ethyl-6-methyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3-cyano-4-dimethylamino-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide];

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(4-methyl-3-nitro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-nitro-2-oxo-3-trifluoromethyl-2H-pyridin-1-yl)-phenyl]-amide];

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(5-amino-6-oxo-6H-[3,4']bipyridinyl-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(5-chloro-3-nitro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

15

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-carbamoyl-2-oxo-5-trifluoromethyl-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

(2R,4R)-5-Bromo-1-(4-[[1-(4-chloro-phenyl)carbamoyl]-4-methoxy-pyrrolidine-2-carbonyl]-amino)-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid isobutyl ester;

20

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-chloro-2-oxo-5-trifluoromethyl-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-bromo-5-chloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(5-bromo-3-nitro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

30

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(3-nitro-2-oxo-5-trifluoromethyl-2H-pyridin-1-yl)-phenyl}-amide};

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(5-methyl-3-nitro-2-oxo-2H-pyridin-1-yl)-phenyl}-amide};

(2R,4R)-5-Chloro-1-(4-{{1-(4-chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl}-amino}-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid methyl ester;

10 (2R,4R)-5-Chloro-1-(4-{{1-(4-chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl}-amino}-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(4-benzyloxy-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide} 1-[(4-chloro-phenyl)-amide];

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(4-amino-5-fluoro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide} 1-[(4-chloro-phenyl)-amide];

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(6-methyl-3-oxo-3H-isoquinolin-2-yl)-phenyl}-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(4-methyl-5-nitro-2-oxo-2H-pyridin-1-yl)-phenyl}-amide};

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(5-acetylamino-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide} 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(4-amino-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide} 1-[(4-chloro-phenyl)-amide];

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(6-oxo-6H-[3,4']bipyridinyl-1-yl)-phenyl}-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(4-bromo-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide} 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(4-oxo-4H-furo[3,2-c]pyridin-5-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(1-oxo-1H-[2,6]naphthyridin-2-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(5-bromo-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(4-oxo-4H-thieno[3,2-c]pyridin-5-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(4-oxo-1,4-dihydro-imidazo[4,5-c]pyridin-5-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(5-chloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(5-fluoro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-1-(4-[[1-(5-Chloro-pyridin-2-yl)carbamoyl]-4-methoxy-pyrrolidine-2-carbonyl]-amino)-3-fluoro-phenyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(6-hydroxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-1-(4-[[1-(5-Chloro-pyridin-2-yl)carbamoyl]-4-methoxy-pyrrolidine-2-carbonyl]-amino)-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid;

(2R,4R)-1-(4-[[1-(5-Chloro-pyridin-2-yl)carbamoyl]-4-methoxy-pyrrolidine-2-carbonyl]-amino)-3-fluoro-phenyl)-6-oxo-1,6-dihydro-pyridine-2-carboxylic acid methyl ester;

(2R,4R)-[1-(4-{{1-(5-Chloro-pyridin-2-ylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-phenyl)-2,5-dioxo-imidazolidin-4-yl]-acetic acid;

(2R,4R)-1-(4-{{1-(5-Chloro-pyridin-2-ylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid methyl ester;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{2-fluoro-4-(2-oxo-3-trifluoromethyl-2H-pyridin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{2-fluoro-4-(4-hydroxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

(2R,4R)-1-(4-{{1-(5-Chloro-pyridin-2-ylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-6-oxo-1,6-dihydro-pyridine-2-carboxylic acid;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{2-fluoro-4-(2-oxo-5-trifluoromethyl-2H-pyridin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{4-(6-cyclopropyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide};

(2R,4R)-1-(4-{{1-(5-Chloro-pyridin-2-ylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid ethyl ester;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{2-fluoro-4-(3-hydroxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{4-(3,3-dimethyl-2-oxo-azetidin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{2-fluoro-4-(7-oxo-7H-thieno[2,3-c]pyridin-6-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-1-aza-spiro[4.5]dec-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-benzoyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-methyl-5-oxo-pyrrolidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-methyl-6-oxo-piperidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-acetylsulfamoyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-thiazolidin-3-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(3-methyl-2-oxo-piperidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-3-sulfamoyl-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(2,5-dioxo-2,5-dihydro-pyrrol-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-amino-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-3-phenoxy-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(2,2-dimethyl-5-oxo-pyrrolidin-1-yl)-2-fluoro-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(3-nitro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

5 (2R,4R)-1-(4-[[1-(5-Chloro-pyridin-2-ylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid tert-butyl ester;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(2,5-dioxo-pyrrolidin-1-yl)-phenyl]-amide];

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-benzothiazol-2-yl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-amino-4,6-dimethyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(3-diethylaminomethyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-aminomethyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

20

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(3-methyl-5-nitro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-Acetic acid 1-(4-[[1-(5-chloro-pyridin-2-ylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridin-3-yl ester;

25

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-[3-(cyclopentanecarbonyl-amino)-2-oxo-2H-pyridin-1-yl]-2-fluoro-phenyl]-amide];

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(5-amino-3-methyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-([2-fluoro-4-[2-oxo-3-(pyridin-4-ylcarbamoyl)-2H-pyridin-1-yl]-phenyl]-amide);

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[4-(3,5-dichloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(3-amino-5-bromo-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[4-(3-cyano-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[4-(3-cyano-2-oxo-5-phenyl-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[4-(3-ethyl-6-methyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[4-(3-cyano-4-dimethylamino-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(4-methyl-3-nitro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(5-nitro-2-oxo-3-trifluoromethyl-2H-pyridin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(5-amino-6-oxo-6H-[3,4']bipyridinyl-1-yl)-2-fluoro-phenyl]-amide} 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(5-chloro-3-nitro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(5-chloro-pyridin-2-yl)-amide];

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(3-carbamoyl-2-oxo-5-trifluoromethyl-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-5-Bromo-1-(4-{{1-(5-chloro-pyridin-2-ylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid isobutyl ester;

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(3-chloro-2-oxo-5-trifluoromethyl-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(3-bromo-5-chloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(5-chloro-pyridin-2-yl)-amide];

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(5-bromo-3-nitro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{2-fluoro-4-(3-nitro-2-oxo-5-trifluoromethyl-2H-pyridin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{2-fluoro-4-(5-methyl-3-nitro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

25 (2R,4R)-5-Chloro-1-(4-{{1-(5-chloro-pyridin-2-ylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid methyl ester;

(2R,4R)-5-Chloro-1-(4-{{1-(5-chloro-pyridin-2-ylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid;

30

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-([4-(4-benzyloxy-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide) 1-[(5-chloro-pyridin-2-yl)-amide];

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-([4-(4-amino-5-fluoro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide) 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-([2-fluoro-4-(5-nitro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide);

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-([2-fluoro-4-(6-methyl-3-oxo-3H-isoquinolin-2-yl)-phenyl]-amide);

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-([2-fluoro-4-(4-methyl-5-nitro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide);

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-([4-(5-acetylamino-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide) 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-([2-fluoro-4-(1-oxo-1H-isoquinolin-2-yl)-phenyl]-amide);

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-([4-(4-amino-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide) 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-([2-fluoro-4-(6-oxo-6H-[3,4']bipyridinyl-1-yl)-phenyl]-amide);

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-([2-fluoro-4-(3-oxo-3H-isoquinolin-2-yl)-phenyl]-amide);

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-([4-(4-bromo-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide) 1-[(5-chloro-pyridin-2-yl)-amide];

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-([2-fluoro-4-(4-oxo-4H-furo[3,2-c]pyridin-5-yl)-phenyl]-amide);

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(1-oxo-1H-[2,6]naphthyridin-2-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(5-bromo-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(4-oxo-4H-thieno[3,2-c]pyridin-5-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(4-oxo-1,4-dihydro-imidazo[4,5-c]pyridin-5-yl)-phenyl]-amide];

4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

3-Hydroxymethyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

5-Methyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

3-Methoxymethyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 5-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

3-Hydroxymethyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 5-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

5-Methyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 5-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

3-Methoxymethyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 5-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[[4-(2-methyl-thiazol-4-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-2-methyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2- {[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3,4,5,6-tetrahydro-2H-[1,4']bipyridinyl-4-ylmethyl)-amide];

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2- {[4-(2-dimethylaminomethyl-imidazol-1-yl)-2-fluoro-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2- {[2-chloro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2- {[2-chloro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide} 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2- {[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2- {[2-fluoro-4-(2-methylamino-imidazol-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2- {[2-fluoro-4-(2-isopropyl-imidazol-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2- {[4-(2-ethyl-imidazol-1-yl)-2-fluoro-phenyl]-amide};

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2- {[4-(2-ethyl-imidazol-1-yl)-2-fluoro-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2- {[2-fluoro-4-(2-isopropyl-imidazol-1-yl)-phenyl]-amide};

25 (2R,4R)-2-Ethyl-4-methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2- {[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide};

4-Methoxy-2-methoxymethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2- {[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide};

2-Hydroxymethyl-4-methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2- {[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide};

30 {1-(4-Chloro-phenylcarbamoyl)-2-[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenylcarbamoyl]-4-methoxy-pyrrolidin-2-yl}-acetic acid methyl ester;

- { 1-(4-Chloro-phenylcarbamoyl)-2-[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenylcarbamoyl]-4-methoxy-pyrrolidin-2-yl}-acetic acid;
- 4-Hydroxymethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 5 { 1-(4-Chloro-phenylcarbamoyl)-5-[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenylcarbamoyl]-pyrrolidin-3-yl}-acetic acid;
- { 1-(4-Chloro-phenylcarbamoyl)-5-[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenylcarbamoyl]-pyrrolidin-3-yl}-acetic acid methyl ester;
- (2R,4R)-4-Ethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- 10 (2R,4R)-4-Isopropyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-fluoro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- 15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(3-fluoro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-chloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];
- 20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-chloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-bromo-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-bromo-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];
- 25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-iodo-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(3-iodo-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- 30 amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(3-ethyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide]; and

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3-ethyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide;
5 a pharmaceutically acceptable salt thereof.

The invention is also directed to a compound which is:

(2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
10 (2R,4R) 4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
(2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
15 (2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
(2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];
(2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
20 (2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
(2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-methyl-imidazol-1-yl)-phenyl]-amide];
25 (2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-2-fluoro-phenyl)-amide];
(2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-methyl-pyrazol-1-yl)-phenyl]-amide];
30 (2R,4R) 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

- (2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-pyrazol-1-yl)-phenyl]-amide];
- (2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(pyrrolidine-1-carbonyl)-phenyl]-amide];
- 5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-
10 pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-methyl-imidazol-1-yl)-phenyl]-amide];
- 15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];
- (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
20 phenyl)-amide] 2-[(4-dimethylcarbamoyl-2-fluoro-phenyl)-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-methyl-pyrazol-1-yl)-phenyl]-amide];
- 5-Methyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
- 25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(4-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[2-chloro-4-(2-
30 oxo-2H-pyridin-1-yl)-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-methyl-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(6-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(3-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(5-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(4-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,5-dihydro-pyrrole-1-carbonyl)-2-fluoro-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,5-dihydro-pyrrole-1-carbonyl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2-methyl-5-oxo-pyrrolidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,5-dihydro-pyrrole-1-carbonyl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(4-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,5-dihydro-pyrrole-1-carbonyl)-2-fluoro-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2-methyl-5-oxo-pyrrolidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,5-dihydro-pyrrole-1-carbonyl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(4-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

-68-

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,5-dihydro-pyrrole-1-carbonyl)-2-fluoro-phenyl]-amide}
or a pharmaceutically acceptable salt thereof.


- 5 The invention is also directed to a pharmaceutical composition comprising a compound of formula I admixed with a carrier, diluent, or excipient.

- The invention is also directed to a method for preventing or treating acute, subacute, and chronic thrombotic disorder in a mammal is comprising
10 administering to a mammal in need thereof a therapeutically effective amount of a compound of any of formulas I-X.

- The invention is also directed to a method of inhibiting Factor Xa in a mammal, comprising administering to a mammal in need of Factor Xa inhibition a
15 Factor Xa inhibition amount of a compound of any of formulas I-X.


 The invention is also directed to a process for preparing a compound of Formula I comprising:

- (a) reacting an amino acid of formula IA



20

with a reagent capable of placing a protecting group P¹ on the amino group of an amino acid to form a protected amino acid given by formula IB:




-69-

wherein P^1 is a protecting group and W^1 is the same as defined above;

(b) converting the acid moiety in the compound of formula IB to a acid
5 halide;


(c) reacting the acid halide with a haloaniline or a haloamino-
heterocycle to form a compound having formula IC:



10

wherein Z^1 is a halogen and A is as defined above;

(d) coupling the compound of formula IC with a compound H-B to
15 give a compound having formula ID:



(e) removing the protecting group from Compound ID and reacting
20 the resulting acid moiety with a C-isocyanate to form a compound of the
invention.

-70-

DETAILED DESCRIPTION OF THE INVENTION

The following definitions are used, unless otherwise described: halo is fluoro, chloro, bromo, or iodo. Alkyl, alkoxy, alkenyl, alkynyl, etc. denote both straight and branched groups; but reference to an individual radical such as

- 5 "propyl" embraces only the straight chain radical, a branched chain isomer such as "isopropyl" being specifically referred to.

The term "alkyl" as used herein refers to a straight or branched hydrocarbon of from 1 to 11 carbon atoms and includes, for example, methyl,
 10 ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, *tert*-butyl, n-pentyl, n-hexyl, and the like. The alkyl group can also be substituted with one or more of the substituents selected from lower alkoxy, lower thioalkoxy, halogen, nitro, cyano, oxo, thio, -OH, -SH, -F, -CF₃, -OCF₃, -NO₂, -CO₂H, -CO₂(C₁-C₆)alkyl,

-NH₂, -NH(C₁-C₆)alkyl, , -CONR⁸R⁹, or -N(C₁-C₆)alkyl)₂. Preferred

- 15 alkyl groups have from 1 to 6 carbon atoms (C₁-C₆ alkyl).

The term "(C₁-C₆)alkyl" as used herein refers to a subset of alkyl which means a straight or branched hydrocarbon radical having from 1 to 6 carbon atoms and includes, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl,
 20 isobutyl, *tert*-butyl, n-pentyl, n-hexyl, and the like. Sometimes herein lower alkyl is referred to as "C₁-C₆alkyl."


The term "(C₁-C₆)hydroxyalkyl" or "hydroxy(C₁-C₆)alkyl" as used herein means a straight or branched alcohol having from 1 to 6 carbon atoms and
 25 includes, for example, hydroxymethyl, hydroxyethyl, hydroxy-propyl and the like.

The term "(C₂-C₆)alkenyl" means a straight or branched unsaturated hydrocarbon radical having from 2 to 6 carbon atoms and includes, for example, ethenyl, 2-propenyl, 1-butenyl, 2-butenyl, 1-pentenyl, 2-pentenyl, 3-methyl-3-

-71-

butenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 3-heptenyl, 1-octenyl, 1-nonenyl, 1-decenyl, 1-undecenyl, 1-dodecenyl, and the like.

The term "alkylene" as used herein refers to a divalent group derived from
 5 a straight or branched chain saturated hydrocarbon having from 1 to 10 carbon atoms by the removal of two hydrogen atoms, for example methylene, 1,2-ethylene, 1,1-ethylene, 1,3-propylene, 2,2-dimethylpropylene, and the like. The alkylene groups of this invention can be optionally substituted. The alkylene group can also be substituted with one or more of the substituents selected from
 10 lower alkyl, lower alkoxy, lower thioalkoxy, halogen, nitro, cyano, oxo, thio, -OH, -SH, -F, -CF₃, -OCF₃, -NO₂, -CO₂H, -CO₂C₁-C₆ alkyl, -NH₂, -NH(C₁-

C₆)alkyl, , -CONR⁸R⁹, or -N(C₁-C₆alkyl)₂. Preferred alkylene groups have from 1 to 6 carbon atoms (C₁-C₆ alkyl).


15 The term "heteroatom" as used herein represents oxygen, nitrogen, or sulfur (O, N, or S) as well as sulfoxyl or sulfonyl (SO or SO₂), unless otherwise indicated. It is to be understood that alkyl chains interrupted by one or more heteroatoms means that a carbon atom of the chain is replaced with a heteroatom having the appropriate valency. Preferably, an alkyl chain is interrupted by 1 to 4
 20 heteroatoms and that two adjacent carbon atoms are not both replaced. Examples of such groups include methoxymethyl, 3-thiomethylpropyl, and 2-thiomethoxyethoxymethyl.

The term "hetero(C₁-C₆)alkyl" as used herein, refers to an alkyl group that
 25 includes one or more heteroatoms such as oxygen, sulfur, or nitrogen (with valence completed by hydrogen or oxygen) in the carbon chain or terminating the carbon chain.

The term "hydrocarbon chain" as used herein refers to a straight
 30 hydrocarbon of from 2 to 6 carbon atoms. The hydrocarbon chain is optionally


-72-

substituted with one or more substituents selected from lower alkyl, lower alkoxy, lower thioalkoxy, halogen, nitro, cyano, oxo, thio, -OH, -SH, -F, -CF₃, -OCF₃,

-NO₂, -CO₂H, -CO₂(C₁-C₆)alkyl, -NH₂, -NH(C₁-C₆)alkyl, ,
-CONR⁸R⁹, or -N(C₁-C₆alkyl)₂.


5

The term "hydrocarbon-heteroatom chain" as used herein refers to a hydrocarbon chain wherein one or more carbon atoms are replaced with a heteroatom. The hydrocarbon-heteroatom chain is optionally substituted with one or more substituents selected from lower alkyl, lower alkoxy, lower thioalkoxy,

10 halogen, nitro, cyano, oxo, thio, , -OH, -SH, -F, -CF₃, -OCF₃, -NO₂,
-CO₂H, -CO₂(C₁-C₆)alkyl, -NH₂, -NH(C₁-C₆)alkyl, -CONR⁸R⁹, or -N(C₁-C₆alkyl)₂.

15 The term "heteroalkylene" as used herein, refers to an alkylene radical as defined above that includes one or more heteroatoms such as oxygen, sulfur, or nitrogen (with valence completed by hydrogen or oxygen) in the carbon chain or terminating the carbon chain.

20 The term "(C₃-C₇)cycloalkyl" means a cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, or cycloheptyl ring, respectively. The (C₃-C₇)cycloalkyl ring may be optionally substituted with one or more substituents selected from lower alkyl, lower alkoxy, lower thioalkoxy, halogen, nitro, cyano, oxo, thio,

, -OH, -SH, -F, -CF₃, -OCF₃, -NO₂, -CO₂H, -CO₂(C₁-C₆)alkyl, -NH₂,
-NH(C₁-C₆)alkyl, -CONR⁸R⁹, or -N(C₁-C₆alkyl)₂.

25

The term "halo(C₁-C₆)alkyl" as used herein means a lower alkyl radical, as defined above, bearing at least one halogen substituent, for example,

-73-

chloromethyl, fluoroethyl, or trifluoromethyl, and the like. Haloalkyl can also include perfluoroalkyl wherein all hydrogens of a loweralkyl group are replaced with fluorides.

5 The term "heterocycle" or "heterocyclo" means a saturated or unsaturated mono- or polycyclic (i.e. bicyclic) fused, bridged, or spiro bicyclic heterocyclic ring system ring incorporating one or more (i.e. 1-4) heteroatoms selected from N, O, and S. It is to be understood that a heterocycle is optionally substituted with -OH, -O(alkyl), SH, S(alkyl), amine, halogen, acid, ester, amide, amidine, alkyl
10 ketone, aldehyde, nitrile, fluoroalkyl, nitro, sulphone, sulfoxide or Cl-6 alkyl. Monocyclic heterocyclic rings contain from about 3 to 12 ring atoms, with from 1 to 5 heteroatoms selected from N, O, and S, and preferably from 3 to 7 member atoms, in the ring. Bicyclic heterocyclics contain from 7 to 17 member atoms, preferably 7 to 12 member atoms, in the ring. Bicyclic heterocyclics contain from
15 about 7 to about 17 ring atoms, preferably from 7 to 12 ring atoms. Bicyclic heterocyclics rings may be fused, spiro, or bridged ring systems. Any of the heterocycles may be substituted with up to three groups as selected from the substitutions provided for alkyl.

20 Examples of heterocyclic groups include cyclic ethers (oxiranes) such as ethyleneoxide, tetrahydrofuran, dioxane, and substituted cyclic ethers, wherein the substituents are those described above for the alkyl and cycloalkyl groups. Typical substituted cyclic ethers include propyleneoxide, phenyloxirane (styrene oxide), cis-2-butene-oxide (2,3-dimethyloxirane), 3-chlorotetrahydrofuran, 2,6-dimethyl-
25 1,4-dioxane, and the like. Heterocyclo groups containing nitrogen are groups such as pyrrolidine, piperidine, pyridin-2-one, piperazine, tetrahydrotriazine, tetrahydropyrazole, and substituted groups such as 3-aminopyrrolidine, 4-methylpiperazin-1-yl, and the like. Typical sulfur containing heterocycles include tetrahydrothiophene, dihydro-1,3-dithiol-2-yl, and hexahydrothiepin-4-yl. Other
30 commonly employed heterocycles include dihydro-oxathiol-4-yl, tetrahydro-oxazolyl, tetrahydro-oxadiazolyl, tetrahydro-dioxazolyl, tetrahydro-oxathiazolyl, hexahydrotriazinyl, tetrahydro-oxazinyl, morpholinyl, thiomorpholinyl,


-74-

tetrahydropyrimidinyl, dioxolinyl, octahydrobenzofuranyl, octahydrobenzimidazolyl, and octahydrobenzothiazolyl. For heterocycles containing sulfur, the oxidized sulfur heterocycles containing SO or SO₂ groups are also included. Examples include the sulfoxide and sulfone forms of tetrahydrothiophene. Examples of unsaturated heterocyclic groups include

5 pyridones (1-H pyridin-2-ones), pyrimidones (1H pyrimidin-2-one, 1H pyrimidin-4-one), pyrazones (1H pyrazin-2-ones), pyridazones (1H pyridazin-3-one), dihydropyrroles, dihydro- and tetrahydro pyridines and the like, as well as any unsaturated heterocyclic groups specifically disclosed herein.

10

The term "aryl" as used herein refers to an aromatic ring which is unsubstituted or optionally substituted by 1 to 4 substituents selected from lower alkyl, lower alkoxy, lower thioalkoxy, halogen, nitro, cyano -OH, -SH, -F, -CF₃,

-OCF₃, -NO₂,  -CO₂H, -CO₂(C₁-C₆)alkyl, -NH₂, -NH(C₁-C₆)alkyl,

15 -CONR⁸R⁹, -, SO₂alkyl, -SO₂NH₂, or -N(C₁-C₆alkyl)₂. Examples include, but are not limited to phenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 2-chloro-3-methylphenyl, 2-chloro-4-methylphenyl, 2-chloro-5-methylphenyl, 3-chloro-2-methylphenyl, 3-chloro-4-methylphenyl, 4-chloro-2-methylphenyl, 4-chloro-3-methylphenyl, 5-chloro-2-methylphenyl, 2,3-dichlorophenyl, 2,5-dichlorophenyl, 3,4-dichlorophenyl, 2,3-dimethylphenyl, 3,4-dimethylphenyl, thienyl, furanyl, pyrrolyl, pyridyl, pyrimidyl,

20 imidazolyl, pyrazinyl, oxazolyl, thiazolyl, naphthyl, benzothienyl, benzofuranyl, indolyl, quinolinyl, isoquinolinyl, and quinazolinyl, and the like.

25

The term "(C₄-C₇)heterocycloalkenyl" refers to a subset of unsaturated heterocyclo, examples of which are provided above.

The term "aralkyl" as used herein refers to an alkyl radical to which is

30 appended an aryl group. Representative arylalkyl groups include benzyl,

-75-

phenylethyl, hydroxybenzyl, fluorobenzyl, fluorophenylethyl, and the like. The arylalkyl groups of this invention can be optionally substituted.

The term "arylene" as used herein refers to a divalent group derived from an aromatic ring. The arylene group can also be substituted with one or more of the substituents listed above for aryl.


The term "heteroaryl" means an aromatic cyclic or fused polycyclic ring system having from 1 to 8 heteroatoms selected from N, O, and S. The heteroaryl groups or fused heteroaryl groups may be unsubstituted or substituted by 1 to 3 substituents selected from those described above for alkyl, alkenyl, and alkynyl, for example, cyanothienyl and formylpyrrolyl.


Typical heteroaryl groups include 2- or 3-thienyl, 2- or 3-furyl, 2- or 3-pyrrolyl, 2-, 4-, or 5-imidazolyl, 3-, 4-, or 5-pyrazolyl, 2-, 4-, or 5-thiazolyl, 3-, 4-, or 5-isothiazolyl, 2-, 4-, or 5-oxazolyl, 3-, 4-, or 5-isoxazolyl, 3- or 5-1,2,4-triazolyl, 4- or 5-1,2,3-triazolyl, tetrazolyl, 2-, 3-, or 4-pyridyl, 3- or 4-pyridazinyl, 3-, 4-, or 5-pyrazinyl, 2-pyrazinyl, 2-, 4-, or 5-pyrimidinyl.

Aromatic fused heteroaryl groups of from 8 to 20 atoms include but are not limited to 1-, 2-, 3-, 5-, 6-, 7-, or 8-indoliziny, 1-, 3-, 4-, 5-, 6-, or 7-isoindolyl, 2-, 3-, 4-, 5-, 6-, or 7-indolyl, 2-, 3-, 4-, 5-, 6-, or 7-indazolyl, 2-, 4-, 5-, 6-, 7-, or 8-puriny, 1-, 2-, 3-, 4-, 6-, 7-, 8-, or 9-quinoliziny, 2-, 3-, 4-, 5-, 6-, 7-, or 8-quinoliyl, 1-, 3-, 4-, 5-, 6-, 7-, or 8-isoquinoliyl, 1-, 4-, 5-, 6-, 7-, or 8-phthalaziny, 2-, 3-, 4-, 5-, or 6-naphthyridiny, 2-, 3-, 5-, 6-, 7-, or 8-quinazoliny, 3-, 4-, 5-, 6-, 7-, or 8-cinnoliny, 2-, 4-, 6-, or 7-pteridiny, 1-, 2-, 3-, 4-, 5-, 6-, 7-, or 8-4aH carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, or 8-carbzaolyl, 1-, 3-, 4-, 5-, 6-, 7-, 8-, or 9-carboliny, 1-, 2-, 3-, 4-, 6-, 7-, 8-, 9-, or 10-phenanthridiny, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8-, or 9-acridiny, 1-, 2-, 4-, 5-, 6-, 7-, 8-, or 9-perimidiny, 2-, 3-, 4-, 5-, 6-, 8-, 9-, or 10-phenathroliny, 1-, 2-, 3-, 4-, 6-, 7-, 8-, or 9-phenaziny, 1-, 2-, 3-, 4-, 6-, 7-, 8-, 9-, or 10-phenothiaziny, 1-, 2-, 3-, 4-, 6-, 7-, 8-, 9-, or 10-phenoxaziny, 2—, 3—, 4—, 5—, 6—, or 1-, 3-, 4-, 5-, 6-, 7-, 8-, 9-, or 10-

-76-

benzisoquinoliny, 2-, 3-, 4-, or thieno[2,3-*b*]furanyl, 2-, 3-, 5-, 6-, 7-, 8-, 9-, 10-, or 11-7*H*-pyrazino[2,3-*c*]carbazolyl, 2-, 3-, 5-, 6-, or 7-2*H*-furo[3,2-*b*]-pyranyl, 2-, 3-, 4-, 5-, 7-, or 8-5*H*-pyrido[2,3-*d*]-*o*-oxazinyl, 1-, 3-, or 5-1*H*-pyrazolo[4,3-*d*]-oxazolyl, 2-, 4-, or 5-4*H*-imidazo[4,5-*d*]thiazolyl, 3-, 5-, or 8-pyrazino[2,3-*d*]pyridazinyl, 2-, 3-, 5-, or 6-imidazo[2,1-*b*]thiazolyl, 1-, 3-, 6-, 7-, 8-, or 9-furo[3,4-*c*]cinnoliny, 1-, 2-, 3-, 4-, 5-, 6-, 8-, 9-, 10, or 11-4*H*-pyrido[2,3-*c*]carbazolyl, 2-, 3-, 6-, or 7-imidazo[1,2-*b*][1,2,4]triazinyl, 7-benzo[*b*]thienyl, 2—, 4—, 5—, 6—, or 7-benzoxazolyl, 2—, 4—, 5—, 6—, or 7-benzimidazolyl, 2—, 4—, 5—, 6—, or 7-benzothiazolyl, 1-, 2-, 4-, 5-, 6-, 7-, 8-, or 9-benzoxapiny, 2-, 4-, 5-, 6-, 7-, or 8-benzoxazinyl, 1-, 2-, 3-, 5-, 6-, 7-, 8-, 9-, 10-, or 11-1*H*-pyrrolo[1,2-*b*][2]benzazapiny. Typical fused heteroaryl groups include, but are not limited to 2—, 3—, 4—, 5—, 6—, 7—, or 8-quinoliny, 1—, 3—, 4—, 5—, 6—, 7—, or 8-isoquinoliny, 2—, 3—, 4—, 5—, 6—, or 7-indolyl, 2—, 3—, 4—, 5—, 6—, or 7-benzo[*b*]thienyl, 2—, 4—, 5—, 6—, or 7-benzoxazolyl, 2—, 4—, 5—, 6—, or 7-benzimidazolyl, 2—, 4—, 5—, 6—, or 7-benzothiazolyl.

Other heteroaryl groups include the following groups wherein “” indicates the point of attachment, and any of which may be substituted as provided



20

The term “monocyclic heteroarylene” as used herein refers to a divalent group derived from a monocyclic heteroaryl. The aryene group can also be substituted with one or more of the substituents listed above for aryl.

25 The term “(C₄-C₇)cycloalkenyl” means a cycloalkyl group having one or more carbon-carbon double or triple bond. Example includes cyclobutene, cyclopentene, cyclohexene, cycloheptene, cyclobutadiene, cyclopentadiene, and the like.

-77-

The symbol "——" means a bond. If an asymmetric carbon is created by such a bond, a particular stereochemistry is not to be implied.

~

The symbol "====" means a double bond.

5

The symbol "~~~~" indicates covalent points of attachment.

The following abbreviations are used herein:

HOAc = Acetic acid

10

DCM = Dichloromethane

TFA = Trifluoroacetic acid

THF = tetrahydrofuran

Me = Methyl

APCI = Atmospheric pressure chemical ionization

15

Et = Ethyl

EtOAc = Ethyl acetate

MeOH = Methanol

HPLC = High Pressure Liquid Chromatography

MPLC = Medium Pressure Liquid Chromatography

20

EEDQ = N-Ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline

PyBOP = Benzotriazol-1-yloxytripyrrolidinophosphonium

hexafluorophosphate

When a bond to a substituent is shown to cross the bond connecting 2
25 atoms in a ring, then such substituent may be bonded to any atom in the ring, provided the atom will accept the substituent without violating its valency. When there appears to be several atoms of the substituent that may bond to the ring atom, then it is the first atom of the listed substituent that is attached to the ring.

30 When a bond from a substituent is shown to cross the bond connecting 2 atoms in a ring of the substituent, then such substituent may be bonded from any atom in the ring that is available.

-78-

When a bond is represented by a line such as "-----" this is meant to represent that the bond may be absent or present, provided that the resultant compound is stable and of satisfactory valency.

5

The term "patient" means all mammals including humans. Examples of patients include humans, cows, dogs, cats, goats, sheep, pigs, and rabbits.

A "therapeutically effective amount" is an amount of a compound of the present invention that when administered to a patient ameliorates a symptom of thrombotic disorders, venous thrombosis, arterial thrombosis, pulmonary embolism, myocardial infarction, cerebral infarction, restenosis, cancer, angina, diabetes, atrial fibrillation, or heart failure. A therapeutically effective amount of a compound of the present invention can be easily determined by one skilled in the art by administering a quantity of a compound to a patient and observing the result. In addition, those skilled in the art are familiar with identifying patients having thrombotic disorders, venous thrombosis, arterial thrombosis, pulmonary embolism, myocardial infarction, cerebral infarction, restenosis, cancer, angina, diabetes, atrial fibrillation, or heart failure.

20

The term "pharmaceutically acceptable salts, esters, amides, and prodrugs" as used herein refers to those carboxylate salts, amino acid addition salts, esters, amides, and prodrugs of the compounds of the present invention which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of patients without undue toxicity, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio, and effective for their intended use, as well as the zwitterionic forms, where possible, of the compounds of the invention. The term "salts" refers to the relatively non-toxic, inorganic and organic acid addition salts of compounds of the present invention. These salts can be prepared in situ during the final isolation and purification of the compounds or by separately reacting the purified compound in its free base form with a suitable organic or inorganic acid and isolating the salt thus formed.

30

-79-


In some situations, compounds of the invention may exist in isomeric form; for example, as tautomers, enantiomers, or diastereomers. Some compounds may exhibit polymorphism. All tautomers, enantiomers, and diastereomers are
 5 incorporated within the definition of the compounds of the invention. It is further to be understood that the present invention encompasses any racemic, optically-active, polymorphic, or stereoisomeric form, or mixtures thereof, of a compound of the invention, which possess the useful properties described herein, it being well known in the art how to prepare optically active forms (for example, by
 10 resolution of the racemic form by recrystallization techniques, by synthesis from optically-active starting materials, by chiral synthesis, or by chromatographic separation using a chiral stationary phase) and how to determine activity or cytotoxicity using the standard tests described herein, or using other similar tests which are well known in the art.

15

Certain compounds of the present invention can exist in unsolvated form as well as solvated form including hydrated form. In general, the solvated form including hydrated form is equivalent to unsolvated form and is intended to be encompassed within the scope of the present invention.

20

Referring now to a compound of Formula I,



I


A specific value for G is H, F, or methyl.

25 A specific value for W¹ is $\text{—}\overset{\text{O}}{\parallel}\text{—CH}_2\text{CH}_2\text{—}$. Other specific values for W¹ are $\text{—CH}_2\text{—CH}_2\text{—CH}_2\text{—}$, $\text{—CH}_2\text{—CH}_2\text{—O—CH}_2\text{—}$,

-80-

- $-\text{CH}_2-\text{HC}=\text{CH}-\text{CH}_2-$, $-\text{CH}_2-\text{CH}_2\text{CR}^5\text{R}^6-$, $-\text{CH}_2\text{CR}^5\text{R}^6\text{CH}_2-$,
 $-\text{CH}_2-\text{CH}_2\text{CR}^7\text{R}^8-$, $-\text{CH}_2\text{CR}^7\text{R}^8\text{CH}_2-$, $-\text{CH}_2-\text{C}=\text{R}^9\text{CH}_2-$,
 $-\text{CH}_2-\text{CH}_2\text{C}=\text{R}^9-$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2-\text{HC}=\text{CH}-$,
 $-\text{CH}_2\text{CH}_2\text{NR}^5-$, $-\text{CH}_2\text{NR}^5\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-\text{NH}-\text{CH}_2-$,
5 $-\text{CO}-\text{CHR}^5-\text{CH}_2-$, $-\text{CO}-\text{CH}_2-\text{CHR}^5-$, $-\text{N}=\text{CR}^5-\text{CH}=-$, $-\text{N}=\text{CR}^5-\text{CH}_2-$,
 $-\text{N}=\text{CH}-\text{CR}^5=$, $-\text{O}-\text{CHR}^5-\text{CH}=-$, $-\text{O}-\text{CHR}^5-\text{CH}_2-$, and $-\text{O}-\text{CH}_2-\text{CR}^5=$,
wherein R^5 is $-\text{OH}$, alkyl, halo(C_1-C_6)alkyl, $-\text{NR}^3\text{R}^4$, $-\text{OR}^2$, halo, $-\text{CN}$, $-\text{CH}_2\text{OR}^2$, $-\text{CH}_2-\text{NR}^3\text{R}^4$, aryl, monocyclic heteroaryl, alkylaryl, $-\text{CONR}^3\text{R}^4$, $-\text{COR}^2$, halo(C_1-C_6)alkyl, or $-\text{CO}_2\text{R}^2$; R^6 is H, alkyl, aralkyl, aryl, or monocyclic
10 heteroaryl; R^7 and R^8 are each independently H, halo, OH, or (C_1-C_6 alkyl)-O;
and R^9 is $=\text{O}$ or $=\text{NOR}^2$;

- A specific value for A is phenyl. Another specific value for A is pyridyl.
Other specific values for A include the following, wherein "~~~~" indicates the
15 points attachment to either B or NH and halo, (C_1-C_6)alkyl, and halo(C_1-C_6)alkyl
are as defined earlier in the definitions section:




A specific value for C is phenyl. Another specific value for C is pyridyl

-81-

and para-halo pyridyl. Other specific values for C include ortho- and para-halo phenyl, ortho-, para-dihalophenyl, ortho, meta-dihalophenyl, para-(C₁-C₆)alkoxyphenyl, para-(C₁-C₆)alkylphenyl, and meta-halo, para-(C₁-C₆)alkylphenyl.

5

One specific value for B is , wherein "~~~~" indicates the point of attachment; Q is NH or is absent; and wherein R' is OH, —O-(C₁-C₆)alkyl, or NR^{3'}R^{4'}, wherein R^{3'} and R^{4'} are independently H, (C₁-C₆)alkyl, or are joined together to form an optionally substituted saturated or unsaturated 5 to 7 membered ring.


10


Another specific value for B is phenyl or phenyl which is substituted at an ortho position with formyl, H₂NSO₂, MeSO₂, MeSO, MeS, (C₁-C₆)alkoxy carbonyl, OH, Ome, cyano, H₂N-CH₂—, HN-(C₁-C₆)alkyl-CH₂—, (C₁-C₆)alkyl₂N-CH₂—.


15

Another specific value for B is pyridyl or pyridyl substituted as provided in the previous position.


20

Another specific value for B is , wherein "~~~~" indicates the point of attachment.




Another specific value for B is , wherein "~~~~" indicates the point of attachment and which is an optionally substituted 4, 5, 6 or 7-

-82-






membered ring. When B is R^x and R^y can be H, halo, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)carboxyalkyl, or taken together can be O=. X^{III} can be CH₂, CH-OH, CH-CO₂(C₁-C₆)alkyl, O, S, NH, or N(C₁-C₆)alkyl, provided that when R^x and R^y taken together are O=, X^{III} is CH₂.

5




Another specific value for B is



or a tautomer thereof such as R^{18} or a tautomer thereof, Y^{II} can be CH₂, CH(C₁-C₆)alkyl, C-CH₂OH, C-CH₂O-(C₁-C₆)alkyl, NH, or N(C₁-C₆)alkyl. R¹⁸ is H, (C₁-C₆)alkyl, hydroxymethyl, CH₂O-(C₁-C₆)alkyl, or NR³R⁴.

10

Another specific value for B is a 5-membered heteroaryl ring which contains at least one nitrogen atom such as imidazolyl, pyrazolyl, triazolyl, tetrazolyl, oxadiazolyl, thiadiazolyl, oxazolyl, isoxazolyl, thiazolyl, or isothiazolyl, any of which may be optionally substituted with halo, (C₁-C₆)alkyl, hydroxy(C₁-C₆)alkyl, CH₂O-(C₁-C₆)alkyl, or halo(C₁-C₆)alkyl. B can also be




, wherein "wavy" indicates the point of attachment;

X^a, Y^a, and Z^a are each independently CR^c or N, wherein R^c is H or (C₁-C₆)alkyl; and R¹⁸ is H, (C₁-C₆)alkyl, hydroxymethyl, CH₂O-(C₁-C₆)alkyl, CH₂NR³R⁴, or,

15


-83-

NR^3R^4 , wherein R^3 and R^4 are as defined herein.



Another specific value for B is

, wherein “~”




indicates the point of attachment. When B is

, J_1 , J_2 , J_3 , and

- 5 J_4 can each be C, or one or more of J_1 , J_2 , J_3 , or J_4 can be N. R^{19} , R^{20} , R^{21} , and R^{22} are each independently H, halo, hydroxy, NH_2 , $\text{NR}^{23}\text{R}^{24}$, NO_2 , SH, $(\text{C}_1\text{-C}_6)\text{alkyl}$, $(\text{C}_1\text{-C}_6)\text{alkoxy}$, halo $(\text{C}_1\text{-C}_6)\text{alkyl}$, $(\text{C}_1\text{-C}_6)\text{alkanoyl}$, $(\text{C}_1\text{-C}_6)\text{alkoxycarbonyl}$, or $(\text{C}_1\text{-C}_6)\text{alkoxy}$, wherein R^{23} and R^{24} are each independently H, $(\text{C}_1\text{-C}_6)\text{alkyl}$, aralkyl, aryl, monocyclic heteroaryl, alkoxycarbonyl, aralkoxycarbonyl, $-\text{SO}_2\text{alkyl}$, or


- 10 joined together to form a saturated or unsaturated 3 to 7 membered ring.

Alternatively, R^{19} and R^{20} , R^{20} and R^{21} , or R^{21} and R^{22} , can be together with the carbons to which they are attached to form a 5, 6, or seven membered saturated or unsaturated cycloalkyl or heterocyclo ring, or an aryl or heteroaryl ring. When any of J_1 , J_2 , J_3 , or J_4 is N, R^{19} , R^{20} , R^{21} or R^{22} , respectively, is absent at that




- 15 position. Thus, specific values for B include

-84-




- 5 (C_6) alkoxycarbonyl, or (C_1-C_6) alkoxy, wherein R^{23} and R^{24} are each independently H, (C_1-C_6) alkyl, aralkyl, aryl, monocyclic heteroaryl, alkoxycarbonyl, aralkoxycarbonyl, $-SO_2$ alkyl, or joined together to form a saturated or unsaturated 3 to 7 membered ring, or R^{19} and R^{20} , R^{20} and R^{21} , or R^{21} and R^{22} , together with the carbons to which they are attached, form a 5, 6, or
- 10 seven membered saturated or unsaturated cycloalkyl or heterocyclo ring, or an aryl or heteroaryl ring.

Reference will now be made to a compound of Formula II.





II

-85-




In , “---” is a bond or is absent. A specific value for X^{II} is CH_2 . Other specific values for X^{II} include CH , O , NH and N . Specific values for R^{10} and R^{11} include H , hydroxy, methoxy, ethoxy, cyano,


trifluoromethyl, fluoro, NH_2 , carboethoxy, $\text{Me}-\text{C}(=\text{O})-\text{NH}-$, $\text{MeSO}_2\text{N}-$, ,
 5 wherein “~” indicates the point of attachment. Also, R^{10} and R^{11} can be taken



together to form $=\text{O}$ or $=\text{NOH}$. Thus, specific values for



-86-



A specific value for G is H, F, or methyl.


5

A specific value for C is phenyl. Another specific value for C is pyridyl.

Other specific values for C include ortho- and p--halo phenyl, ortho, p--dihalophenyl, ortho, meta-dihalophenyl, p-(C₁-C₆)alkoxyphenyl, p-(C₁-C₆)alkylphenyl, and meta-halo, p--(C₁-C₆)alkylphenyl.


10

A specific value for A is phenyl. Another specific value for A is pyridyl. Other specific values for A include the following, wherein "~~~~" indicates the points attachment to either B or NH and halo, (C₁-C₆)alkyl, and halo(C₁-C₆)alkyl are as defined earlier in the definitions section:



15


-87-





- One specific value for B is $\sim\sim\sim\text{Q}-\text{C}(=\text{O})-\text{R}^1$, wherein " $\sim\sim\sim$ " indicates the point of attachment; Q is NH or is absent; and wherein R¹ is OH, —O-(C₁-C₆)alkyl, or NR^{3'}R^{4'}, wherein R^{3'} and R^{4'} are independently H, (C₁-C₆)alkyl, or are joined together to form an optionally substituted saturated or unsaturated 5 to 7 membered ring.


- Another specific value for B is phenyl or phenyl which is substituted at an ortho position with formyl, H₂NSO₂, MeSO₂, MeSO, MeS, (C₁-C₆)alkoxycarbonyl, cyano, H₂N-CH₂—, HN-(C₁-C₆)alkyl-CH₂—, (C₁-C₆)alkyl₂N-CH₂—.

- Another specific value for B is pyridyl or pyridyl substituted as provided in the previous position.





Another specific value for B is , wherein " $\sim\sim\sim$ " indicates the point of attachment.




- Another specific value for B is , wherein " $\sim\sim\sim$ " indicates the point of attachment and which is an optionally substituted 4, 5, 6 or 7-

-88-









membered ring. When B is , R^x and R^y can be H, halo, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, (C_1-C_6) carboxyalkyl, or taken together can be $O=$. X^{III} can be CH_2 , $CH-OH$, $CH-CO_2(C_1-C_6)$ alkyl, O, S, NH, or $N(C_1-C_6)$ alkyl, provided that when R^x and R^y taken together are $O=$, X^{III} is CH_2 .

5




Another specific value for B is



or  or . When B is  or a tautomer thereof, Y^{II} can be CH_2 , $CH(C_1-C_6)$ alkyl, $C-CH_2OH$, $C-CH_2O-(C_1-C_6)$ alkyl, NH, or $N(C_1-C_6)$ alkyl. R^{18} is H, (C_1-C_6) alkyl, hydroxymethyl, $CH_2O-(C_1-C_6)$ alkyl, or NR^3R^4 .

10

Another specific value for B is a 5-membered heteroaryl ring which contains at least one nitrogen atom such as imidazolyl, pyrazolyl, triazolyl, tetrazolyl, oxadiazolyl, thiadiazolyl, oxazolyl, isoxazolyl, thiazolyl, or isothiazolyl, any of which may be optionally substituted with halo, (C_1-C_6) alkyl, hydroxy (C_1-C_6) alkyl, $CH_2O-(C_1-C_6)$ alkyl, or halo (C_1-C_6) alkyl. B can also be



, wherein "wavy" indicates the point of attachment;


X^a , Y^a , and Z^a are each independently CR^c or N, wherein R^c is H or (C_1-C_6) alkyl;

and

R^{18} is H, (C_1-C_6) alkyl, hydroxymethyl, $CH_2O-(C_1-C_6)$ alkyl, $CH_2NR^3R^4$,


-89-

or NR^3R^4 , wherein R^3 and R^4 are as defined herein.



Another specific value for B is


, wherein “~”



indicates the point of attachment. When B is


, J_1 , J_2 , J_3 , and

- 5 J_4 can each be C, or one or more of J_1 , J_2 , J_3 , or J_4 can be N. R^{19} , R^{20} , R^{21} , and R^{22} are each independently H, halo, hydroxy, NH_2 , $\text{NR}^{23}\text{R}^{24}$, NO_2 , SH, $(\text{C}_1\text{-C}_6)\text{alkyl}$, $(\text{C}_1\text{-C}_6)\text{alkoxy}$, halo $(\text{C}_1\text{-C}_6)\text{alkyl}$, $(\text{C}_1\text{-C}_6)\text{alkanoyl}$, $(\text{C}_1\text{-C}_6)\text{alkoxycarbonyl}$, or $(\text{C}_1\text{-C}_6)\text{alkoxy}$, wherein R^{23} and R^{24} are each independently H, $(\text{C}_1\text{-C}_6)\text{alkyl}$, aralkyl, aryl, monocyclic heteroaryl, alkoxycarbonyl, aralkoxycarbonyl, $-\text{SO}_2\text{alkyl}$, or
- 10 joined together to form a saturated or unsaturated 3 to 7 membered ring. Alternatively, R^{19} and R^{20} , R^{20} and R^{21} , or R^{21} and R^{22} , can be together with the carbons to which they are attached to form a 5, 6, or seven membered saturated or unsaturated cycloalkyl or heterocyclo ring, or an aryl or heteroaryl ring. When any of J_1 , J_2 , J_3 , or J_4 is N, R^{19} , R^{20} , R^{21} or R^{22} , respectively, is absent at that




- 15 position. Thus, specific values for B include

-90-




- 5 C₆)alkoxycarbonyl, or (C₁-C₆)alkoxy, wherein R²³ and R²⁴ are each independently H, (C₁-C₆)alkyl, aralkyl, aryl, monocyclic heteroaryl, alkoxycarbonyl, aralkoxycarbonyl, -SO₂alkyl, or joined together to form a saturated or unsaturated 3 to 7 membered ring, or R¹⁹ and R²⁰, R²⁰ and R²¹, or R²¹ and R²², together with the carbons to which they are attached, form a 5, 6, or
- 10 seven membered saturated or unsaturated cycloalkyl or heterocyclo ring, or an aryl or heteroaryl ring.

Reference will now be made to a compound of Formula III:





III

-91-




In , “---” is a bond or is absent. A specific value for X^{II} is CH_2 . Other specific values for X^{II} include CH, O, NH and N. Specific values for R^{10} and R^{11} include H, hydroxy, methoxy, ethoxy, cyano,

- 5 trifluoromethyl, fluoro, NH_2 , carboethoxy, $Me-\overset{O}{\parallel}-NH-$, $MeSO_2N-$, , wherein “~” indicates the point of attachment. Also, R^{10} and R^{11} taken together





can be $=O$ or $=NOH$. Thus, specific values for

include




-92-



In

- 5 halo. R₁₂ and R₁₃ are each independently H, halo, (C₁-C₆)alkyl, or (C₁-




C₆)alkoxy. Specific values for pyridyl, p-chlorophenyl, p-fluorophenyl, p-bromophenyl, p-methoxyphenyl, p-toluy, p-chloropyridyl, ortho, p-difluorophenyl, meta, p-difluorophenyl, and meta fluoro, paratoluy.


10

A specific value for G is H, F, or methyl.

- A specific value for A is phenyl. Another specific value for A is pyridyl. Other specific values for A include the following, wherein “~~~~” indicates the points attachment to either B or NH and halo, (C₁-C₆)alkyl, and halo(C₁-C₆)alkyl are as defined earlier in the definitions section:
- 15


-93-




One specific value for B is , wherein “~” indicates the point of attachment; Q is NH or is absent; and wherein R^{1'} is OH, —O-(C₁-C₆)alkyl, or NR^{3'}R^{4'}, wherein R^{3'} and R^{4'} are independently H, (C₁-C₆)alkyl, or are joined together to form an optionally substituted saturated or unsaturated 5 to 7 membered ring.


Another specific value for B is phenyl or phenyl which is substituted at an ortho position with formyl, H₂NSO₂, MeSO₂, MeSO, MeS, (C₁-C₆)alkoxycarbonyl, cyano, H₂N-CH₂—, HN-(C₁-C₆)alkyl-CH₂—, (C₁-C₆)alkyl₂N-CH₂—.


Another specific value for B is pyridyl or pyridyl substituted as provided in the previous position.





Another specific value for B is , wherein “~” indicates the point of attachment.


-94-



Another specific value for B is , wherein "~~~~" indicates the point of attachment and which is an optionally substituted 4, 5, 6 or 7-






membered ring. When B is , R^x and R^y can be H, halo, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, (C_1-C_6) carboxyalkyl, or taken together can be $O=$. X^{III} can be CH_2 , $CH-OH$, $CH-CO_2(C_1-C_6)alkyl$, O, S, NH, or $N(C_1-C_6)alkyl$, provided that when R^x and R^y taken together are $O=$, X^{III} is CH_2 .






Another specific value for B is

 or a tautomer thereof such as




10

or  or . When B is  or a tautomer thereof, Y^{II} can be CH_2 , $CH(C_1-C_6)alkyl$, $C-CH_2OH$, $C-CH_2O-(C_1-C_6)alkyl$, NH, or $N(C_1-C_6)alkyl$. R^{18} is H, $(C_1-C_6)alkyl$, hydroxymethyl, $CH_2O-(C_1-C_6)alkyl$, or NR^3R^4 .

Another specific value for B is a 5-membered heteroaryl ring which
15 contains at least one nitrogen atom such as imidazolyl, pyrazolyl, triazolyl, tetrazolyl, oxadiazolyl, thiadiazolyl, oxazolyl, isoxazolyl, thiazolyl, or isothiazolyl, any of which may be optionally substituted with halo, $(C_1-C_6)alkyl$,

-95-

hydroxy(C₁-C₆)alkyl, CH₂O-(C₁-C₆)alkyl, or halo(C₁-C₆)alkyl. B can also be




, wherein “~” indicates the point of attachment;

X^a, Y^a, and Z^a are each independently CR^c or N, wherein R^c is H or (C₁-C₆)alkyl;


and R¹⁸ is H, (C₁-C₆)alkyl, hydroxymethyl, CH₂O-(C₁-C₆)alkyl, CH₂NR³R⁴, or

5 NR³R⁴, wherein R³ and R⁴ are as defined herein.



Another specific value for B is

, wherein “~”



indicates the point of attachment. When B is , J₁, J₂, J₃, and

J₄ can each be C, or one or more of J₁, J₂, J₃, or J₄ can be N. R¹⁹, R²⁰, R²¹, and R²²

10 are each independently H, halo, hydroxy, NH₂, NR²³R²⁴, NO₂, SH, (C₁-C₆)alkyl,

(C₁-C₆)alkoxy, halo(C₁-C₆)alkyl, (C₁-C₆)alkanoyl, (C₁-C₆)alkoxycarbonyl, or (C₁-


C₆)alkoxy, wherein R²³ and R²⁴ are each independently H, (C₁-C₆)alkyl, aralkyl,

aryl, monocyclic heteroaryl, alkoxycarbonyl, aralkoxycarbonyl, -SO₂alkyl, or



joined together to form a saturated or unsaturated 3 to 7 membered ring.

15 Alternatively, R¹⁹ and R²⁰, R²⁰ and R²¹, or R²¹ and R²², can be together with the carbons to which they are attached to form a 5, 6, or seven membered saturated or unsaturated cycloalkyl or heterocyclo ring, or an aryl or heteroaryl ring. When any of J₁, J₂, J₃, or J₄ is N, R¹⁹, R²⁰, R²¹ or R²², respectively, is absent at that

-96-



position. Thus, specific values for B include



, wherein “~” indicates the point of attachment; and R^{19} ,

R^{20} , and R^{21} , R^{22} are each independently H, halo, hydroxy, NH_2 , $NR^{23}R^{24}$, NO_2 ,

5 SH, $(C_1-C_6)alkyl$, $(C_1-C_6)alkoxy$, halo $(C_1-C_6)alkyl$, $(C_1-C_6)alkanoyl$, $(C_1-$

$C_6)alkoxycarbonyl$, or $(C_1-C_6)alkoxy$, wherein R^{23} and R^{24} are each

independently H, $(C_1-C_6)alkyl$, aralkyl, aryl, monocyclic heteroaryl,

alkoxycarbonyl, aralkoxycarbonyl, $-SO_2alkyl$, or joined together to form a

saturated or unsaturated 3 to 7 membered ring, or R^{19} and R^{20} , R^{20} and R^{21} , or R^{21}


10 and R^{22} , together with the carbons to which they are attached, form a 5, 6, or

seven membered saturated or unsaturated cycloalkyl or heterocyclo ring, or an


aryl or heteroaryl ring.

Reference will now be made to a compound of Formula IV:

-97-





IV



In


, "----" is a bond or is absent. A specific value for X^{II} is CH_2 . Other specific values for X^{II} include CH , O , NH and N . Specific values for R^{10} and R^{11} include H , hydroxy, methoxy, ethoxy, cyano,

- 5 trifluoromethyl, fluoro, NH_2 , carboethoxy, $Me-C(=O)-NH-$, $MeSO_2N-$, , wherein "~~~~" indicates the point of attachment. Also, R^{10} and R^{11} taken together





can be $=O$ or $=NOH$. Thus, specific values for


include



-98-




- 5 In R^{13} , Z is N, CH, C(C₁-C₆)alkyl, C-(C₁-C₆)alkoxy, or C-halo. R¹² and R¹³ are each independently H, halo, (C₁-C₆)alkyl, (C₁-C₆)alkoxy.




Specific values for R^{13} therefore include phenyl, pyridyl, p-chlorophenyl, p-fluorophenyl, p-bromophenyl, p-methoxyphenyl, p-tolyl, p-

-99-

chloropyridyl, ortho, p-difluorophenyl, meta, para-difluorophenyl, and meta fluoro, p-methyl phenyl.



- In R^{14} , X and Y are each independently N, CH, C-halo, C(C₁-C₆)alkyl, C-halo(C₁-C₆)alkyl, or C-heterocyclo. R^{14} and R^{15} are each independently H, halo, (C₁-C₆)alkyl, halo(C₁-C₆)alkyl, or heterocyclo. Specific



A specific value for G is H, F, or methyl.

- One specific value for B is $\text{---Q---C(=O)---R}^{1'}$, wherein "—" indicates the point of attachment; Q is NH or is absent; and wherein $\text{R}^{1'}$ is OH, —O-(C₁-C₆)alkyl, or $\text{NR}^{3'}\text{R}^{4'}$, wherein $\text{R}^{3'}$ and $\text{R}^{4'}$ are independently H, (C₁-C₆)alkyl, or are joined together to form a optionally substituted saturated or unsaturated 5 to

-100-


7 membered ring.

Another specific value for B is phenyl or phenyl which is substituted at an ortho position with formyl, H_2NSO_2 , MeSO_2 , MeSO , MeS , $(\text{C}_1\text{-C}_6)\text{alkoxycarbonyl}$,
 5 cyano, $\text{H}_2\text{N-CH}_2\text{—}$, $\text{HN-(C}_1\text{-C}_6\text{)alkyl-CH}_2\text{—}$, $(\text{C}_1\text{-C}_6\text{)alkyl}_2\text{N-CH}_2\text{—}$.

Another specific value for B is pyridyl or pyridyl substituted as provided in the previous position.


10

Another specific value for B is




, wherein “~” indicates the

point of attachment.



Another specific value for B is , wherein “~” indicates the point of attachment and which is an optionally substituted 4, 5, 6 or 7-




15




membered ring. When B is , R^x and R^y can be H, halo, $(\text{C}_1\text{-C}_6)\text{alkyl}$, $(\text{C}_1\text{-C}_6)\text{alkoxy}$, $(\text{C}_1\text{-C}_6)\text{carboxyalkyl}$, or taken together can be O= . X^{III} can be CH_2 , CH-OH , $\text{CH-CO}_2(\text{C}_1\text{-C}_6)\text{alkyl}$, O, S, NH, or $\text{N(C}_1\text{-C}_6\text{)alkyl}$, provided that when R^x and R^y taken together are O= , X^{III} is CH_2 .

-101-

Another specific value for B is



or a tautomer thereof such as




When B is or a tautomer thereof, Y^{II} can be CH_2 , $CH(C_1-C_6)alkyl$, $C-CH_2OH$, $C-CH_2O-(C_1-C_6)alkyl$, NH , or $N(C_1-C_6)alkyl$. R^{18} is H , $(C_1-C_6)alkyl$, hydroxymethyl, $CH_2O-(C_1-C_6)alkyl$, or NR^3R^4 .

5

Another specific value for B is a 5-membered heteroaryl ring which contains at least one nitrogen atom such as imidazolyl, pyrazolyl, triazolyl, tetrazolyl, oxadiazolyl, thiadiazolyl, oxazolyl, isoxazolyl, thiazolyl, or isothiazolyl, any of which may be optionally substituted with halo, $(C_1-C_6)alkyl$, hydroxy $(C_1-C_6)alkyl$, $CH_2O-(C_1-C_6)alkyl$, or halo $(C_1-C_6)alkyl$. B can also be

10




, wherein "wavy" indicates the point of attachment;

X^a , Y^a , and Z^a are each independently CR^c or N , wherein R^c is H or $(C_1-C_6)alkyl$; and

R^{18} is H , $(C_1-C_6)alkyl$, hydroxymethyl, $CH_2O-(C_1-C_6)alkyl$, or NR^3R^4 .


15



Another specific value for B is

, wherein "wavy" indicates the

-102-




point of attachment. When B is \mathbb{R}^{20} , J_1, J_2, J_3 , and J_4 can each be

C, or one or more of J₁, J₂, J₃, or J₄ can be N. R¹⁹, R²⁰, R²¹, and R²² are each independently H, halo, hydroxy, NH₂, NR²³R²⁴, NO₂, SH, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl, (C₁-C₆)alkanoyl, (C₁-C₆)alkoxycarbonyl, or (C₁-


5 C₆)alkoxy, wherein R²³ and R²⁴ are each independently H, (C₁-C₆)alkyl, aralkyl, aryl, monocyclic heteroaryl, alkoxy carbonyl, aralkoxy carbonyl, -SO₂alkyl, or joined together to form a saturated or unsaturated 3 to 7 membered ring.

Alternatively, R¹⁹ and R²⁰, R²⁰ and R²¹, or R²¹ and R²², can be together with the carbons to which they are attached to form a 5, 6, or seven membered saturated or

10 unsaturated cycloalkyl or heterocyclo ring, or an aryl or heteroaryl ring. When any of J₁, J₂, J₃, or J₄ is N, R¹⁹, R²⁰, R²¹ or R²², respectively, is absent at that position.



Thus, specific values for B include



"~" indicates the point of attachment; and R¹⁹, R²⁰, and R²¹, R²² are each



15 independently H, halo, hydroxy, NH_2 , $\text{NR}^{23}\text{R}^{24}$, NO_2 , SH, $(\text{C}_1\text{-C}_6)\text{alkyl}$, $(\text{C}_1\text{-C}_6)\text{alkoxy}$, halo $(\text{C}_1\text{-C}_6)\text{alkyl}$, $(\text{C}_1\text{-C}_6)\text{alkanoyl}$, $(\text{C}_1\text{-C}_6)\text{alkoxycarbonyl}$, or $(\text{C}_1\text{-C}_6)\text{alkoxy}$, wherein R^{23} and R^{24} are each independently H, $(\text{C}_1\text{-C}_6)\text{alkyl}$, aralkyl, aryl, monocyclic heteroaryl, alkoxycarbonyl, aralkoxycarbonyl, $-\text{SO}_2\text{alkyl}$, or

-103-


joined together to form a saturated or unsaturated 3 to 7 membered ring, or R^{19} and R^{20} , R^{20} and R^{21} , or R^{21} and R^{22} , together with the carbons to which they are attached, form a 5, 6, or seven membered saturated or unsaturated cycloalkyl or heterocyclo ring, or an aryl or heteroaryl ring.

5

Reference will now be made to a compound of Formula V:




In , “---” is a bond or is absent. A specific value for X^{II} is CH_2 . Other specific values for X^{II} include CH , NH and N . Specific values
10 for R^{10} and R^{11} include H , hydroxy, methoxy, ethoxy, cyano, trifluoromethyl,

fluoro, NH_2 , carboethoxy, $Me-C(=O)-NH-$, $MeSO_2N-$, , wherein “~” indicates the point of attachment. Also, R^{10} and R^{11} taken together can be $=O$ or

-104-


=NOH. Thus, specific values for

include




5

, or




In R^{12} and R^{13} are each independently H, halo, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, or C-halo. Z is N, CH, C(C₁-C₆)alkyl, C-(C₁-C₆)alkoxy, or C-halo.

-105-




Specific values for R^{12} and R^{13} therefore include phenyl, pyridyl, p-chlorophenyl, p-fluorophenyl, p-bromophenyl, p-methoxyphenyl, p-tolyl, p-chloropyridyl, ortho, p-difluorophenyl, meta, p-difluorophenyl, and meta fluoro, p-methyl phenyl.


5



In R^{14} and R^{15} , X and Y are each independently N, CH, C-halo, C(C₁-C₆)alkyl, C-halo(C₁-C₆)alkyl, or C-heterocyclo. R^{14} and R^{15} are each independently H, halo, (C₁-C₆)alkyl, halo(C₁-C₆)alkyl, or heterocyclo. Specific




10









A specific value for G is H, F, or methyl.

15


-106-





fused heteroaryl ring, provided that when  is present, Y' is C. X' and Y' are each independently N, CH, C-halo, C-halo, C-(C₁-C₆)alkyl, C-halo(C₁-C₆)alkyl, C-(C₁-C₆)alkoxy, C-OH, C-SO₂Me, C-SO₂NH₂, C-SOMe, C-SMe, C-CO₂(C₁-C₆)alkyl, C-CN, or C-CH₂NH₂. Specific values for




-107-





Reference will now be made to a compound of Formula VI.



5 In , “----” is a bond or is absent. A specific value for X^{II} is CH_2 . Other specific values for X^{II} include CH , NH and N . Specific values for R^{10} and R^{11} include H , hydroxy, methoxy, ethoxy, cyano, trifluoromethyl,


fluoro, NH_2 , carboethoxy, $Me-\overset{O}{\parallel}-NH-$, $MeSO_2N-$, , wherein “~” indicates the point of attachment. Also, R^{10} and R^{11} taken together can be $=O$ or





10 $=NOH$. Thus, specific values for


include

-108-




5 In , Z is N, CH, C(C₁-C₆)alkyl, C-(C₁-C₆)alkoxy, or C-halo. R¹² and R¹³ are each independently H, halo, (C₁-C₆)alkyl, (C₁-C₆)alkoxy.




Specific values for  therefore include phenyl, pyridyl, p-chlorophenyl, p-fluorophenyl, p-bromophenyl, p-methoxyphenyl, p-tolyl, p-

chloropyridyl, ortho, p-difluorophenyl, meta, p-difluorophenyl, and meta fluoro, p-methyl phenyl.







In C_B , X and Y are each independently N, CH, C-halo, C(C₁-C₆)alkyl, C-halo(C₁-C₆)alkyl, or C-heterocyclo. R¹⁴ and R¹⁵ are each independently H, halo, (C₁-C₆)alkyl, halo(C₁-C₆)alkyl, or heterocyclo. Specific



values for


include


include , , 



10


A specific value for G is H, F, or methyl.



In , Q is NH or is absent. R¹ is OH, -O-(C₁-C₆)alkyl, (C₃-C₇)cycloalkyl, (C₃-C₇)heterocyclo, (C₄-C₇)cycloalkenyl, (C₄-C-
7)heterocycloalkenyl, aryl, monocyclic heteroaryl, or -NR³R⁴, wherein R³ and R⁴ are independently H, (C₁-C₆)alkyl, aralkyl, aryl, monocyclic heteroaryl,


-110-

alkoxycarbonyl, aralkoxycarbonyl, $-\text{SO}_2\text{alkyl}$, or joined together to form a saturated or unsaturated 5 to 7 membered ring. Thus, specific values for




5

Reference will now be made to a compound of Formula VII.



VII


-111-




In

, "----" is a bond or is absent. A specific value for


X^{11} is CH_2 . Other specific values for X^{11} include CH, O, NH and N. Specific values for R^{10} and R^{11} include H, hydroxy, methoxy, ethoxy, cyano,

trifluoromethyl, fluoro, NH_2 , carboethoxy, $Me-C(=O)-NH-$, $MeSO_2N-$, ,
 5 wherein "~~~~" indicates the point of attachment. Also, R^{10} and R^{11} taken together






can be =O or =NOH. Thus, specific values for


include




-112-




In , Z is N, CH, C(C₁-C₆)alkyl, C(C₁-C₆)alkoxy, or C-halo. R¹² and R¹³ are each independently H, halo, (C₁-C₆)alkyl, (C₁-C₆)alkoxy.



- 5 Specific values for therefore include phenyl, pyridyl, p-chlorophenyl, p-fluorophenyl, p-bromophenyl, p-methoxyphenyl, p-toluy, p-chloropyridyl, ortho, p-difluorophenyl, meta, p-difluorophenyl, and meta fluoro, p-methyl phenyl.




- 10 In , X and Y are each independently N, CH, C-halo, C(C₁-C₆)alkyl, C-halo(C₁-C₆)alkyl, or C-heterocyclo. R¹⁴ and R¹⁵ are each independently H, halo, (C₁-C₆)alkyl, halo(C₁-C₆)alkyl, or heterocyclo. Specific




5

A specific value for G is H, F, or methyl.




In compounds of Formula VII,  is an optionally substituted


4, 5, 6 or 7-membered ring, wherein R^x and R^y are H, halo, hydroxymethyl, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, or wherein R^x and R^y are H, halo, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)carboxyalkyl, or taken together are O=. X^{III} is CH₂, CH-OH, CH-CO₂(C₁-C₆)alkyl, O, S, NH, or N(C₁-C₆)alkyl, provided that when R^x and R^y taken together are O=, X^{III} is CH₂. Thus, specific values for



-114-




. Any of the specific values for




- may be further optionally substituted with one, two or three
 5 substituent groups selected from the group consisting of halo, (C₁-C₆)alkyl, halo(C₁-C₆)alkyl, or hydroxymethyl.

Reference will now be made to a compound of Formula VIII.




VIII


-115-



In


, "----" is a bond or is absent. A specific value for X'' is CH_2 . Other specific values for X'' include CH , O , NH and N . Specific values for R'' and R''' include H , hydroxy, methoxy, ethoxy, cyano,

trifluoromethyl, fluoro, NH_2 , carboethoxy, $\text{Me}-\text{C}(=\text{O})-\text{NH}-$, $\text{MeSO}_2\text{N}-$, ,
 5 wherein "~~~~" indicates the point of attachment. Also, R'' and R''' taken together





can be $=\text{O}$ or $=\text{NOH}$. Thus, specific values for


include




-116-



In R^{12} and R^{13} , Z is N, CH, C(C₁-C₆)alkyl, C(C₁-C₆)alkoxy, or C-halo. R^{12} and R^{13} are each independently H, halo, (C₁-C₆)alkyl, (C₁-C₆)alkoxy.




- 5 Specific values for R^{12} and R^{13} therefore include phenyl, pyridyl, p-chlorophenyl, p-fluorophenyl, p-bromophenyl, p-methoxyphenyl, p-toluy, p-chloropyridyl, ortho, p-difluorophenyl, meta, p-difluorophenyl, and meta fluoro, p-methyl phenyl.






- 10 In R^{14} and R^{15} , X and Y are each independently N, CH, C-halo, C(C₁-C₆)alkyl, C-halo(C₁-C₆)alkyl, or C-heterocyclo. R^{14} and R^{15} are each independently H, halo, (C₁-C₆)alkyl, halo(C₁-C₆)alkyl, or heterocyclo. Specific

-117-



5

In compounds of Formula VIII,



R_{18} , Y^{II} is CH, CR^{18} , or N. R^{18} is H, (C_1-C_6) alkyl, hydroxymethyl,


$CH_2O-(C_1-C_6)$ alkyl, or NR^3R^4 , wherein R^3 and R^4 are each independently H,

10 (C_1-C_6) alkyl, aralkyl, aryl, monocyclic heteroaryl, alkoxycarbonyl,

-118-


aralkoxycarbonyl, -SO₂alkyl, or joined together to form a saturated or unsaturated

3 to 7 membered ring. Thus, specific values for R_{18} include




5 A specific value for G is H, F, or methyl.


Reference will now be made to a compound of Formula IX.



10 X^{II} is CH₂. Other specific values for X^{II} include CH, O, NH and N. Specific values for R¹⁰ and R¹¹ include H, hydroxy, methoxy, ethoxy, cyano,


-119-

trifluoromethyl, fluoro, NH_2 , carboethoxy, $\text{Me}-\text{C}(=\text{O})-\text{NH}-$, $\text{MeSO}_2\text{N}-$, , wherein "~~~~" indicates the point of attachment. Also, R^{10} and R^{11} taken together




can be $=\text{O}$ or $=\text{NOH}$. Thus, specific values for


include




-120-




R^{12} , Z is N, CH, C(C₁-C₆)alkyl, or C-halo. R^{12} and R^{13} are each independently H, halo, (C₁-C₆)alkyl, (C₁-C₆)alkoxy. Specific values for





R^{13} therefore include phenyl, pyridyl, p-chlorophenyl, p-fluorophenyl, p-bromophenyl, p-methoxyphenyl, p-tolyl, p-chloropyridyl, ortho, p-difluorophenyl, meta, p-difluorophenyl, and meta fluoro, p-methyl phenyl.




X and Y are each independently N, CH, C-halo, C(C₁-C₆)alkyl, C-halo(C₁-C₆)alkyl, or C-heterocyclo. R^{14} and R^{15} are each independently H, halo, (C₁-C₆)alkyl, halo(C₁-C₆)alkyl, or heterocyclo. Specific



-121-




, X^a , Y^a , and Z^a are each independently CH, CR^{18} or N; and R^{18} is H, (C_1-C_6) alkyl, hydroxymethyl, $CH_2O-(C_1-C_6)$ alkyl, $CH_2NR^3R^4$, or NR^3R^4 , wherein R^3 and R^4 are each independently H, (C_1-C_6) alkyl, aralkyl, aryl, monocyclic heteroaryl, alkoxy carbonyl, aralkoxy carbonyl, $-SO_2$ alkyl, or joined together to form a saturated or unsaturated 3 to 7 membered ring. Thus, specific





include imidazolyl, pyrazolyl, triazolyl, tetrazolyl, oxadiazolyl, thiadiazolyl, oxazolyl, isoxazolyl, thiazolyl, and isothiazolyl, any of which may be optionally substituted with halo, (C_1-C_6) alkyl, hydroxy (C_1-C_6) alkyl,

$CH_2O-(C_1-C_6)$ alkyl, or halo (C_1-C_6) alkyl. Other values for R^{18}





A specific value for G is H, F, or methyl.

Reference will now be made to a compound of Formula X.



5 In § , “---” is a bond or is absent. A specific value for X^{II} is CH₂. Other specific values for X^{II} include CH, O, NH and N. Specific values for R¹⁰ and R¹¹ include H, hydroxy, methoxy, ethoxy, cyano,


trifluoromethyl, fluoro, NH_2 , carboethoxy, $\text{Me}-\text{C}(=\text{O})-\text{NH}-$, $\text{MeSO}_2\text{N}-$, ,
wherein “~” indicates the point of attachment. Also, R^{10} and R^{11} taken together




10 can be =O or =NOH. Thus, specific values for

include


-123-



5




In
halo. R¹² and R¹³ are each independently H, halo, (C₁-C₆)alkyl, -(C₁-C₆)alkoxy, or C-




Specific values for R^{13} therefore include phenyl, pyridyl, p-chlorophenyl, p-fluorophenyl, p-bromophenyl, p-methoxyphenyl, p-toluyyl, p-chloropyridyl, ortho, p-difluorophenyl, meta, p-difluorophenyl, and meta fluoro, p-methyl phenyl.


5




In R^{14} , R^{15} , X and Y are each independently N, CH, C-halo, C(C₁-C₆)alkyl, C-halo(C₁-C₆)alkyl, or C-heterocyclo. R^{14} and R^{15} are each independently H, halo, (C₁-C₆)alkyl, halo(C₁-C₆)alkyl, or heterocyclo. Specific




-125-



In , J₁, J₂, J₃, and J₄ are each C, or one of J₁, J₂, J₃, and J₄ is N. R¹⁹, R²⁰, R²¹, and R²² are each independently H, halo, hydroxy, NH₂, NR²³R²⁴, NO₂, SH, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl, (C₁-C₆)alkanoyl, (C₁-C₆)alkoxycarbonyl, or (C₁-C₆)alkoxy, wherein R²³ and R²⁴ are each independently H, (C₁-C₆)alkyl, aralkyl, aryl, monocyclic heteroaryl, alkoxy, alkoxy, or joined together to form a saturated or unsaturated 3 to 7 membered ring, or R¹⁹ and R²⁰, R²⁰ and R²¹, or R²¹ and R²², together with the carbons to which they are attached, form a 5, 6, or 7 membered saturated or unsaturated cycloalkyl or heterocyclo ring, or an aryl or heteroaryl ring. Also, when any of J₁, J₂, J₃, or J₄ is N, R¹⁹, R²⁰, R²¹ or R²², respectively, is absent at that position. Thus, specific values for



-126-



A specific value for G is H, F, or methyl.

5 PREPARATION OF COMPOUNDS OF THE INVENTION

Processes and novel intermediates for preparing compounds of formula I are provided as further embodiments of the invention and are illustrated by the following procedures in which the meanings of the generic radicals are as given above unless otherwise qualified. Certain compounds of formula I are useful as
10 intermediates for preparing other compounds of formula I.

It is also noted that compounds of formula I can be prepared using protecting groups. It is to be noted that the appropriate use and choice of protecting groups is well-known by one skilled in the art, and is not limited to the
15 specific examples below. It is also to be understood that such groups not only serve to protect chemically reactive sites, but also to enhance solubility or otherwise change physical properties. A good general reference for protecting group preparation and deprotection is Protecting Groups in Organic Synthesis by T.W. Green and P.G. Wuts. A number of general reactions such as oxidations and
20 reductions etc. are not shown in detail but can be done by methods understood by one skilled in the art. General transformations are well-reviewed in Comprehensive Organic Transformation by Richard Larock, and the series” published by Wiley-Interscience. In general, the starting materials are obtained from commercial sources unless otherwise indicated.

25

It will be appreciated by those skilled in the art that compounds of the invention having one or more chiral centers may exist in and be isolated in optically active and racemic forms. Some compounds may exhibit polymorphism. It is to be understood that the present invention encompasses any racemic,

-127-

optically-active, polymorphic, geometric, or stereoisomeric form, or mixtures thereof, of a compound of the invention, which possess the useful properties described herein, it being well known in the art how to prepare optically active forms (for example, by resolution of the racemic form by recrystallization techniques, by synthesis from optically-active starting materials, by chiral synthesis, or by chromatographic separation using a chiral stationary phase) and how to determine activity or cytotoxicity using the standard tests described herein, or using other similar tests which are well known in the art.

10 It will be further appreciated by the skilled artisan that the following schemes depict the synthesis of some compounds of the invention. It is to be understood, however, that compounds of the invention other than those specifically disclosed can be prepared using the strategies depicted in the schemes.


15 One method for making compounds of the invention is disclosed in Scheme 1. The route is useful for a wide variety of starting materials with variable W^1 groups, provided the appropriate protecting group is utilized, if necessary. The scheme is also employed for both racemic mixtures and enantiomerically pure compounds. The method comprises reacting an amino acid having Formula A with a reagent capable of forming a protecting group on the amino group of an amino acid to form compound with Formula A-1. In Scheme 1, P^1 is a protecting group and W^1 is the same as defined above. The compound of Formula A-1 is next converted into an acid halide. Alternatively the carboxylic acid may be activated by a number of coupling reagents, but not limited to, such as BOP, PYBOP, HATU, EEDQ, or CDI. The acid halide is then reacted with a haloaniline or a haloaminoheterocycle to form a halide having Formula D, where Y is a halogen and A is as defined above. The compound with Formula D is then subjected to a coupling reaction with a compound having B to give a compound of Formula E. The protecting group is then removed from compound E and the resulting compound reacted with an isocyanate having C to form a compound of Formula I. Finally, as relating to Scheme 1 and subsequent schemes, it is to be noted that compounds of the invention wherein G has any of the provided

-128-

meanings other than H can be easily prepared by techniques available to the skilled artisan. For example, compounds wherein G is alkyl can be prepared by alkylation of I, or by alkylation of a corresponding ester of A, B D, or E.

5


Scheme 1



Scheme 2 provides a more detailed approach to the preparation of compounds of the invention. The synthetic route starts by protecting an amino acid A with di-*tert*-butyl dicarbonate in tetrahydrofuran with 2M NaOH as the base. The resulting BOC-protected amino acid B-1 is converted into the acid chloride using oxalyl chloride, with pyridine added to prevent deprotection. The acid chloride or acid is then immediately reacted with a bromoaniline or bromoaminoheterocycle of choice. The resulting bromide is then typically subjected to a Suzuki coupling with a boronic acid, although other coupling conditions may be used to provide D-1. Compound D-1 is then deprotected with 33% trifluoroacetic acid in dichloromethane and allowed to react with an isocyanate in the presence of an amine base such as triethylamine in tetrahydrofuran or the like. This route is useful for preparing compounds containing biaryl A-B groups; i.e., where both A and B are aryl.

-129-


Scheme 2



- 5 Another method of making compounds of the invention is depicted in Scheme 3. The route summarized in Scheme 3 is useful for synthesizing a wide variety of A-B groups. The chemistry is similar to Scheme 1, but the second step (i.e., the conversion of B to D) introduces different substituted aniline or aminoheterocycle. By selecting an appropriately substituted aniline or
- 10 aminoheteroaryl group, a wide variety of B groups can be introduced. For example, when Z is a cyano group, the corresponding imidazole and imidazoline compounds can be made using techniques known to the skilled artisan. When Z is a methyl ester or the like, the corresponding amides and ketones can be synthesized via a number of methods available to the skilled artisan. For example,
- 15 an amide is prepared by hydrolyzing the ester with lithium hydroxide or TMSOK or the like, treating the resulting carboxylic acid with a typical peptide coupling reagent such as HATU or PYBOP followed by introduction of the desired amine.

-130-


Scheme 3



Compounds of the invention may also be prepared as provided in Scheme


4. Thus, a mixture of **F** (where **Z** is a halide and **Y** is an amino or nitro group), **B** (where **B** is a nitrogen-containing ring, an amine or amide), K_3PO_4 , CuI , and *trans*-diaminocyclohexane is heated in dioxane at reflux to obtain compound **G**. In the case where **Y** is a nitro group, the moiety is reduced to the amino group with $RaNi$ and $EtOH$ under a hydrogen atmosphere. The appropriate aniline **G**, EEDQ, triethylamine and carboxylic acid **H** are heated at reflux in chloroform to produce **J**. Finally, a solution of compound **J**, TFA and DCM is stirred at room temperature for 1 h and then concentrated under reduced pressure. The resulting oil is dissolved in THF and cooled to 0 °C followed by the addition of triethylamine and the appropriate isocyanate to produce invention compound **K**.

Scheme 4



-131-


Scheme 5 provides an approach to compounds of the invention wherein X^{II}




in (c.f., Formulas II-X), is N and “----” is a bond. Thus, aniline **M** is converted to an acrylamide **N** by the addition of acryloyl chloride and a base, such as saturated sodium bicarbonate, in a solvent such as ethyl acetate at room temperature. Alternatively, the aniline may be converted to **N** by the addition of an acrylic acid and adding a coupling reagent such as dicyclohexylcarbodi-imide (DCC), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU), or EEDQ. To the acrylamide **N** is then added an excess of trimethylsilyl diazomethane in a solvent such as ethyl acetate or dichloromethane. The resulting dihydropyrazole **O** may then be treated with an isocyanate, in the presence of base such as pyridine or triethylamine, in a solvent such as dichloromethane, to afford compound **P**.

15


Scheme 5.1



-132-




An alternative approach to molecules wherein X^{II} in (c.f., Formulas II-X), is N and “----” is a bond is provided in **Scheme 5.2**. The chemistry for the preparation of compounds Q-T is similar to that described in **Scheme 5.1**. Compound T may be converted to an invention compound by a Suzuki coupling with a boronic acid, although other coupling conditions commonly known to the skilled practitioner may also be used. This route is particularly useful for compounds containing a biaryl A-B group. In situations where a *tert*-butylsulphonamide is present in B, which occurs in some of the particularly preferred compounds, the sulphonamide may be formed by stirring with trifluoroacetic acid for 16 h.

Scheme 5.2


-133-

Another approach to the preparation of invention compounds wherein X^{II}



in (c.f., Formulas II-X), is N and "—" is a bond is provided in **Scheme 5.3**. The synthetic routes commences with formation of acrylamide, **U** as disclosed in **Scheme 5.1**. A diazoacetate such as ethyl diazoacetate is then mixed with acrylamide **U** to afford the appropriately substituted dihydropyrazole **V**. Similar chemistry to that described in **Scheme 5.1** allows conversion of **V** to **W**. Reduction of the ester moiety in **W** is then achieved using a reducing agent such as super-hydride to afford the hydroxyl containing invention compound **X**.

10


Scheme 5.3

Alternatively, invention compounds wherein X^{II} in (c.f., Formulas II-X), is N and "—" is a bond is provided in **Scheme 5.4**. Here,

-134-

- the acrylic ester **Y** is reacted with a diazo compound to form the dihydropyrazole **Z**. Addition of a chloroformate affords compounds of formula **AA**. Deprotection of the ester is achieved by acid, such as TFA or hydrogen chloride to afford compounds of formula **BB**. Addition of an aniline with a coupling reagent such as EEDQ and base affords compounds of formula **CC**. Deprotection of the carbamate with Pd/C and hydrogen affords compounds of formula **DD** which may then be reacted with isocyanates to afford invention compounds of formula **FF**.

Scheme 5.4



10

Other methods for making compounds of the invention or variations of the above Schemes are provided in the Examples section.

-135-

Not all compounds of the invention falling into a given class may be compatible with some of the reaction conditions described. Such restrictions are readily apparent to those skilled in the art of organic synthesis, and alternative methods must then be used.

5

Some of the compounds of Formula I are capable of further forming pharmaceutically acceptable acid-addition and/or base salts. All of these forms are within the scope of the present invention. Thus, pharmaceutically acceptable acid addition salts of the compounds of Formula I include salts derived from nontoxic
10 inorganic acids such as hydrochloric, nitric, phosphoric, sulfuric, hydrobromic, hydriodic, hydrofluoric, phosphorous, and the like, as well as the salts derived from nontoxic organic acids, such as aliphatic mono- and dicarboxylic acids, phenyl-substituted alkanoic acids, hydroxy alkanoic acids, alkanedioic acids, aromatic acids, aliphatic and aromatic sulfonic acids, etc. Such salts thus include
15 sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, nitrate, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, acetate, trifluoroacetate, propionate, caprylate, isobutyrate, oxalate, malonate, succinates suberate, sebacate, fumarate, maleate, mandelate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, phthalate, benzensulfonate,
20 toluenesulfonate, phenylacetate, citrate, lactate, maleate, tartrate, methanesulfonate, and the like. Also contemplated are salts of amino acids such as arginate and the like and gluconate, galacturonate (see, for example, Berge, S.M. et. al., "Pharmaceutical Salts," *Journal of Pharmaceutical Science*, 1977;66:1-19).

25 The acid addition salt of said basic compounds are prepared by contacting the free base form with a sufficient amount of the desired acid to produce the salt in the conventional manner.

Pharmaceutically acceptable base addition salts are formed with metals or
30 amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are N,N'-dibenzylethylenediamine, chloroprocaine,

-136-

choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methyl glucamine, and procaine (see, for example, Berge, S.M., supra., 1977).

The base addition salts of said acidic compounds are prepared by
5 contacting the free acid form with a sufficient amount of the desired base to produce the salt in the conventional manner.

Examples of pharmaceutically acceptable, non-toxic esters of the compounds of this invention include C₁-C₆ alkyl esters wherein the alkyl group is
10 a straight or branched chain. Acceptable esters also include C₅-C₇ cycloalkyl esters as well as arylalkyl esters such as, but not limited to benzyl. C₁-C₄ alkyl esters are preferred. Esters of the compounds of the present invention may be prepared according to conventional methods.

15 Examples of pharmaceutically acceptable, non-toxic amides of the compounds of this invention include amides derived from ammonia, primary C₁-C₆ alkyl amines and secondary C₁-C₆ dialkyl amines wherein the alkyl groups are straight or branched chain. In the case of secondary amines, the amine may also be in the form of a 5- or 6-membered heterocycle containing one nitrogen
20 atom. Amides derived from ammonia, C₁-C₃ alkyl primary amines and C₁-C₂ dialkyl secondary amines are preferred. Amides of the compounds of the invention may be prepared according to conventional methods.

Certain of the compounds of the present invention can exist in unsolvated
25 forms as well as solvated forms, including hydrated forms. In general, the solvated forms, including hydrated forms, are equivalent to unsolvated forms and are intended to be encompassed within the scope of the present invention.

Certain of the compounds of the present invention possess one or more
30 chiral centers and each center may exist in the *R*(D) or *S*(L) configuration. The

-137-

present invention includes all enantiomeric and epimeric forms, as well as the appropriate mixtures thereof.

The compounds of Formula I can be formulated as pharmaceutical
5 compositions and administered to a mammalian host, such as a human patient in a variety of forms adapted to the chosen route of administration, i.e., orally or parenterally, by intravenous, intramuscular, topical or subcutaneous routes.

Thus, the present compounds may be systemically administered, e.g.,
10 orally, in combination with a pharmaceutically acceptable vehicle such as an inert diluent or an assimilable edible carrier. They may be enclosed in hard or soft shell gelatin capsules, may be compressed into tablets, or may be incorporated directly with the food of the patient's diet. For oral therapeutic administration, the active compound may be combined with one or more excipients and used in the form of
15 ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.1% of active compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 60% of the weight of a given unit dosage form. The amount of active compound in
20 such therapeutically useful compositions is such that an effective dosage level will be obtained.

The tablets, troches, pills, capsules, and the like may also contain the following: binders such as gum tragacanth, acacia, corn starch or gelatin;
25 excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, fructose, lactose or aspartame or a flavoring agent such as peppermint, oil of wintergreen, or cherry flavoring may be added. When the unit dosage form is a capsule, it may contain, in addition to
30 materials of the above type, a liquid carrier, such as a vegetable oil or a polyethylene glycol. Various other materials may be present as coatings or to otherwise modify the physical form of the solid unit dosage form. For instance,

-138-

tablets, pills, or capsules may be coated with gelatin, wax, shellac or sugar and the like. A syrup or elixir may contain the active compound, sucrose or fructose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavoring such as cherry or orange flavor. Of course, any material used in
5 preparing any unit dosage form should be pharmaceutically acceptable and substantially non-toxic in the amounts employed. In addition, the active compound may be incorporated into sustained-release preparations and devices.

The active compound may also be administered intravenously or
10 intraperitoneally by infusion or injection. Solutions of the active compound or its salts can be prepared in water, optionally mixed with a nontoxic surfactant. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, triacetin, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of
15 microorganisms.

The pharmaceutical dosage forms suitable for injection or infusion can include sterile aqueous solutions or dispersions or sterile powders comprising the active ingredient which are adapted for the extemporaneous preparation of sterile
20 injectable or infusible solutions or dispersions, optionally encapsulated in liposomes. In all cases, the ultimate dosage form must be sterile, fluid and stable under the conditions of manufacture and storage. The liquid carrier or vehicle can be a solvent or liquid dispersion medium comprising, for example, water, ethanol, a polyol (for example, glycerol, propylene glycol, liquid polyethylene glycols, and
25 the like), vegetable oils, nontoxic glyceryl esters, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the formation of liposomes, by the maintenance of the required particle size in the case of dispersions or by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal
30 agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, buffers or sodium chloride. Prolonged absorption of the

-139-

injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions are prepared by incorporating the active compound in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filter sterilization. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze drying techniques, which yield a powder of the active ingredient plus any additional desired ingredient present in the previously sterile-filtered solutions.

For topical administration, the present compounds may be applied in pure form, i.e., when they are liquids. However, it will generally be desirable to administer them to the skin as compositions or Formulations, in combination with a dermatologically acceptable carrier, which may be a solid or a liquid.

Useful solid carriers include finely divided solids such as talc, clay, microcrystalline cellulose, silica, alumina and the like. Useful liquid carriers include water, alcohols or glycols or water-alcohol/glycol blends, in which the present compounds can be dissolved or dispersed at effective levels, optionally with the aid of non-toxic surfactants. Adjuvants such as fragrances and additional antimicrobial agents can be added to optimize the properties for a given use. The resultant liquid compositions can be applied from absorbent pads, used to impregnate bandages and other dressings, or sprayed onto the affected area using pump-type or aerosol sprayers.

Thickeners such as synthetic polymers, fatty acids, fatty acid salts and esters, fatty alcohols, modified celluloses or modified mineral materials can also be employed with liquid carriers to form spreadable pastes, gels, ointments, soaps, and the like, for application directly to the skin of the user.

-140-

Examples of useful dermatological compositions which can be used to deliver the compounds of Formula I to the skin are known to the art; for example, see Jacquet et al. (U.S. Pat. No. 4,608,392), Geria (U.S. Pat. No. 4,992,478), Smith et al. (U.S. Pat. No. 4,559,157) and Wortzman (U.S. Pat. No. 4,820,508).

5

Useful dosages of the compounds of Formula I can be determined by comparing their in vitro activity, and in vivo activity in animal models. Methods for the extrapolation of effective dosages in mice, and other animals, to humans are known to the art; for example, see U.S. Pat. No. 4,938,949.

10

Generally, the concentration of the compound(s) of Formula I in a liquid composition, such as a lotion, will be from about 0.1-25 wt-%, preferably from about 0.5-10 wt-%. The concentration in a semi-solid or solid composition such as a gel or a powder will be about 0.1-5 wt-%, preferably about 0.5-2.5 wt-%.

15

The amount of the compound, or an active salt or derivative thereof, required for use in treatment will vary not only with the particular salt selected but also with the route of administration, the nature of the condition being treated and the age and condition of the patient and will be ultimately at the discretion of the attendant physician or clinician.

20

The compounds of the present invention can be administered to a patient at dosage levels in the range of about 0.1 to about 2,000 mg per day. For a normal human adult having a body weight of about 70 kilograms, a dosage in the range of about 0.01 to about 10 mg per kilogram of body weight per day is preferable. However, the specific dosage used can vary. For example, the dosage can depend on a numbers of factors including the requirements of the patient, the severity of the condition being treated, and the pharmacological activity of the compound being used. The determination of optimum dosages for a particular patient is well-known to those skilled in the art.

25

30

-141-

Ideally, the active ingredient should be administered to achieve peak plasma concentrations of the active compound of from about 0.5 nM (nonmolar) to about 75 μ M, preferably, about 1 nM to 50 μ M, most preferably, about 0.1 to about 5 μ M. This may be achieved, for example, by the intravenous injection of a
5 0.05 to 5% solution of the active ingredient, optionally in saline, or orally administered as a bolus containing about 10-500 mg of the active ingredient. Desirable blood levels may be maintained by multiple oral dosing, or continuous infusion to provide about 0.01-5.0 mg/kg/hr or by intermittent infusions containing about 0.4-15 mg/kg of the active ingredient(s).

10

The desired dose may conveniently be presented in a single dose or as divided doses administered at appropriate intervals, for example, as two, three, four or more sub-doses per day. The sub-dose itself may be further divided, e.g., into a number of discrete loosely spaced administrations; such as multiple
15 inhalations from an insufflator or by application of a plurality of drops into the eye.

The following examples illustrate the various embodiments of the present invention. Those skilled in the art will recognize many variations that are within
20 the spirit of the present invention and scope of the claims.

BIOLOGICAL ASSAYS

The invention compounds have demonstrated factor Xa inhibitory activity in the standard assays commonly employed by those skilled in the art.

25

A. Determination of Factor Xa IC₅₀

The ability of compounds to act as inhibitors of human factor Xa catalytic activity is assessed by determination of that concentration of test substance that inhibits by 50% (IC₅₀) the ability of human factor Xa to cleave the chromogenic
30 substrate S2765 (N-CBz-D-Arg-L-Gly-L-Arg-p-nitroanilide. 2HCl, DiaPharma). The IC₅₀ was determined at 3pM and 30pM concentrations human factor Xa

-142-

(Enzyme Research Laboratories). These concentrations were achieved by diluting a 21.087 μM stock solution of factor Xa in the appropriate amount of a buffer comprising 10 μM HEPES, 150 μM NaCl, 0.1% BSA, pH 7.4 (HBSA buffer).

Accordingly, 5 μL of the compound to be tested in DMSO (2% final) is added to
 5 the factor Xa/buffer solution and incubated for 60 minutes at room temperature.

The IC_{50} is determined by monitoring the increase in absorbance at 390 nm excitation, 460 nm emission, with a 455 nm cutoff, in a fluorometric plate reader. Results of the IC_{50} at 3 pM and 30 pM enzyme concentrations are

10 provided in Table 1.

Table 1

EXAMPLE	FXA 3pM IC_{50} (μM)	FXA 30 pM IC_{50} (μM)
(R)-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]	0.0199	0.0270
Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]		0.0355
Pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]		0.2568
Pyrrolidine-1,2-dicarboxylic acid 1-[(2,4-difluoro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]		9.1175
Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-p-tolylamide	0.3890	
Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-[(4-methoxy-phenyl)-amide]	0.8730	
Pyrrolidine-1,2-dicarboxylic acid 1-[(4-bromo-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]	0.0333	

Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-phenylamide	0.9510	
Pyrrolidine-1,2-dicarboxylic acid 1-[(3,4-difluorophenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]	0.6460	
Pyrrolidine-1,2-dicarboxylic acid 1-[(3-fluoro-4-methyl-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]	0.5150	
Pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]	0.0804	
Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]	0.0172	
Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[5-(2-sulfamoyl-phenyl)-pyridin-2-yl]-amide]	0.0730	
Pyrrolidine-1,2-dicarboxylic acid 2-[(3-chloro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(5-chloro-pyridin-2-yl)-amide]	0.1010	
Pyrrolidine-1,2-dicarboxylic acid 2-[(3-chloro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide]	0.0166	
Pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[5-(2-sulfamoyl-phenyl)-pyridin-2-yl]-amide]	0.2770	
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]	0.0007	
Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-amide]	0.1560	
Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-biphenyl-4-yl)-amide]	0.0273	

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3-methyl-biphenyl-4-yl)-amide]	0.0591	
(2R,4R)-4-methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]	0.000374833	
(2R)-4-oxo-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]	0.007596667	
(2R,4S)-4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]	0.004443333	
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-p-tolylamide	0.01065	
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(4-fluoro-phenyl)-amide]	0.0028575	
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]	0.0071125	
Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3-trifluoromethyl-biphenyl-4-yl)-amide]	0.0806	
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]		0.0014
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2-oxo-piperidin-1-yl)-phenyl]-amide]	0.0025	
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]	0.0025	
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-		0.0041

oxo-piperidin-1-yl)-phenyl]-amide}		
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide}		0.0149
(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide}		0.0084
(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide}		0.0045
(2R,4R)-4-Propoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide}		0.0008
(2R,4R)-4-Propoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide}		0.0020
(2R,4R)-4-Cyano-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide}		0.0050
(2R,4R)-4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide}		0.0510
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(2-oxo-azetidin-1-yl)-phenyl]-amide}		0.0110
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{2-fluoro-4-(2-oxo-azetidin-1-yl)-phenyl]-amide}		0.1223
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{2-fluoro-4-(2-oxo-pyrrolidin-1-yl)-phenyl]-amide}		0.0040
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(2-oxo-azepan-1-yl)-phenyl]-amide}		0.0294

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(2-oxo-azepan-1-yl)-phenyl]-amide}		0.0041
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide}		0.0300
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide}		0.0013
(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide}		0.0004
(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide}		0.00062
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-phenyl]-amide}		0.231
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(2,3-dimethyl-5-oxo-2,5-dihydro-pyrazol-1-yl)-phenyl]-amide}		0.2560
4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(2-hydroxymethyl-pyrrolidin-1-yl)-phenyl]-amide}		0.0110
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(pyrrolidin-1-yl)-phenyl]-amide}		0.0640
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(pyrazol-1-yl)-phenyl]-amide}	0.0344	
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-pyrazol-1-yl-phenyl]-amide}		0.0434
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-[1,2,4]triazol-1-yl-		0.0619

-147-

phenyl)-amide]		
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-[1,2,3]triazol-2-yl-phenyl)-amide]		0.1903
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-pyrazol-1-yl-phenyl)-amide]		0.041000
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-pyrazol-1-yl-phenyl)-amide]		0.032000
(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-pyrazol-1-yl-phenyl)-amide]		0.029000
(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-pyrazol-1-yl-phenyl)-amide]		0.018000
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-methyl-pyrazol-1-yl)-phenyl]-amide]		0.082000
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-methyl-pyrazol-1-yl)-phenyl]-amide]		0.021000
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-methyl-pyrazol-1-yl)-phenyl]-amide]		0.009600
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(5-methyl-pyrazol-1-yl)-phenyl]-amide]		0.015000
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3,5-dimethyl-pyrazol-1-yl)-phenyl]-amide]		0.049000
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3,5-dimethyl-pyrazol-1-yl)-phenyl]-amide]		0.020000
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3,5-dimethyl-pyrazol-1-yl)-2-fluoro-phenyl]-amide]		0.010000

(2R,4S)-4-Amino-pyrrolidine-1,2-dicarboxylic acid 1- [(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'- methanesulfonyl-biphenyl-4-yl)-amide]		0.010000
(2R,4R)-4-Amino-pyrrolidine-1,2-dicarboxylic acid 1- [(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'- methanesulfonyl-biphenyl-4-yl)-amide]		0.025925
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl- biphenyl-4-yl)-amide]		0.000770
(2R)-4-Hydroxyimino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'- methanesulfonyl-biphenyl-4-yl)-amide]		0.002108
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'- methylsulfamoyl-biphenyl-4-yl)-amide]		0.001922
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-dimethylsulfamoyl- 3-fluoro-biphenyl-4-yl)-amide]		0.001188
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'- methanesulfonyl-biphenyl-4-yl)-amide]		0.001021
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl- biphenyl-4-yl)-amide]		0.000491
(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1- [(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'- methanesulfonyl-biphenyl-4-yl)-amide]		0.000701
(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1- [(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl- biphenyl-4-yl)-amide]		0.000276
(2R,4S)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'- methanesulfonyl-biphenyl-4-yl)-amide]		0.001685
(2R,4R)-4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1- [(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'- methanesulfonyl-biphenyl-4-yl)-amide]		0.001200
(2R)-4,4-Difluoro-pyrrolidine-1,2-dicarboxylic acid 1- [(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'- methanesulfonyl-biphenyl-4-yl)-amide]		0.005360

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(4-fluoro-phenyl)-amide]		0.001873
(2R,4R)-4-Acetylamino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]		0.010545
(2R,4R)-4-Methanesulfonylamino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]		0.001735
(2R,4S)-4-(1H-Tetrazol-5-yl)-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]		0.038850
(2R,4S)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]		0.011770
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-methyl-imidazol-1-yl)-phenyl]-amide]		0.006210
(2R,4R)-4-Cyano-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]		0.055475
(2R,4R)-4-(1H-Tetrazol-5-yl)-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]		0.007973
(2R,4R)-4-Trifluoromethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]		0.032950
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-cyano-3-fluoro-biphenyl-4-yl)-amide]		0.001950
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[(2'-aminomethyl-3-fluoro-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide]		0.001633
(2R,4R)-4'-[[1-(4-Chloro-phenyl)carbamoyl]-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3'-fluoro-biphenyl-2-carboxylic acid methyl ester		0.007112
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,5-dihydro-pyrrole-1-carbonyl)-phenyl]-amide]		0.00029

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-2-fluoro-phenyl)-amide]		0.015
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-phenyl]-amide]		0.0081
Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-phenyl]-amide]		0.081
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2-methyl-pyrrolidine-1-carbonyl)-phenyl]-amide]		0.024
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(ethyl-methyl-carbamoyl)-phenyl]-amide]		0.037
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-phenyl)-amide]		0.010
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2R-methyl-pyrrolidine-1-carbonyl)-phenyl]-amide]		0.011
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2S-methyl-pyrrolidine-1-carbonyl)-phenyl]-amide]		0.082
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(pyrrolidine-1-carbonyl)-phenyl]-amide]		0.0096
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-2-pyrrolidin-1-yl-phenyl]-amide]		0.035
(2R,4R)-4-[[1-(4-Chloro-phenylcarbamoyl)-4-hydroxy-pyrrolidine-2-carbonyl]-amino]-3-pyrrolidin-1-yl-benzoic acid methyl ester		0.020
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(azetidine-1-carbonyl)-phenyl]-amide] 1-[(4-chloro-phenyl)-amide]		0.061
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(pyrrolidine-1-carbonyl)-phenyl]-amide]		0.084

-151-

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-phenyl]-amide]		0.081
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(4-dimethylcarbamoyl-phenyl)-amide]		0.044
(2R,4R)-4-[[1-(4-Chloro-phenylcarbamoyl)-4-hydroxy-pyrrolidine-2-carbonyl]-amino]-3-dimethylamino-benzoic acid methyl ester		0.029
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-phenyl]-amide]		0.0073
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-phenyl)-amide]		0.0049
(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-phenyl]-amide]		0.0057
(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-phenyl)-amide]		0.0039
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(pyrrolidine-1-carbonyl)-phenyl]-amide]		0.0062
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-2-fluoro-phenyl)-amide]		0.0061
(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(pyrrolidine-1-carbonyl)-phenyl]-amide]		0.0044
(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-2-fluoro-phenyl)-amide]		0.0029
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-methyl-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]		0.0052

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-methyl-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]		0.0020
Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-quinolin-8-yl-phenyl)-amide]		0.066
Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3,5-difluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]		0.0098
Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-2-methyl-biphenyl-4-yl)-amide]		0.018
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-biphenyl-4-yl)-amide]		0.00073
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfinyl-biphenyl-4-yl)-amide]		0.0039
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-methyl-2'-methylsulfanyl-biphenyl-4-yl)-amide]		0.053
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]		0.0012
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3-methyl-biphenyl-4-yl)-amide]		0.00068
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3-trifluoromethyl-biphenyl-4-yl)-amide]		0.0015
5-Methyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]		0.005623
3-Hydroxymethyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]		0.011267
4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]		0.008015

-153-

(R) 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]		0.007738
4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl)-amide]		0.044850
4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]		0.008718
1-(4-Chloro-phenylcarbamoyl)-5-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-4,5-dihydro-1H-pyrazole-3-carboxylic acid ethyl ester		0.040850
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methoxy-biphenyl-4-yl)-amide]		0.058900
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-hydroxy-biphenyl-4-yl)-amide]		0.074100
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-iodo-phenyl)-amide]		0.058575
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl)-amide]		0.000404
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl)-amide]		0.001153
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-(3-hydroxy-2-oxo-piperidin-1-yl)-phenyl)-amide]		0.008498
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-(2-oxo-tetrahydro-pyrimidin-1-yl)-phenyl)-amide]		0.007665
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-(2-oxo-imidazolidin-1-yl)-phenyl)-amide]		0.036425
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-(2-oxo-oxazolidin-3-yl)-phenyl)-amide]		0.045525

(2R,4R)-1-(4-{{1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl}-amino}-3-fluoro-phenyl)-2-oxo-piperidine-3-carboxylic acid ethyl ester		0.011000
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(3-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl}-amide}		0.000862
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{2-fluoro-4-(3-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl}-amide}		0.005075
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{4-(3-methyl-2-oxo-2H-pyridin-1-yl)-phenyl}-amide}		0.000383
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{4-(3-methyl-2-oxo-2H-pyridin-1-yl)-phenyl}-amide}		0.001793
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{(4-morpholin-4-yl)-phenyl}-amide}		0.043050
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{4-(2-methyl-5-oxo-pyrrolidin-1-yl)-phenyl}-amide}		0.002303
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{4-(2-methyl-5-oxo-pyrrolidin-1-yl)-phenyl}-amide}		0.011975
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(4-methyl-2-oxo-2H-pyridin-1-yl)-phenyl}-amide}		0.002757
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{2-fluoro-4-(4-methyl-2-oxo-2H-pyridin-1-yl)-phenyl}-amide}		0.012100
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(5-methyl-2-oxo-2H-pyridin-1-yl)-phenyl}-amide}		0.000743
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{2-fluoro-4-(5-methyl-2-oxo-2H-pyridin-1-yl)-phenyl}-amide}		0.003500

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(4-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide]		0.002170
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(4-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide]		0.004967
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide]		0.00025
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(3-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide]		0.00148
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(6-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide]		0.001210
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(5-chloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide]		0.00058
4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(4-tert-butyl-phenyl)-amide] 1-[(4-chloro-phenyl)-amide]		0.0260
4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3,5'-difluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]		0.0006
4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)-amide]		0.0650
4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide]		0.0070
4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(2,3-dimethyl-5-oxo-2,5-dihydro-pyrazol-1-yl)-phenyl]-amide]		0.0580
4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,3-dimethyl-5-oxo-2,5-dihydro-pyrazol-1-yl)-phenyl]-amide]		0.0800
4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide]		0.0050

-156-

4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide]-TFA Salt		0.0900
4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide]		0.0170
4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide]		0.0030
4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-(6-oxo-6H-pyridazin-1-yl)-phenyl)-amide]		0.0420

Similarly, Table 2 provides the percent inhibition at 3 pM concentration of factor Xa and compound concentrations of 4 μ M and 1 μ M.

5

Table 2

Name	Fxa 3PM %Inhibition at 4 μ M	Fxa 3PM %Inhibition at 1 μ M
Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-[(4-isopropyl-phenyl)-amide]	19.22	
Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-[(4-trifluoromethyl-phenyl)-amide]	25.1	
Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-[(3-methoxy-phenyl)-amide]	36.21	
Pyrrolidine-1,2-dicarboxylic acid 1-[(4-ethyl-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]	38.14	
Pyrrolidine-1,2-dicarboxylic acid 1-[(3-acetyl-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]	21.04	
Pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-2-methyl-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]	19.84	
Pyrrolidine-1,2-dicarboxylic acid 1-[(2-fluoro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]	37.21	
Pyrrolidine-1,2-dicarboxylic acid 1-[(3-chloro-4-fluoro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]	41.07	

-157-

Pyrrolidine-1,2-dicarboxylic acid 1-[(3-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]	55.97	
Pyrrolidine-1,2-dicarboxylic acid 1-[(4-cyano-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]	51.86	
Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-2-methyl-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]	38.88	
Pyrrolidine-1,2-dicarboxylic acid 1-[(3,4-dichloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]	41.71	
Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-[(3-trifluoromethyl-phenyl)-amide]	21.61	
Pyrrolidine-1,2-dicarboxylic acid 1-[(4-dimethylamino-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]	19.93	
(2S)-Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]		46.5
Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-cyano-phenyl)-amide]		16.58
Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(1H-tetrazol-5-yl)-phenyl]-amide]		36.64
(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-azetidin-1-yl)-phenyl]-amide]		54
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-[1,2,3]triazol-1-yl-phenyl)-amide]		26% at 1uM
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(4-acetylamino-phenyl)-amide] 1-[(4-chloro-phenyl)-amide]		0.0793
(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(cyclopentanecarbonylamino)-phenyl]-amide]		36% at 1uM
4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-pyrimidin-5-yl-phenyl)-amide]		32% at 1 uM


-158-

The foregoing biological tests establish that the compounds of the present invention are potent inhibitors of the Factor Xa. Accordingly, the compounds of the present invention are useful in pharmaceutical formulations for preventing and treating thrombotic disorders. Such disorders include venous thrombosis, deep vein thrombosis, thrombophlebitis, arteriolembolism, coronary and cerebral arterial thrombosis, cerebral embolism, kidney embolism, pulmonary embolism, first or recurrent myocardial infarction, unstable angina, cerebral infarction, stroke, and atherosclerosis.

To further assist in understanding the present invention, the following non-limiting examples of such factor Xa inhibitory compounds are provided. The following examples, of course, should not be construed as specifically limiting the present invention, variations presently known or later developed, which would be within the purview of one skilled in the art and considered to fall within the scope of the present invention as described herein. Preferred synthetic routes for intermediates involved in the synthesis as well as the resulting anti-thrombotic compounds of the present invention follow.

Example 1

(R)-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]



Step 1: (R)-2-(4-Bromo-2-fluoro-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid 9H-fluoren-9-ylmethyl ester

-159-

Fmoc-D-Pro (1.2 g, 3.56 mmol) was suspended in 10 mL dry dichloromethane (DCM), cooled in an ice water bath, and then thionyl chloride (0.78 mL, 10.7 mmol) was added slowly. After addition was complete, the ice water bath was removed and the reaction was stirred at ambient temperature overnight. The solution was concentrated to dryness. The oil was redissolved in 20 mL dry DCM under an Ar atmosphere, and 4-bromo-2-fluoroaniline (0.81 g, 4.3 mmol) was added, followed by dry pyridine (0.87 mL, 10.7 mmol). The mixture was stirred at ambient temperature for 4 hours, then concentrated. The residue was redissolved in 100 mL EtOAc, washed with 1M HCl (3x50 mL), brine (2x50 mL), dried with MgSO₄, filtered and concentrated to yield the title product as a sticky oil. (1.8 g, 99%) APCI (AP+): 509.2, 511.2 (M+H)⁺.

Step 2: (R)-Pyrrolidine-2-carboxylic acid (2'-*tert*-butylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide

(R)-2-(4-Bromo-2-fluoro-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid 9H-fluoren-9-ylmethyl ester (1.78 g, 3.49 mmol), 2-(*N-tert*-butyl)phenylsulfonamide boronic acid (1.35 g, 5.24 mmol), and tetrabutylammonium bromide (60 mg, 0.17 mmol) were combined in 40 mL toluene. 3.5 mL of an aqueous 2M sodium carbonate solution was then added, followed by tetrakis(triphenylphosphine) palladium(0) (200 mg, 0.17 mmol). The resulting solution was heated at reflux overnight, and cooled to room temperature, partitioned between EtOAc (250 mL) and water (150 mL). The combined organic layers were washed with water (2x100 mL), brine, dried with MgSO₄, filtered and concentrated. Purification by column chromatography on silica gel eluting with 10% EtOAc in hexanes, then EtOAc, and then to MeOH/NH₄OH/EtOAc (5:1:94) provided the title compound as an oil. (1.03 g, 70%) APCI (AP+): 420.3 (M+H)⁺.

Step 3: (R)-pyrrolidine-1,2-dicarboxylic acid 2-[(2'-*tert*-butylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide]

(R)-Pyrrolidine-2-carboxylic acid (2'-*tert*-butylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide (0.25 g, 0.6 mmol) and 4-chlorophenylisocyanate (90 mg,

-160-

0.6 mmol) were combined in 10 mL DCM and stirred at ambient temperature for one hour. The solution was concentrated. The residue was purified on a silica gel column eluting with 50% EtOAc in hexanes to provide the title compound as a white foam. (0.225 g, 66%) APCI (AP-): 571.3, 573.3 (M-H)⁻.

5

Step 4: (R)-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]


(R)-pyrrolidine-1,2-dicarboxylic acid 2-[(2'-*tert*-butylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide] (0.22 g, 0.38 mmol) was stirred in 20 mL trifluoroacetic acid at ambient temperature overnight. The solution was concentrated, then purified on a silica gel column eluting with 50% EtOAc in hexanes, then 75% EtOAc in hexanes. The pure fractions were combined and concentrated, then redissolved in acetonitrile/water and lyophilized to yield title compound as a white solid. (0.15 g, 76%) APCI (AP-): 515.2, 517.2 (M-H)⁻.

15

Example 2

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]

20



Step 1: Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

DL-proline (10 g, 86.9 mmol) was dissolved in 120 mL of THF: H₂O (2:1), and 87 mL of a 2M NaOH solution was added, followed by Boc₂O (24.6 g, 113 mmol). The reaction mixture was stirred at ambient temperature overnight.

25

-161-

The THF was removed in vacuo. The remaining aqueous mixture was acidified to pH 3 with aqueous citric acid, extracted twice with EtOAc. The combined organic layers were washed with water, brine, dried with MgSO₄, filtered and concentrated to yield the title compound as a white solid. (18.7 g, quant.) APCI (AP-): 214.1 (M-H)⁺.

Step 2: 2-(2'-*tert*-Butylsulfamoyl-3-fluoro-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (1.6 g, 7.4 mmol) was dissolved in 40 mL dry diethyl ether under Ar. The solution was cooled in an ice water bath as dry pyridine (3 mL, 37.2 mmol) was added, followed by the dropwise addition of oxalyl chloride (1.6 mL, 18.6 mmol). A precipitate formed immediately. The reaction was stirred vigorously at 0°C for one hour, then at ambient temperature for one hour. 50 mL diethyl ether were added, and the mixture was filtered. The filtered off solids were washed with diethyl ether. The filtrates were combined and concentrated to give an off-white oil. The oil was redissolved in 40 mL dry DCM under Ar, cooled to 0°C, and 3 mL of pyridine were added, followed by 4'-amino-3'-fluoro-biphenyl-2-sulfonic acid *tert*-butylamide (2 g, 6.2 mmol). The reaction was allowed to warm to ambient temperature, stirred for 3 hours then concentrated. The concentrate was redissolved in 250 mL EtOAc, washed with 10% citric acid (2x100 mL), water (3x100 mL), brine (100 mL), dried with MgSO₄, filtered and concentrated. The concentrate was purified on a silica gel column eluted with 30% moving to 50% EtOAc in hexanes. Combined and concentrated pure fractions to yield title compound as a white solid. (1.4 g, 42%) APCI (AP-): 518.2 (M-H)⁺.

Step 3: Pyrrolidine-2-carboxylic acid (2'-*tert*-butylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide

2-(2'-*tert*-Butylsulfamoyl-3-fluoro-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (1.3 g, 2.5 mmol) was dissolved in 40 mL DCM, cooled in an ice bath, added 10 mL trifluoroacetic acid, stirred at 0°C for 30

-162-

minutes then at ambient temperature for one hour. Concentrated solution, redissolved in EtOAc (150 mL), washed with saturated NaHCO₃ (2x100 mL), water (100 mL), brine (100 mL), dried with MgSO₄, filtered and concentrated to yield title compound as a white solid. (1 g, 91%) APCI (AP⁻): 418.2 (M-H)⁻.

5

Step 4: Pyrrolidine-1,2-dicarboxylic acid 2-[(2'-*tert*-butylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide]

Pyrrolidine-2-carboxylic acid (2'-*tert*-butylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide (0.3 g, 0.72 mmol) was dissolved in 10 mL dry THF under Ar, added 4-chlorophenylisocyanate (0.11 g, 0.72 mmol) and stirred at ambient temperature for one hour. Concentrated solution to yield title compound. (0.42 g, quant.)
APCI
(AP⁻): 571.2, 573.2 (M-H)⁻.

15 **Step 5: Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]**


Pyrrolidine-1,2-dicarboxylic acid 2-[(2'-*tert*-butylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide] (0.4 g, 0.7 mmol) was stirred in 10 mL trifluoroacetic acid at ambient temperature overnight, concentrated solution, then purified on a silica gel column eluted with 50% moving to 75% EtOAc in hexanes. Combined and concentrated pure fractions, redissolved in acetonitrile/water and lyophilized to yield title compound as a white fluffy powder. (0.26 g, 72%) APCI (AP⁻): 515.1, 517.2 (M-H)⁻.

25

Example 3

Pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]

-163-



Step 1: Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

DL-proline (10 g, 86.9 mmol) was dissolved in 120 mL of THF: H₂O (2:1), added 87 mL of a 2M NaOH solution followed by Boc₂O (24.6 g, 113 mmol). Stirred at ambient temperature overnight. Removed THF in vacuo, acidified water to pH 3 with citric acid, extracted twice with EtOAc, washed organics with water, brine, dried with MgSO₄, filtered and concentrated to yield title compound as a white solid. (18.7 g, quant.) APCI (AP-): 214.1 (M-H)⁻.

10

Step 2: 2-(2'-*tert*-Butylsulfamoyl-3-fluoro-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (1.6 g, 7.4 mmol) was dissolved in 40 mL dry diethyl ether under Ar, cooled in an ice water bath, added dry pyridine (3 mL, 37.2 mmol) followed by the dropwise addition of oxalyl chloride (1.6 mL, 18.6 mmol). A precipitate forms immediately. The reaction was stirred vigorously at 0°C for one hour, then at ambient temperature for one hour. Added 50 mL diethyl ether, filtered off solids washing with diethyl ether. Concentrated the filtrates to an off-white oil. Redissolved oil in 40 mL dry DCM under Ar, cooled to 0°C, added 3 mL pyridine followed by 4'-amino-3'-fluoro-biphenyl-2-sulfonic acid *tert*-butylamide (2 g, 6.2 mmol). The reaction was allowed to warm to ambient temperature, stirred for 3 hours then concentrated. Redissolved in 250 mL EtOAc, washed with 10% citric acid (2x100 mL), water (3x100 mL), brine (100 mL), dried with MgSO₄, filtered and concentrated.

25 Purified on a silica gel column eluted with 30% moving to 50% EtOAc in

-164-

hexanes. Combined and concentrated pure fractions to yield title compound as a white solid. (1.4 g, 42%) APCI (AP-): 518.2 (M-H)⁻.

Step 3: Pyrrolidine-2-carboxylic acid (2'-*tert*-butylsulfamoyl-3-fluoro-

5 biphenyl-4-yl)-amide

2-(2'-*tert*-Butylsulfamoyl-3-fluoro-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (1.3 g, 2.5 mmol) was dissolved in 40 mL DCM, cooled in an ice bath, added 10 mL trifluoroacetic acid, stirred at 0°C for 30 minutes then at ambient temperature for one hour. Concentrated solution, 10 redissolved in EtOAc (150 mL), washed with saturated NaHCO₃ (2x100 mL), water (100 mL), brine (100 mL), dried with MgSO₄, filtered and concentrated to yield title compound as a white solid. (1 g, 91%) APCI (AP-): 418.2 (M-H)⁻.

Step 3a: (5-Chloro-pyridin-2-yl)-carbamic acid 4-nitro-phenyl ester

15 2-Amino-5-chloropyridine (1.05 g, 8.2 mmol) was suspended in 15 mL dry DCM under Ar, added dry pyridine (0.66 mL, 8.2 mmol), cooled to 0°C and added 4-nitrophenol chloroformate. A white precipitate forms rapidly. Stirred at ambient temperature for 1 hour, filtered off solid, washed with water and then DCM. A white solid recovered which is consistent with title compound. (2.1 g, 20 88%) ¹H-NMR (D6-DMSO)

Step 4: Pyrrolidine-1,2-dicarboxylic acid 2-[(2'-*tert*-butylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide] 1-[(5-chloro-pyridin-2-yl)-amide]

Pyrrolidine-2-carboxylic acid (2'-*tert*-butylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide (245 mg, 0.83 mmol) and (5-Chloro-pyridin-2-yl)-carbamic acid 4-nitro-phenyl ester (0.35 g, 0.83 mmol) were combined in 10 mL dry DMF under Ar, added diisopropylethylamine (0.15 mL, 0.83 mmol) and heated at 80°C for 2 hours. Cooled solution, taken up in EtOAc (200 mL), washed with sat. K₂CO₃ (3x150 mL), sat. NaHCO₃ (3x150 mL), brine, dried with MgSO₄, filtered and

-165-

concentrated to yield title compound as a light pink solid. (0.45 g, 94%) APCI (AP-): 572.3, 574.3 (M-H)⁻.


Step 5: Pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]

Pyrrolidine-1,2-dicarboxylic acid 2-[(2'-*tert*-butylsulfamoyl)-3-fluoro-biphenyl-4-yl)-amide] 1-[(5-chloro-pyridin-2-yl)-amide] (0.45 g, 0.78 mmol) was stirred in 10 mL trifluoroacetic acid at ambient temperature for 4 hours. Reaction is sluggish. Heated to 80°C for 30 minutes. Cooled solution, concentrated, redissolved in 40 mL toluene and concentrated. Purified on a silica gel column eluted with 75% EtOAc in hexanes. Combined and concentrated pure fractions, redissolved in acetonitrile/water and lyophilized to yield title compound as a white fluffy powder. (0.277 g, 68%) APCI (AP-): 516.1, 518.1 (M-H)⁻.

15

Example 4

Pyrrolidine-1,2-dicarboxylic acid 1-[(2,4-difluoro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]



20

Step 1: Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

DL-proline (10 g, 86.9 mmol) was dissolved in 120 mL of THF: H₂O (2:1), added 87 mL of a 2M NaOH solution followed by Boc₂O (24.6 g, 113 mmol). Stirred at ambient temperature overnight. Removed THF in vacuo,

-166-

acidified water to pH 3 with citric acid, extracted twice with EtOAc, washed organics with water, brine, dried with MgSO₄, filtered and concentrated to yield title compound as a white solid. (18.7 g, quant.) APCI (AP⁻): 214.1 (M-H)⁻.

5 **Step 2: 2-(2'-*tert*-Butylsulfamoyl-3-fluoro-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester**

Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (1.6 g, 7.4 mmol) was dissolved in 40 mL dry diethyl ether under Ar, cooled in an ice water bath, added
10 dry pyridine (3 mL, 37.2 mmol) followed by the dropwise addition of oxalyl chloride (1.6 mL, 18.6 mmol). A precipitate forms immediately. The reaction was stirred vigorously at 0°C for one hour, then at ambient temperature for one hour. Added 50 mL diethyl ether, filtered off solids washing with diethyl ether. Concentrated the filtrates to an off-white oil. Redissolved oil in 40 mL dry DCM
15 under Ar, cooled to 0°C, added 3 mL pyridine followed by 4'-amino-3'-fluoro-biphenyl-2-sulfonic acid *tert*-butylamide (2 g, 6.2 mmol). The reaction was allowed to warm to ambient temperature, stirred for 3 hours then concentrated. Redissolved in 250 mL EtOAc, washed with 10% citric acid (2x100 mL), water (3x100 mL), brine (100 mL), dried with MgSO₄, filtered and concentrated.
20 Purified on a silica gel column eluted with 30% moving to 50% EtOAc in hexanes. Combined and concentrated pure fractions to yield title compound as a white solid. (1.4 g, 42%) APCI (AP⁻): 518.2 (M-H)⁻.

25 **Step 3: Pyrrolidine-2-carboxylic acid (2'-*tert*-butylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide**

2-(2'-*tert*-Butylsulfamoyl-3-fluoro-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (1.3 g, 2.5 mmol) was dissolved in 40 mL DCM, cooled in an ice bath, added 10 mL trifluoroacetic acid, stirred at 0°C for 30 minutes then at ambient temperature for one hour. Concentrated solution,
30 redissolved in EtOAc (150 mL), washed with saturated NaHCO₃ (2x100 mL),

-167-

water (100 mL), brine (100 mL), dried with MgSO_4 , filtered and concentrated to yield title compound as a white solid. (1 g, 91%) APCI (AP-): 418.2 (M-H)⁻.


Step 4: Pyrrolidine-1,2-dicarboxylic acid 1-[(2,4-difluoro-phenyl)-amide] 2-
5 [(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]

To a solution of pyrrolidine-2-carboxylic acid (2'-*tert*-butylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide (0.12 mmol/mL, 0.3 mmol) in 2.5 mL of THF was added 2,4-difluorophenylisocyanate. The reaction was shaken on an orbital shaker at ambient temperature for 2 hours, then 50 mg of trisamine resin (2.4
10 mmol/g) was added and shaken for 1 hour. Filtered off resin, washing with THF. The THF was blown down with a stream of nitrogen. Added 2 mL of trifluoroacetic acid and the reaction was shaken at ambient temperature overnight. The trifluoroacetic acid was blown down with a stream of nitrogen. Redissolved in 2 mL EtOAc and loaded onto a silica gel column. Eluted compound with 60%
15 EtOAc in hexanes. Combined and concentrated pure fractions, redissolved in acetonitrile/water and lyophilized to a white fluffy powder. (0.13 g, 84%) APCI (AP-): 517.2 (M-H)⁻.

Example 5

20

Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-
amide] 1-*p*-tolylamide



25 Step 1: Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

-168-

DL-proline (10 g, 86.9 mmol) was dissolved in 120 mL of THF: H₂O (2:1), added 87 mL of a 2M NaOH solution followed by Boc₂O (24.6 g, 113 mmol). Stirred at ambient temperature overnight. Removed THF in vacuo, acidified water to pH 3 with citric acid, extracted twice with EtOAc, washed
5 organics with water, brine, dried with MgSO₄, filtered and concentrated to yield title compound as a white solid. (18.7 g, quant.) APCI (AP-): 214.1 (M-H)⁻.

Step 2: 2-(2'-*tert*-Butylsulfamoyl-3-fluoro-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

10 Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (1.6 g, 7.4 mmol) was dissolved in 40 mL dry diethyl ether under Ar, cooled in an ice water bath, added dry pyridine (3 mL, 37.2 mmol) followed by the dropwise addition of oxalyl chloride (1.6 mL, 18.6 mmol). A precipitate forms immediately. The reaction was stirred vigorously at 0°C for one hour, then at ambient temperature for one
15 hour. Added 50 mL diethyl ether, filtered off solids washing with diethyl ether. Concentrated the filtrates to an off-white oil. Redissolved oil in 40 mL dry DCM under Ar, cooled to 0°C, added 3 mL pyridine followed by 4'-amino-3'-fluoro-biphenyl-2-sulfonic acid *tert*-butylamide (2 g, 6.2 mmol). The reaction was allowed to warm to ambient temperature, stirred for 3 hours then concentrated.
20 Redissolved in 250 mL EtOAc, washed with 10% citric acid (2x100 mL), water (3x100 mL), brine (100 mL), dried with MgSO₄, filtered and concentrated. Purified on a silica gel column eluted with 30% moving to 50% EtOAc in hexanes. Combined and concentrated pure fractions to yield title compound as a white solid. (1.4 g, 42%) APCI (AP-): 518.2 (M-H)⁻.

25

Step 3: Pyrrolidine-2-carboxylic acid (2'-*tert*-butylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide

2-(2'-*tert*-Butylsulfamoyl-3-fluoro-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (1.3 g, 2.5 mmol) was dissolved in 40 mL DCM,
30 cooled in an ice bath, added 10 mL trifluoroacetic acid, stirred at 0°C for 30 minutes then at ambient temperature for one hour. Concentrated solution,

-169-

redissolved in EtOAc (150 mL), washed with saturated NaHCO₃ (2x100 mL), water (100 mL), brine (100 mL), dried with MgSO₄, filtered and concentrated to yield title compound as a white solid. (1 g, 91%) APCI (AP-): 418.2 (M-H)⁻.


5 **Step 4: Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-p-tolylamide**

To a solution of pyrrolidine-2-carboxylic acid (2'-*tert*-butylsulfamoyl-3'-fluoro-biphenyl-4-yl)-amide (0.12 mmol/mL, 0.24 mmol) in 2 mL of THF was added p-tolylisocyanate. The reaction was shaken on an orbital shaker at ambient
10 temperature for 2 hours, then 50 mg of trisamine resin (2.4 mmol/g) was added and shaken for 1 hour. Filtered off resin, washing with THF. The THF was blown down with a stream of nitrogen. Added 2 mL of trifluoroacetic acid and the reaction was shaken at ambient temperature overnight. The trifluoroacetic acid was blown down with a stream of nitrogen. Loaded onto a silica gel column
15 and eluted using an automated ISCO system with gradient of 40-100% EtOAc in hexanes over 40 minutes. Combined and concentrated pure fractions, redissolved in acetonitrile/water and lyophilized to a white fluffy powder. (0.084 g, 73%) APCI (AP-): 495.1 (M-H)⁻.

20

Example 6

Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-[(4-methoxy-phenyl)-amide]




-170-

This compound was prepared using the same procedures as found for Example 5 with p-methoxyphenylisocyanate substituted for p-tolylisocyanate in step 4. (0.089 g, 72%) APCI (AP-): 511.1 (M-H)⁻.

5

Example 7

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-bromo-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]




10

This compound was prepared using the same procedures as found for Example 5 with 4-bromophenylisocyanate substituted for p-tolylisocyanate in step 4. (0.097 g, 72%) APCI (AP-): 559.0, 561.0 (M-H)⁻.

Example 8

15

Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-phenylamide




20

This compound was prepared using the same procedures as found for Example 5 with phenylisocyanate substituted for p-tolylisocyanate in step 4. (0.084 g, 73%) APCI (AP-): 481.1 (M-H)⁻.

-171-

Example 9

Pyrrolidine-1,2-dicarboxylic acid 1-[(3,4-difluoro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]




5

This compound was prepared using the same procedures as found for Example 5 with 3,4-difluorophenylisocyanate substituted for p-tolylisocyanate in step 4. (0.079 g, 63%) APCI (AP-): 517.0 (M-H)⁻.

10

Example 10

Pyrrolidine-1,2-dicarboxylic acid 1-[(3-fluoro-4-methyl-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]



15


This compound was prepared using the same procedures as found for Example 5 with 3-fluoro-4-methylphenylisocyanate substituted for p-tolylisocyanate in step 4. (0.083 g, 67%) APCI (AP-): 513.0 (M-H)⁻.

20

Example 11

-172-

Pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]



5 Step 1: Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

DL-proline (10 g, 86.9 mmol) was dissolved in 120 mL of THF: H₂O (2:1), added 87 mL of a 2M NaOH solution followed by Boc₂O (24.6 g, 113 mmol). Stirred at ambient temperature overnight. Removed THF in vacuo, acidified water to pH 3 with citric acid, extracted twice with EtOAc, washed
10 organics with water, brine, dried with MgSO₄, filtered and concentrated to yield title compound as a white solid. (18.7 g, quant.) APCI (AP-): 214.1 (M-H)⁻.

Step 2: 2-(4-Bromo-2-fluoro-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

15 Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (3.8 g, 18 mmol) was dissolved in 150 mL dry diethyl ether under Ar, cooled in an ice water bath, added dry pyridine (8.6 mL, 106 mmol) followed by the dropwise addition of oxalyl chloride (4.6 mL, 53 mmol). A precipitate formed immediately. The reaction was stirred vigorously at 0°C for one hour, then at ambient temperature for one hour.
20 100 mL diethyl ether was then added. The solids were filtered off and washed with diethyl ether. The filtrates were concentrated to an off-white oil. The solid was redissolved oil in 50 mL dry DCM under Ar, cooled to 0°C, and 6 mL pyridine was added, followed by 4-bromo-2-fluoroaniline (3.4 g, 17.7 mmol). The reaction was allowed to warm to ambient temperature, stirred for 3 hours then
25 concentrated. The concentrate was redissolved in 250 mL EtOAc, washed with 10% HCl (3x100 mL), brine (100 mL), dried with MgSO₄, filtered and

concentrated. The residue was purified on a silica gel column eluted with 20% moving to 40% EtOAc in hexanes to yield title compound as a white solid. (3.2 g, 47%) APCI (AP-): 385.1, 387.1 (M-H)⁻.

5 **Step 3: 2-(3-Fluoro-2'-methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester**

2-(4-Bromo-2-fluoro-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (2 g, 4.8 mmol), 2-(methylthio)benzene boronic acid (0.98 g, 5.8 mmol), and tetrabutylammonium bromide (78 mg, 0.24 mmol) were combined in
10 30 mL toluene, added 5 mL of a 2M aqueous Na₂CO₃ solution followed by tetrakis(triphenyl)phosphine palladium(0) (0.28 g, 0.24 mmol). The reaction was heated at reflux for 4 hours, cooled, concentrated, redissolved in EtOAc (250 mL), washed with water (3x200 mL), brine (200 mL), dried with MgSO₄, filtered, and concentrated in the presence of 4 g coarse silica. The residue was purified by
15 column chromatography on silica gel eluting with 20% EtOAc in hexanes to yield title compound as a yellow foam. (1.84 g, 88%) APCI (AP-): 429.2 (M-H)⁻.

Step 4: 2-(3-Fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

20 2-(3-Fluoro-2'-methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (0.57 g, 1.32 mmol) was dissolved in 10 mL EtOAc, and *m*-chloroperoxybenzoic acid (1.5 g, 5.3 mmol) was added in one portion to the resulting mixture. The mixture was stirred at ambient temperature for 3 hours, then diluted with 100 mL EtOAc, washed with sat. NaHCO₃ (3x100
25 mL), brine, dried with MgSO₄, filtered and concentrated. The residue was purified by column chromatography on silica gel eluting with 33% moving to 50% EtOAc in hexanes to yield title compound as a yellow foam. (0.48 g, 75%) APCI (AP-): 461.2 (M-H)⁻.

30 **Step 5: Pyrrolidine-2-carboxylic acid (3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide**

-174-

2-(3-Fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (0.45 g, 0.93 mmol) was dissolved in 15 mL DCM and 15 mL trifluoroacetic acid and the resulting mixture was stirred at ambient temperature for 1 hour. The mixture was concentrated, and the residue was
5 redissolved in 100 mL EtOAc, washed with sat. NaCl, dried with MgSO₄, filtered and concentrated to yield title compound as an oil. (0.27 g, 80%) APCI (AP-): 361.1 (M-H)⁻.


Step 6: Pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]

Pyrrolidine-2-carboxylic acid (3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide (0.27 g, 0.75 mmol) and (5-Chloro-pyridin-2-yl)-carbamic acid 4-nitrophenyl ester (0.22 g, 0.0.75 mmol; see Example 3, step 3a) were combined in 10 mL DMF and heated at 80°C for 4 hours, then cooled. 200 mL of water were then
15 added and a precipitate formed. The solids were filtered off, washed with water, redissolved in EtOAc, dried with MgSO₄, filtered and concentrated. The residue was purified by column chromatography on silica gel eluting with 70% EtOAc in hexanes to yield title compound. (220 mg, 57%) APCI (AP-): 515.1, 517.1 (M-H)⁻.

20

Example 12

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]



25

-175-

Step 1: Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

DL-proline (10 g, 86.9 mmol) was dissolved in 120 mL of THF: H₂O (2:1), and 87 mL of a 2M NaOH solution was added to the resulting mixture, followed by Boc₂O (24.6 g, 113 mmol). The mixture was stirred at ambient
5 temperature overnight. THF was removed in vacuo, and the remaining water was acidified to pH 3 with citric acid, then extracted twice with EtOAc. The combined organic layers were washed with water, brine, dried with MgSO₄, filtered and concentrated to yield title compound as a white solid. (18.7 g, quant.) APCI (AP-): 214.1 (M-H)⁻.

10

Step 2: 2-(4-Bromo-2-fluoro-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (3.8 g, 18 mmol) was dissolved in 150 mL dry diethyl ether under Ar, and the resulting solution was
15 cooled in an ice water bath, while dry pyridine (8.6 mL, 106 mmol) was added, followed by the dropwise addition of oxalyl chloride (4.6 mL, 53 mmol). A precipitate formed immediately. The reaction was stirred vigorously at 0°C for one hour, then at ambient temperature for one hour. 100 mL of diethyl ether were then added, and the solids were removed by filtration washing with diethyl
20 ether. Concentrated the filtrates to an off-white oil. Redissolved oil in 50 mL dry DCM under Ar, cooled to 0°C, added 6 mL pyridine followed by 4-bromo-2-fluoroaniline (3.4 g, 17.7 mmol). The reaction was allowed to warm to ambient temperature, stirred for 3 hours then concentrated. Redissolved in 250 mL EtOAc, washed with 10% HCl (3x100 mL), brine (100 mL), dried with MgSO₄,
25 filtered and concentrated. Purified on a silica gel column eluted with 20% moving to 40% EtOAc in hexanes. Combined and concentrated pure fractions to yield title compound as a white solid. (3.2 g, 47%) APCI (AP-): 385.1, 387.1 (M-H)⁻.

30

Step 3: 2-(3-Fluoro-2'-methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

-176-

2-(4-Bromo-2-fluoro-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (2 g, 4.8 mmol), 2-(methylthio)benzene boronic acid (0.98 g, 5.8 mmol), and tetrabutylammonium bromide (78 mg, 0.24 mmol) were combined in 30 mL toluene, added 5 mL of a 2M aqueous Na₂CO₃ solution followed by
5 tetrakis(triphenyl)phosphine palladium(0) (0.28 g, 0.24 mmol). Heated reaction at reflux for 4 hours, cooled, concentrated, redissolved in EtOAc (250 mL), washed with water (3x200 mL), brine (200 mL), dried with MgSO₄, filtered, and concentrated in the presence of 4 g coarse silica. Loaded silica onto a silica gel column, eluted with 20% EtOAc in hexanes. Combined and concentrated pure
10 fractions to yield title compound as a yellow foam. (1.84 g, 88%) APCI (AP-): 429.2 (M-H)⁻.

Step 4: 2-(3-Fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

15 2-(3-Fluoro-2'-methanesulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (1.2 g, 2.8 mmol) was dissolved in 30 mL EtOAc, added *m*-chloroperoxybenzoic acid (3.2 g, 11 mmol) in one portion and stirred at ambient temperature for 3 hours. A 10% aqueous solution (50 mL) of Na₂S₂O₃ was added to the vigorously stirring reaction in order to quench peroxide. After
20 30 minutes the solution was diluted with 100 mL EtOAc, separated layers, washed organics with sat. NaHCO₃ (3x100 mL), brine, dried with MgSO₄, filtered and concentrated. Purified on a silica gel column eluted with 50% EtOAc in hexanes. Combined and concentrated pure fractions to yield title compound as a yellow foam. (1.01 g, 78%) APCI (AP-): 461.1 (M-H)⁻.

25

Step 5: Pyrrolidine-2-carboxylic acid (3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide

2-(3-Fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (1 g, 2 mmol) was dissolved in 20 mL DCM and
30 10 mL trifluoroacetic acid and stirred at ambient temperature for 1 hour.

-177-

Concentrated solution, redissolved in 100 mL CHCl_3 , repeated CHCl_3 co-concentration 3 times to yield title compound as an oil. (quant. yield) APCI (AP-): 361.1 (M-H)⁻.


5 **Step 6: Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]**

Pyrrolidine-2-carboxylic acid (3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide (0.23 g, 0.64 mmol) was dissolved in 10 mL THF, added diisopropylethylamine (0.33 mL, 1.9 mmol), followed by 4-chlorophenylisocyanate (97 mg, 0.64 mmol). Stirred reaction at ambient temperature for 1 hour then concentrated. Purified on a silica gel column eluted with 70% EtOAc in hexanes. Pure fractions were combined and concentrated, redissolved in acetonitrile/water and lyophilized to yield title compound as a white powder. (0.23 g, 70%) APCI (AP-): 514.1, 516.1 (M-H)⁻.

15

Example 13

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[5-(2-sulfamoyl-phenyl)-pyridin-2-yl]-amide]



20

Step 1: Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

DL-proline (10 g, 86.9 mmol) was dissolved in 120 mL of THF: H₂O (2:1), added 87 mL of a 2M NaOH solution followed by Boc₂O (24.6 g, 113 mmol). Stirred at ambient temperature overnight. Removed THF in vacuo, acidified water to pH 3 with citric acid, extracted twice with EtOAc, washed

25

-178-

organics with water, brine, dried with MgSO₄, filtered and concentrated to yield title compound as a white solid. (18.7 g, quant.) APCI (AP-): 214.1 (M-H)⁻.

Step 2: 2-(5-Bromo-pyridin-2-ylcarbamoyl)-pyrrolidine-1-carboxylic acid

5 *tert*-butyl ester

Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (1.97 g, 9.2 mmol) was dissolved in 150 mL dry diethyl ether under Ar, cooled in an ice water bath, added dry pyridine (3.7 mL, 45.8 mmol) followed by the dropwise addition of oxalyl chloride (2 mL, 22.9 mmol). A precipitate forms immediately. The reaction was stirred vigorously at 0°C for 2 hours, then at ambient temperature for one hour. Added 100 mL diethyl ether, filtered off solids washing with diethyl ether. Concentrated the filtrates to an off-white oil. Redissolved oil in 100 mL dry DCM under Ar, cooled to 0°C, added 3 mL pyridine followed by 2-amino-5-bromopyridine (1.32 g, 7.6 mmol). The reaction was allowed to warm to ambient temperature, stirred for 3 hours then concentrated. Redissolved in 250 mL EtOAc, washed with 10% HCl (3x100 mL), brine (100 mL), dried with MgSO₄, filtered and concentrated to yield title compound as a yellow sticky solid. (2.74 g, 97%) APCI (AP-): 368.0, 370.0 (M-H)⁻.

20 Step 3: 2-[5-(2-*tert*-Butylsulfamoyl-phenyl)-pyridin-2-ylcarbamoyl]-pyrrolidine-1-carboxylic acid *tert*-butyl ester

2-(5-Bromo-pyridin-2-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (0.92 g, 2.48 mmol), 2-(*N-tert*-butyl)phenylsulfonamide boronic acid (0.64 g, 2.48 mmol), and tetrabutylammonium bromide (40 mg, 0.12 mmol) were combined in 10 mL toluene, added 2 mL of an aqueous 2M sodium carbonate solution followed by the addition of tetrakis(triphenylphosphine) palladium(0) (144 mg, 0.12 mmol). Refluxed solution for 4 hours, cooled to room temperature, partitioned between EtOAc (150 mL) and water (100 mL), washed organics with water (3x100 mL), brine, dried with MgSO₄, filtered and concentrated. Purified on a silica gel column eluted with 40% moving to 50% EtOAc in hexanes.

-179-

Combined and concentrated pure fractions to yield title compound as a white solid. (0.92 g, 74%) APCI (AP-): 501.1 (M-H)⁻.

Step 4: Pyrrolidine-2-carboxylic acid [5-(2-*tert*-butylsulfamoyl-phenyl)-pyridin-2-yl]-amide

2-[5-(2-*tert*-Butylsulfamoyl-phenyl)-pyridin-2-ylcarbamoyl]-pyrrolidine-1-carboxylic acid *tert*-butyl ester (0.92 g, 1.83 mmol) was dissolved in 30 mL DCM, added 10 mL trifluoroacetic acid and stirred at ambient temperature for one hour. Concentrated solution, redissolved in CHCl₃ and re-concentrated. Repeated co-concentration twice to yield title compound as an oil. (quant. yield) APCI (AP-): 401.1 (M-H)⁻.

Step 5: Pyrrolidine-1,2-dicarboxylic acid 2-{[5-(2-*tert*-butylsulfamoyl-phenyl)-pyridin-2-yl]-amide} 1-[(4-chloro-phenyl)-amide]

Pyrrolidine-2-carboxylic acid [5-(2-*tert*-butylsulfamoyl-phenyl)-pyridin-2-yl]-amide (0.44 g, 0.59 mmol) was dissolved in 10 mL THF, added diisopropylethylamine (0.41 mL, 2.4 mmol) followed by 4-chlorophenylisocyanate (91 mg, 0.59 mmol). Stirred at ambient temperature for 2 hours. Diluted reaction with 50 mL EtOAc, washed with 10% citric acid (3x50 mL), water (2x50 mL), brine, dried with MgSO₄, filtered and concentrated to yield title compound. (0.28 g, 85%) APCI (AP+): 556.2, 558.2 (M-H)⁺.

Step 6: Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[5-(2-sulfamoyl-phenyl)-pyridin-2-yl]-amide}

Pyrrolidine-1,2-dicarboxylic acid 2-{[5-(2-*tert*-butylsulfamoyl-phenyl)-pyridin-2-yl]-amide} 1-[(4-chloro-phenyl)-amide] (0.28 g, 0.5 mmol) was stirred in 10 mL trifluoroacetic acid at ambient temperature for 20 hours. The solution was concentrated and purified on a silica gel column using an automated system with gradient of 40-100% EtOAc in hexanes over 40 minutes. Combined and concentrated pure fractions, redissolved in acetonitrile/water and lyophilized to


-180-

yield title compound as a light yellow powder. (0.156 g, 62%) APCI (AP+): 500.0, 501.9 (M+H)⁺.

Example 14

5

Pyrrolidine-1,2-dicarboxylic acid 2-[(3-chloro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(5-chloro-pyridin-2-yl)-amide]



10 Step 1: Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

DL-proline (10 g, 86.9 mmol) was dissolved in 120 mL of THF: H₂O (2:1), added 87 mL of a 2M NaOH solution followed by Boc₂O (24.6 g, 113 mmol). Stirred at ambient temperature overnight. Removed THF in vacuo, acidified water to pH 3 with citric acid, extracted twice with EtOAc, washed
15 organics with water, brine, dried with MgSO₄, filtered and concentrated to yield title compound as a white solid. (18.7 g, quant.) APCI (AP-): 214.1 (M-H)⁻.

Step 2: 2-(4-Bromo-2-chloro-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

20 Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (2.2 g, 10 mmol) was dissolved in 150 mL dry diethyl ether under Ar, cooled in an ice water bath, added dry pyridine (2.8 mL, 34 mmol) followed by the dropwise addition of oxalyl chloride (1.5 mL, 17 mmol). A precipitate forms immediately. The reaction was stirred vigorously at 0°C for 1 hours, then at ambient temperature for two hours.
25 Added 100 mL diethyl ether, filtered off solids washing with diethyl ether. Concentrated the filtrates to an off-white oil. Redissolved oil in 100 mL dry DCM

-181-

under Ar, cooled to 0°C, added 3 mL pyridine followed by 2-chloro-4-bromoaniline (1.8 g, 8.5 mmol). The reaction was allowed to warm to ambient temperature and stirred over the weekend. Concentrated reaction, redissolved in 250 mL EtOAc, washed with 10% citric acid (3x100 mL), water, (2x100 mL),
5 brine (100 mL), dried with MgSO₄, filtered and concentrated. Purified on a silica gel column eluted with an automated system with gradient of 0-60% EtOAc in hexanes. Combined and concentrated pure fractions to yield title compound as a yellow solid. (2.7 g, 79%) APCI (AP-): 401.0, 403.0, 405.0 (M-H)⁻.

10 **Step 3: 2-(3-Chloro-2'-methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester**

2-(4-Bromo-2-chloro-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (1 g, 2.5 mmol), 2-(methylthio)benzene boronic acid (0.5 g, 3 mmol), and tetrabutylammonium bromide (40 mg, 0.12 mmol) were combined in
15 10 mL toluene, added 2 mL of a 2M aqueous Na₂CO₃ solution followed by tetrakis(triphenylphosphine) palladium(0) (0.14 g, 0.12 mmol). Heated reaction at reflux for 5 hours, cooled, dissolved in EtOAc (150 mL), washed with water (3x50 mL), brine (50 mL), dried with MgSO₄, filtered, and concentrated. Purified
20 on a silica gel column eluted using an automated system with gradient of 0-50% EtOAc in hexanes over 40 minutes. Combined and concentrated pure fractions to yield title compound as a white foam. (1.01 g, 91%) APCI (AP-): 445.1, 447.1 (M-H)⁻.

25 **Step 4: 2-(3-Chloro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester**

2-(3-Chloro-2'-methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (1 g, 2 mmol) was dissolved in 20 mL EtOAc, added *m*-chloroperoxybenzoic acid (2.6 g, 8.9 mmol) in one portion and stirred at ambient temperature for 3 hours. A 10% aqueous solution (30 mL) of Na₂S₂O₃
30 was added to the vigorously stirring reaction in order to quench peroxide. After 20 minutes the solution was diluted with 100 mL EtOAc, separated layers, washed

-182-

organics with sat. NaHCO_3 (3x100 mL), brine, dried with MgSO_4 , filtered and concentrated to yield title compound as a clear oil. (1 g, 100%) APCI (AP-): 477.1, 479.1 (M-H)⁻.

5 **Step 5: Pyrrolidine-2-carboxylic acid (3-chloro-2'-methanesulfonyl-biphenyl-4-yl)-amide**

2-(3-Chloro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (1 g, 2 mmol) was dissolved in 20 mL DCM and 20 mL trifluoroacetic acid and stirred at ambient temperature for 2 hours.

- 10 Concentrated solution, redissolved in 100 mL CHCl_3 , repeated CHCl_3 co-concentration 4 times to yield title compound as a yellow foam. (quant. yield) APCI (AP+): 379.1, 381.1 (M+H)⁺.

15 **Step 6: Pyrrolidine-1,2-dicarboxylic acid 2-[(3-chloro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(5-chloro-pyridin-2-yl)-amide]**


- Pyrrolidine-2-carboxylic acid (3-chloro-2'-methanesulfonyl-biphenyl-4-yl)-amide (0.11 g, 0.29 mmol) and (5-Chloro-pyridin-2-yl)-carbamic acid 4-nitrophenyl ester (0.085 g, 0.29 mmol; see Example 3, step 3a) were combined in 3 mL DMF and heated at 50°C for 2 hours, dissolved in 100 mL EtOAc, washed with
- 20 sat. NaHCO_3 (3x50 mL), brine, dried with MgSO_4 , filtered and concentrated. Purified on a silica gel column eluting using an automated system with gradient of 0-100% EtOAc in hexanes over 40 minutes. Combined and concentrated pure fractions, redissolved in acetonitrile/water and lyophilized to yield title compound as a white powder. (110 mg, 71%) APCI (AP+): 533.0, 535.0 (M+H)⁺.

25

Example 15

Pyrrolidine-1,2-dicarboxylic acid 2-[(3-chloro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide]

-183-



Step 1: Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

DL-proline (10 g, 86.9 mmol) was dissolved in 120 mL of THF: H₂O (2:1), added 87 mL of a 2M NaOH solution followed by Boc₂O (24.6 g, 113 mmol). Stirred at ambient temperature overnight. Removed THF in vacuo, acidified water to pH 3 with citric acid, extracted twice with EtOAc, washed organics with water, brine, dried with MgSO₄, filtered and concentrated to yield title compound as a white solid. (18.7 g, quant.) APCI (AP-): 214.1 (M-H)⁻.

10

Step 2: 2-(4-Bromo-2-chloro-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (2.2 g, 10 mmol) was dissolved in 150 mL dry diethyl ether under Ar, cooled in an ice water bath, added dry pyridine (2.8 mL, 34 mmol) followed by the dropwise addition of oxalyl chloride (1.5 mL, 17 mmol). A precipitate forms immediately. The reaction was stirred vigorously at 0°C for 1 hours, then at ambient temperature for two hours. Added 100 mL diethyl ether, filtered off solids washing with diethyl ether. Concentrated the filtrates to an off-white oil. Redissolved oil in 100 mL dry DCM under Ar, cooled to 0°C, added 3 mL pyridine followed by 2-chloro-4-bromoaniline (1.8 g, 8.5 mmol). The reaction was allowed to warm to ambient temperature and stirred over the weekend. Concentrated reaction, redissolved in 250 mL EtOAc, washed with 10% citric acid (3x100 mL), water, (2x100 mL), brine (100 mL), dried with MgSO₄, filtered and concentrated. Purified on a silica gel column eluted with an automated system with gradient of 0-60% EtOAc in

25

-184-

hexanes. Combined and concentrated pure fractions to yield title compound as a yellow solid. (2.7 g, 79%) APCI (AP-): 401.0, 403.0, 405.0 (M-H)⁻.

Step 3: 2-(3-Chloro-2'-methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

2-(4-Bromo-2-chloro-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (1 g, 2.5 mmol), 2-(methylthio)benzene boronic acid (0.5 g, 3 mmol), and tetrabutylammonium bromide (40 mg, 0.12 mmol) were combined in 10 mL toluene, added 2 mL of a 2M aqueous Na₂CO₃ solution followed by tetrakis(triphenylphosphine) palladium(0) (0.14 g, 0.12 mmol). Heated reaction at reflux for 5 hours, cooled, dissolved in EtOAc (150 mL), washed with water (3x50 mL), brine (50 mL), dried with MgSO₄, filtered, and concentrated. Purified on a silica gel column eluted using an automated system with gradient of 0-50% EtOAc in hexanes over 40 minutes. Combined and concentrated pure fractions to yield title compound as a white foam. (1.01 g, 91%) APCI (AP-): 445.1, 447.1 (M-H)⁻.

Step 4: 2-(3-Chloro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

2-(3-Chloro-2'-methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (1 g, 2 mmol) was dissolved in 20 mL EtOAc, added *m*-chloroperoxybenzoic acid (2.6 g, 8.9 mmol) in one portion and stirred at ambient temperature for 3 hours. A 10% aqueous solution (30 mL) of Na₂S₂O₃ was added to the vigorously stirring reaction in order to quench peroxide. After 20 minutes the solution was diluted with 100 mL EtOAc, separated layers, washed organics with sat. NaHCO₃ (3x100 mL), brine, dried with MgSO₄, filtered and concentrated to yield title compound as a clear oil. (1 g, 100%) APCI (AP-): 477.1, 479.1 (M-H)⁻.

Step 5: Pyrrolidine-2-carboxylic acid (3-chloro-2'-methanesulfonyl-biphenyl-4-yl)-amide

-185-

2-(3-Chloro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (1 g, 2 mmol) was dissolved in 20 mL DCM and 20 mL trifluoroacetic acid and stirred at ambient temperature for 2 hours. Concentrated solution, redissolved in 100 mL CHCl_3 , repeated CHCl_3 co-concentration 4 times to yield title compound as a yellow foam. (quant. yield) APCI (AP+): 379.1, 381.1 (M+H)⁺.


Step 6: Pyrrolidine-1,2-dicarboxylic acid 2-[(3-chloro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide]

10 Pyrrolidine-2-carboxylic acid (3-chloro-2'-methanesulfonyl-biphenyl-4-yl)-amide (0.16 g, 0.42 mmol) was dissolved in 5 mL THF, added diisopropylethylamine (0.22 mL, 1.3 mmol) followed by 4-chlorophenylisocyanate (65 mg, 0.42 mmol). Stirred at ambient temperature for 20 hours. The reaction was concentrated and purified using an automated system with gradient of 0-100% EtOAc in hexanes over 40 minutes. Combined and concentrated pure fractions, redissolved in acetonitrile/water and lyophilized to yield title compound as a white powder. (0.175 g, 78%) APCI (AP+): 532.0, 534.0(M-H)⁺.

20

Example 16

Pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[5-(2-sulfamoyl-phenyl)-pyridin-2-yl]-amide]



25

Step 1: Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

-186-

DL-proline (10 g, 86.9 mmol) was dissolved in 120 mL of THF: H₂O (2:1), added 87 mL of a 2M NaOH solution followed by Boc₂O (24.6 g, 113 mmol). Stirred at ambient temperature overnight. Removed THF in vacuo, acidified water to pH 3 with citric acid, extracted twice with EtOAc, washed
5 organics with water, brine, dried with MgSO₄, filtered and concentrated to yield title compound as a white solid. (18.7 g, quant.) APCI (AP-): 214.1 (M-H)⁻.

Step 2: 2-(5-Bromo-pyridin-2-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

10 Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (1.97 g, 9.2 mmol) was dissolved in 150 mL dry diethyl ether under Ar, cooled in an ice water bath, added dry pyridine (3.7 mL, 45.8 mmol) followed by the dropwise addition of oxalyl chloride (2 mL, 22.9 mmol). A precipitate forms immediately. The reaction was stirred vigorously at 0°C for 2 hours, then at ambient temperature for one hour.
15 Added 100 mL diethyl ether, filtered off solids washing with diethyl ether. Concentrated the filtrates to an off-white oil. Redissolved oil in 100 mL dry DCM under Ar, cooled to 0°C, added 3 mL pyridine followed by 2-amino-5-bromopyridine (1.32 g, 7.6 mmol). The reaction was allowed to warm to ambient temperature, stirred for 3 hours then concentrated. Redissolved in 250 mL
20 EtOAc, washed with 10% HCl (3x100 mL), brine (100 mL), dried with MgSO₄, filtered and concentrated to yield title compound as a yellow sticky solid. (2.74 g, 97%) APCI (AP-): 368.0, 370.0 (M-H)⁻.

Step 3: 2-[5-(2-*tert*-Butylsulfamoyl-phenyl)-pyridin-2-ylcarbamoyl]-pyrrolidine-1-carboxylic acid *tert*-butyl ester

25 2-(5-Bromo-pyridin-2-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (0.92 g, 2.48 mmol), 2-(*N*-*tert*-butyl)phenylsulfonamide boronic acid (0.64 g, 2.48 mmol), and tetrabutylammonium bromide (40 mg, 0.12 mmol) were combined in 10 mL toluene, added 2 mL of an aqueous 2M sodium carbonate
30 solution followed by the addition of tetrakis(triphenylphosphine) palladium(0) (144 mg, 0.12 mmol). Refluxed solution for 4 hours, cooled to room temperature,

-187-

partitioned between EtOAc (150 mL) and water (100 mL), washed organics with water (3x100 mL), brine, dried with MgSO₄, filtered and concentrated. Purified on a silica gel column eluted with 40% moving to 50% EtOAc in hexanes. Combined and concentrated pure fractions to yield title compound as a white solid. (0.92 g, 74%) APCI (AP-): 501.1 (M-H)⁻.

Step 4: Pyrrolidine-2-carboxylic acid [5-(2-*tert*-butylsulfamoyl-phenyl)-pyridin-2-yl]-amide

2-[5-(2-*tert*-Butylsulfamoyl-phenyl)-pyridin-2-ylcarbamoyl]-pyrrolidine-1-carboxylic acid *tert*-butyl ester (0.92 g, 1.83 mmol) was dissolved in 30 mL DCM, added 10 mL trifluoroacetic acid and stirred at ambient temperature for one hour. Concentrated solution, redissolved in CHCl₃ and re-concentrated. Repeated co-concentration twice to yield title compound as an oil. (quant. yield) APCI (AP-): 401.1 (M-H)⁻.


Step 5: Pyrrolidine-1,2-dicarboxylic acid 2-[[5-(2-*tert*-butylsulfamoyl-phenyl)-pyridin-2-yl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide]

Pyrrolidine-2-carboxylic acid [5-(2-*tert*-butylsulfamoyl-phenyl)-pyridin-2-yl]-amide (0.17 g, 0.25 mmol) and (5-Chloro-pyridin-2-yl)-carbamic acid 4-nitro-phenyl ester (0.074 g, 0.25 mmol; see Example 3, step 3a) were combined in 3 mL DMF, added diisopropylethylamine (0.18 mL, 1 mmol) and heated at 50°C for 2 hours. Dissolved in 100 mL EtOAc, washed with sat. NaHCO₃ (3x50 mL), brine, dried with MgSO₄, filtered and concentrated. Purified on a silica gel column eluting using an automated system with gradient of 30-100% EtOAc in hexanes over 40 minutes. Combined and concentrated pure fractions to yield title compound as a white solid. (90 mg, 64%) APCI (AP-): 555.2, 557.2 (M-H)⁻.

Example 17

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]

-188-



Step 1: (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

- 5 Cis-4-hydroxy-D-proline (1.1 g, 8.4 mmol) was dissolved in 18 mL of THF/ H₂O (2:1), added 2M aqueous NaOH solution (6.3 mL, 13 mmol), followed by Boc₂O. Reaction was stirred at ambient temperature over the weekend. Solution was taken up in 150 mL EtOAc, washed with 10% HCl (3x100 mL), dried organics with MgSO₄, filtered and concentrated to yield title compound as a
- 10 white foam. (1.5 g, 77%) APCI (AP-): 230.1 (M-H)⁻.

Step 2: (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

- (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester
- 15 (1.5 g, 6.5 mmol) was dissolved in 20 mL DMF, added imidazole (1.1 g, 16 mmol), dimethylaminopyridine (80 mg, 0.65 mmol), and *tert*-butyldimethylsilyl chloride (1.08 g, 7.1 mmol). Stirred at ambient temperature overnight, dissolved in 150 mL EtOAc, washed with 10% citric acid (3x100 mL), water (2x100 mL), brine, dried with MgSO₄, filtered, and concentrated. Redissolved in 50 mL
- 20 CHCl₃ and reconcentrated to yield title compound as a clear oil. (1.54 g, 69%) APCI (AP-): 344.2 (M-H)⁻.

Step 3: (2R,4R)-2-(4-Bromo-2-fluoro-phenylcarbamoyl)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

-189-

(2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (1.54 g, 4.5 mmol) was dissolved in 50 mL diethyl ether under an Ar atmosphere, cooled in an ice water bath, added dry pyridine (1.4 mL, 17.8 mmol), followed by the dropwise addition of oxalyl chloride (0.58 mL, 6.7 mmol). Reaction was stirred at 0°C for 1 hour, then at ambient temperature for 1 hour. Diluted reaction with 50 mL diethyl ether, filtered off solids and concentrated filtrates. Redissolved filtrate in 30 mL dry DCM under an Ar atmosphere, cooled in an ice water bath, added 1 mL dry pyridine followed by 4-bromo-2-fluoroaniline (0.85 g, 4.5 mmol). Stirred reaction at ambient temperature for 20 hours. Concentrated solution, redissolved in EtOAc (150 mL), washed with 10% citric acid (3x100 mL), water (2x100 mL), brine, dried with MgSO₄, filtered and concentrated. Purified on a silica gel column eluting with an automated system with gradient of 0-60% EtOAc in hexanes over 40 minutes. Combined and concentrated pure fractions to yield title compound as a clear oil. (0.96 g, 42%) APCI (AP-): 515.1, 517.1 (M-H)⁺.

Step 4: (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

(2R,4R)-2-(4-Bromo-2-fluoro-phenylcarbamoyl)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (0.95 g, 1.84 mmol), 2-(methylthio)benzene boronic acid (0.37 g, 2.2 mmol), and tetrabutylammonium bromide (30 mg, 0.09 mmol) were combined in 15 mL toluene, added 2 mL of a 2M aqueous Na₂CO₃ solution followed by tetrakis(triphenyl)phosphine palladium(0) (0.11 g, 0.09 mmol). Heated reaction at reflux for 5 hours, cooled, dissolved in EtOAc (100 mL), washed with water (3x50 mL), brine (50 mL), dried with MgSO₄, filtered, and concentrated. Purified on a silica gel column eluted using an automated system with gradient of 0-60% EtOAc in hexanes over 40 minutes. Combined and concentrated pure fractions to yield title compound as a light yellow foam. (0.77 g, 74%) APCI (AP-): 559.3 (M-H)⁺.

-190-

Step 5: (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

(2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methylsulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (0.84 g, 1.35 mmol) was dissolved in 20 mL EtOAc, added added *m*-chloroperoxybenzoic acid (1.6 g, 5.4 mmol) in one portion and stirred at ambient temperature for 3 hours. A 10% aqueous solution (50 mL) of Na₂S₂O₃ was added to the vigorously stirring reaction in order to quench peroxide. After 20 minutes the solution was diluted with 100 mL EtOAc, separated layers, washed organics with sat. NaHCO₃ (3x100 mL), water (2x100 mL), brine, dried with MgSO₄, filtered and concentrated to yield title compound as a light brown foam. (0.8 g, 100%) APCI (AP-): 591.3 (M-H)⁻.

Step 6: (2R,4R)-4-Hydroxy-pyrrolidine-2-carboxylic acid (3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide

(2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (0.8 g, 1.35 mmol) was dissolved in 30 mL DCM, added 10 mL trifluoroacetic acid and stirred reaction at ambient temperature for 90 minutes. Solution was concentrated to yield title compound as an amber oil. (100% crude yield) APCI (AP-): (M-H)⁻.

Step 7: (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chlorophenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]

(2R,4R)-4-Hydroxy-pyrrolidine-2-carboxylic acid (3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide (0.51 g, 1.35 mmol) was dissolved in 20 mL THF, added diisopropylethylamine (0.94 mL, 5.4 mmol) followed by 4-chlorophenylisocyanate (0.21 g, 1.35 mmol). Stirred at ambient temperature for 2 hours. The reaction was dissolved in 100 mL EtOAc, washed with 10% citric acid (3x50 mL), water (2x50 mL), brine, dried with MgSO₄, filtered and concentrated.


-191-

Purified using an automated system with gradient of 50-100% EtOAc in hexanes over 40 minutes. Compound was still eluting at end of run. Switched gradient to 0-15% MeOH in EtOAc over 40 minutes. Combined and concentrated pure fractions to yield title compound as a white solid. (0.54 g, 75%) APCI (AP+):

5 530.1, 532.1(M-H)⁺.

Example 18

10 **Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-amide]**



Step 1: Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

DL-proline (10 g, 86.9 mmol) was dissolved in 120 mL of THF: H₂O (2:1), added 87 mL of a 2M NaOH solution followed by Boc₂O (24.6 g, 113 mmol). Stirred at ambient temperature overnight. Removed THF in vacuo, acidified water to pH 3 with citric acid, extracted twice with EtOAc, washed organics with water, brine, dried with MgSO₄, filtered and concentrated to yield title compound as a white solid. (18.7 g, quant.) APCI (AP-): 214.1 (M-H)⁻.

20

Step 2: 2-(4-Cyano-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (2.1 g, 9.8 mmol) was dissolved in 100 mL dry diethyl ether under an argon atmosphere, added dry pyridine (3.3 mL, 32.5 mmol), cooled to 0°C then added oxalyl chloride (1.4 mL, 16.3 mmol) dropwise. A precipitate forms immediately upon addition of oxalyl

25

-192-

chloride and therefore reaction requires vigorous stirring. Stirred at 0°C for one hour than at ambient temperature for 2 hours. Dry diethyl ether (100 mL) was added and the solids were filtered off, washing with more diethyl ether. The filtrates were concentrated, then redissolved in dry DCM under argon, cooled to 5 0°C, added 1 mL dry pyridine then added 4-amino-benzonitrile (0.96 g, 8.1 mmol) in one portion. Stirred at ambient temperature for 2 hours, concentrated, redissolved in ethyl acetate (250 mL), washed with 10% citric acid (3x100 mL), water, brine, dried with MgSO₄, filtered and concentrated to yield title compound as a white foam. (2.4 g, 93%) APCI (AP-): 314.1 (M-H)⁻.

10

Step 3: Pyrrolidine-2-carboxylic acid (4-cyano-phenyl)-amide

2-(4-Cyano-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (3 g, 8 mmol) was dissolved in 90 mL of a 1:2 mixture of trifluoroacetic acid and dichloromethane. The reaction was stirred at ambient temperature for one 15 hour then concentrated to yield title compound as an amber oil which was carried onto next step as is. APCI (AP-): 214.1 (M-H)⁻.

Step 4: Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-cyano-phenyl)-amide]

20 Pyrrolidine-2-carboxylic acid (4-cyano-phenyl)-amide (0.9 g, 4 mmol) was dissolved in 25 mL tetrahydrofuran, added diisopropylethylamine (3.5 mL, 20 mmol), followed by the addition of 4-chlorophenylisocyanate (0.61 g, 4 mmol). The reaction was stirred at ambient temperature for one hour then concentrated solution. Redissolved in 100 mL ethyl acetate, washed with 10% citric acid 25 (2x100 mL), water, brine, dried with MgSO₄, filtered and concentrated. Recrystallized from ethyl acetate and hexanes to yield title compound as a white powder. (1.1 g, 75%) APCI (AP-): 367.1, 369.1 (M-H)⁻.

Step 5: Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl)-amide]


30

-193-

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-cyano-phenyl)-amide] (0.2 g, 0.54 mmol) was dissolved in 2 mL dry methanol under argon, cooled in an ice bath, bubbled in HCl(g) for 10 minutes, removed argon line and HCl line and sealed off flask. Stirred at 0°C for 20 minutes, then at
5 ambient temperature for 4 hours. The solution was concentrated in vacuo to a yellow foam. The foam was redissolved in 4 mL dry methanol under argon, added N-methylethylenediamine (0.048 mL, 0.54 mmol) and stirred at ambient temperature overnight. The solution was concentrated in vacuo then purified by preparatory HPLC. Combined and lyophilized pure fractions to yield the TFA salt
10 of the title compound as a white fluffy powder. (0.2 g, 69%) APCI (AP-): 424.1, 426.1 (M-H)⁻.

Example 19

15 **Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-biphenyl-4-yl)-amide]**



Step 1: Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

20 DL-proline (10 g, 86.9 mmol) was dissolved in 120 mL of THF: H₂O (2:1), added 87 mL of a 2M NaOH solution followed by Boc₂O (24.6 g, 113 mmol). Stirred at ambient temperature overnight. Removed THF in vacuo, acidified water to pH 3 with citric acid, extracted twice with EtOAc, washed organics with water, brine, dried with MgSO₄, filtered and concentrated to yield
25 title compound as a white solid. (18.7 g, quant.) APCI (AP-): 214.1 (M-H)⁻.

-194-

Step 2: 2-(4-Bromo-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (5 g, 23 mmol) was dissolved in 250 mL dry diethyl ether under Ar, cooled in an ice water bath, added
5 dry pyridine (11.3 mL, 139 mmol) followed by the dropwise addition of oxalyl chloride (6.1 mL, 69 mmol). A precipitate forms immediately. The reaction was stirred vigorously at 0°C for one hour, then at ambient temperature for one hour. Added 100 mL diethyl ether, filtered off solids washing with diethyl ether. Concentrated the filtrates to an off white oil. Redissolved (3.66 g, 15 mmol) oil in
10 40 mL dry DCM under Ar, cooled to 0°C, added 4.6 mL pyridine followed by 4-bromoaniline (2.7 g, 15 mmol). The reaction was allowed to warm to ambient temperature, stirred for 3 hours then concentrated. Redissolved in 100 mL EtOAc, washed with 1 N HCl (2x100 mL), brine (100 mL), dried with MgSO₄, filtered and concentrated to yield title compound. (4.56 g, 80%) APCI (AP-):
15 369.0, 370.0 (M-H)⁻.

Step 3: 2-(2'-Methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

2-(4-Bromo-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl
20 ester (2.57 g, 6.9 mmol), 2-(methylthio)benzene boronic acid (1.4 g, 8.3 mmol), and tetrabutylammonium bromide (110 mg, 0.33 mmol) were combined in 30 mL toluene, added 6.9 mL of a 2M aqueous Na₂CO₃ solution followed by tetrakis(triphenylphosphine) palladium(0) (0.40 g, 0.34 mmol). Heated reaction at reflux overnight, cooled, concentrated, redissolved in EtOAc (250 mL), washed
25 with water (3x200 mL), brine (200 mL), dried with MgSO₄, filtered, and concentrated. Purified on a silica gel column eluted with 20% moving to 40% EtOAc in hexanes. Combined and concentrated pure fractions to yield title compound. (1.75 g, 60%) APCI (AP-): 411.1 (M-H)⁻.

30 **Step 4: 2-(2'-Methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester**

-195-

2-(2'-Methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (1.0 g, 2.4 mmol) was dissolved in 10 mL acetonitrile, added oxone (2.98 g, 4.8 mmol) in one portion and stirred at ambient temperature for 1 week and then concentrated. Redissolved in 100 mL EtOAc and 100 mL sat.

- 5 NaHCO₃, separated layers, washed organics with brine, dried with MgSO₄, filtered and concentrated. Purified on a silica gel column eluted with 20% moving to 40% EtOAc in hexanes. Combined and concentrated pure fractions to yield title compound. (0.80 g, 75%) APCI (AP-): 443.1 (M-H)⁻.

10 **Step 5: Pyrrolidine-2-carboxylic acid (2'-methanesulfonyl-biphenyl-4-yl)-amide**

2-(2'-Methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (0.8 g, 1.7 mmol) was dissolved in 25 mL DCM and 5 mL trifluoroacetic acid and stirred at ambient temperature for one hour and

- 15 concentrated. Redissolved in 100 mL EtOAc and 100 mL sat. NaHCO₃, separated layers, washed organics with brine, dried with MgSO₄, filtered and concentrated to yield title compound as an oil (quant. yield) APCI (AP-): 343.1 (M-H)⁻.

20 **Step 6: Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-biphenyl-4-yl)-amide]**


Pyrrolidine-2-carboxylic acid (2'-methanesulfonyl-biphenyl-4-yl)-amide (0.61 g, 1.8 mmol) was dissolved in 20 mL THF, added diisopropylethylamine (0.92 mL, 5.3 mmol), followed by 4-chlorophenylisocyanate (270 mg, 1.7 mmol). Stirred reaction at ambient temperature for one hour then concentrated.

- 25 Crystallized from EtOAc in hexanes and filtered to yield title compound as a white powder. (0.71 g, 80%) APCI (AP-): 496.1, 498.1 (M-H)⁻.

-196-

Example 20

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3-methyl-biphenyl-4-yl)-amide]



5

Step 1: Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

DL-proline (10 g, 86.9 mmol) was dissolved in 120 mL of THF: H₂O (2:1), added 87 mL of a 2M NaOH solution followed by Boc₂O (24.6 g, 113 mmol). Stirred at ambient temperature overnight. Removed THF in vacuo, acidified water to pH 3 with citric acid, extracted twice with EtOAc, washed organics with water, brine, dried with MgSO₄, filtered and concentrated to yield title compound as a white solid. (18.7 g, quant.) APCI (AP-): 214.1 (M-H)⁻.

10

Step 2: 2-(4-Bromo-2-methyl-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

15

Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (2 g, 9.3 mmol) was dissolved in 100 mL dry diethyl ether under Ar, cooled in an ice water bath, added dry pyridine (4.5 mL, 55 mmol) followed by the dropwise addition of oxalyl chloride (2.4 mL, 27 mmol). A precipitate forms immediately. The reaction was stirred vigorously at 0°C for one hour, then at ambient temperature for one hour. Added 100 mL diethyl ether, filtered off solids washing with diethyl ether. Concentrated the filtrates to an off white oil. Redissolved (0.95 g, 4.0 mmol) oil in 40 mL dry DCM under Ar, cooled to 0°C, added 1.3 mL pyridine followed by 4-bromo-2-methylaniline (0.75 g, 4.0 mmol). The reaction was allowed to warm to ambient temperature, stirred overnight then concentrated. Redissolved in 100 mL EtOAc, washed with 1 N HCl (3x50 mL), brine (100 mL), dried with MgSO₄,

20

25

-197-

filtered and concentrated to yield title compound. (1.09 g, 70%) APCI (AP-): 381.0, 383.0 (M-H)⁻.

Step 3: 2-(3-Methyl-2'-methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

2-(4-Bromo-2-methyl-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (1.07 g, 2.7 mmol), 2-(methylthio)benzene boronic acid (0.56 g, 3.3 mmol), and tetrabutylammonium bromide (322 mg, 0.13 mmol) were combined in 20 mL toluene, added 2.8 mL of a 2M aqueous Na₂CO₃ solution followed by tetrakis(triphenylphosphine) palladium(0) (0.161 g, 0.13 mmol). Heated reaction at reflux 4 hours, cooled, concentrated, redissolved in EtOAc (250 mL), washed with water (3x200 mL), brine (200 mL), dried with MgSO₄, filtered, and concentrated. Purified on a silica gel column eluted with 20% EtOAc in hexanes. Combined and concentrated pure fractions to yield title compound. (0.59 g, 50%) APCI (AP-): 425.1 (M-H)⁻.

Step 4: 2-(2'-Methanesulfonyl-3-methyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

2-(3-Methyl-2'-methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (0.57 g, 1.3 mmol) was dissolved in 10 mL EtOAc, added added *m*-chloroperoxybenzoic acid (1.76 g, 6.1 mmol) in one portion and stirred at ambient temperature for 3 hours. Reaction not complete. Concentrated and redissolved in 10 mL acetonitrile and added oxone (1.5 g, 2.4 mmol) in one portion and stirred at ambient temperature overnight and then concentrated. Redissolved in 100 mL EtOAc and 100 mL sat. NaHCO₃, separated layers, washed organics with brine, dried with MgSO₄, filtered and concentrated. Purified on a silica gel column eluted with 20% moving to 40% EtOAc in hexanes. Combined and concentrated pure fractions to yield title compound. (0.44 g, 72%) APCI (AP-): 457.1 (M-H)⁻.

-198-


Step 5: Pyrrolidine-2-carboxylic acid (2'-methanesulfonyl-3-methyl-biphenyl-4-yl)-amide

2-(2'-Methanesulfonyl-3-methyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (0.4 g, 0.87 mmol) was dissolved in 15 mL DCM and 5 mL trifluoroacetic acid and stirred at ambient temperature for 2 hour and concentrated. Redissolved in 100 mL EtOAc and 100 mL sat. NaHCO₃, separated layers, washed organics with brine, dried with MgSO₄, filtered and concentrated to yield title compound as an oil (0.29 g, 93%) APCI (AP-): 357.1 (M-H)⁻.

10 **Step 6: Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3-methyl-biphenyl-4-yl)-amide]**

Pyrrolidine-2-carboxylic acid (2'-methanesulfonyl-3-methyl-biphenyl-4-yl)-amide (0.29 g, 0.80 mmol) was dissolved in 10 mL THF, added diisopropylethylamine (0.42 mL, 2.4 mmol), followed by 4-chlorophenylisocyanate (124 mg, 0.80 mmol). Stirred reaction at ambient temperature for one hour then concentrated. Purified on a silica gel column eluted with 20% moving to 40% EtOAc in hexanes. Combined and concentrated pure fractions to yield title compound. (0.32 g, 78%) APCI (AP-): 510.1, 512.1 (M-H)⁻.

20

Example 21**(2R,4R)-4-methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]**

-199-

Step 1: (2R,4R)-4-hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

Cis-4-hydroxy-D-proline (1.1 g, 8.4 mmol) was dissolved in 18 mL of THF/ H₂O (2:1), added 2M aqueous NaOH solution (6.3 mL, 13 mmol), followed
5 by Boc₂O. Reaction was stirred at ambient temperature over the weekend. Solution was taken up in 150 mL EtOAc, washed with 10% HCl (3x100 mL), dried organics with MgSO₄, filtered and concentrated to yield title compound as a white foam. (1.5 g, 77%) APCI (AP⁻): 230.1 (M-H)⁻.

10 Step 2: (2R,4R)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (1.5 g, 6.5 mmol) was dissolved in 2 mL DMF, added imidazole (1.1 g, 16 mmol), dimethylaminopyridine (80mg, 0.65 mmol), and *tert*-butyldimethylsilyl chloride
15 (1.08 g, 7.1 mmol). Stirred at ambient temperature overnight, dissolved in 150 mL EtOAc, washed with 10% citric acid (3x100 mL), water (2x100 mL), brine, dried with MgSO₄, filtered, and concentrated. Redissolved in 50 mL CHCl₃ and reconcentrated to yield title compound as a clear oil. (1.54 g, 69%) APCI (AP⁻): 344.2 (M-H)⁻.

20

Step 3: (2R,4R)-2-(4-bromo-2-fluoro-phenylcarbamoyl)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

(2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (1.54 g, 4.5 mmol) was dissolved in 50 mL diethyl ether
25 under an Ar atmosphere, cooled in an ice water bath, added dry pyridine (1.4 mL, 17.8 mmol), followed by the dropwise addition of oxalyl chloride (0.58 mL, 6.7 mmol). Reaction was stirred at 0°C for 1 hour, then at ambient temperature for 1 hour. Diluted reaction with 50 mL diethyl ether, filtered off solids and concentrated filtrates. Redissolved filtrate in 30 mL dry DCM under an Ar
30 atmosphere, cooled in an ice water bath, added 1 mL dry pyridine followed by 4-bromo-2-fluoroaniline (0.85 g, 4.5 mmol). Stirred reaction at ambient temperature

-200-

for 20 hours. Concentrated solution, redissolved in EtOAc (150 mL), washed with 10% citric acid (3x100 mL), water (2x100 mL), brine, dried with MgSO₄, filtered and concentrated. Purified on a silica gel column eluting with an automated system with gradient of 0-60% EtOAc in hexanes over 40 minutes.

- 5 Combined and concentrated pure fractions to yield title compound as a clear oil. (0.96 g, 42%) APCI (AP-): 515.1, 517.1 (M-H)⁻.

Step 4: (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

- 10 (2R,4R)-2-(4-Bromo-2-fluoro-phenylcarbamoyl)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (0.95 g, 1.84 mmol), 2-(methylthio)benzene boronic acid (0.37 g, 2.2 mmol), and tetrabutylammonium bromide (30mg, 0.09 mmol) were combined in 15 mL toluene, added 2 mL of a
- 15 2M aqueous Na₂CO₃ solution followed by tetrakis(triphenyl)phosphine palladium(0) (0.11 g, 0.09 mmol). Heated reaction at reflux for 5 hours, cooled, dissolved in EtOAc (100 mL), washed with water (3x50 mL), brine (50 mL), dried with MgSO₄, filtered, and concentrated. Purified on a silica gel column eluted
- 20 minutes. Combined and concentrated pure fractions to yield title compound as a light yellow foam. (0.77 g, 74%) APCI (AP-): 559.3 (M-H)⁻.

Step 5: (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

- 25 (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (0.84 g, 1.35 mmol) was dissolved in 20 mL EtOAc, added added *m*-chloroperoxybenzoic acid (1.6 g, 5.4 mmol) in one portion and stirred at ambient temperature for 3
- 30 hours. A 10% aqueous solution (50 mL) of Na₂S₂O₃ was added to the vigorously stirring reaction in order to quench peroxide. After 20 minutes the solution was

-201-

diluted with 100 mL EtOAc, separated layers, washed organics with sat. NaHCO_3 (3x100 mL), water (2x100 mL), brine, dried with MgSO_4 , filtered and concentrated to yield title compound as a light brown foam. (0.8 g, 100%) APCI (AP-): 591.3 (M-H)⁻.

5

Step 6: (2R,4R)-4-hydroxy-pyrrolidine-2-carboxylic acid (3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide

(2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (0.8 g, 1.35 mmol) was dissolved in 30 mL DCM, added 10 mL trifluoroacetic acid and stirred reaction at ambient temperature for 90 minutes. Solution was concentrated to yield title compound as an amber oil. (100% crude yield) APCI (AP-): 434.1 (M-H)⁻.

15 **Step 7: (2R,4R)-4-hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chlorophenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]**

(2R,4R)-4-Hydroxy-pyrrolidine-2-carboxylic acid (3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide (0.51 g, 1.35 mmol) was dissolved in 20 mL THF, added diisopropylethylamine (0.94 mL, 5.4 mmol) followed by 4-chlorophenylisocyanate (0.21 g, 1.35 mmol). Stirred at ambient temperature for 2 hours. The reaction was dissolved in 100 mL EtOAc, washed with 10% citric acid (3x50 mL), water (2x50 mL), brine, dried with MgSO_4 , filtered and concentrated. Purified using an automated system with gradient of 50-100% EtOAc in hexanes over 40 minutes. Compound was still eluting at end of run. Switched gradient to 25 0-15% MeOH in EtOAc over 40 minutes. Combined and concentrated pure fractions to yield title compound as a white solid. (0.54 g, 75%) APCI (AP+): 530.1, 532.1(M+H)⁺.

30 **Step 8: (2R,4R)-4-methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chlorophenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]**

-202-


(2R,4R)-4-hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] (0.070 g, 0.13 mmol) was dissolved in 1.5 mL dry acetonitrile under an Argon atmosphere. Freshly prepared Ag₂O (45mg, 0.19 mmol) was added followed by methyl iodide (9 mL, 0.14 mmol). The reaction was stirred for 2 hours with no reaction seen by HPLC. Added 10 equivalents each of Ag₂O and methyl iodide. After 2 hours the reaction was complete by HPLC. The solids were filtered through celite, washing with ethyl acetate. The filtrates were concentrated, then purified by preparatory HPLC. Pure fractions were combined and lyophilized to yield title compound.

10 (28mg, 40%) APCI (AP-): 544.0, 546.1 (M-H)⁻.

Example 22

(2R)-4-oxo-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]

15



Step 1: (2R,4R)-4-hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

Cis-4-hydroxy-D-proline (15 g, 114 mmol) was dissolved in 150 mL of THF/ H₂O (2:1), added 2M aqueous NaOH solution (86μmL, 172 mmol), followed by Boc₂O (27 g, 126 mmol). Reaction was stirred at ambient temperature overnight. The solution was acidified with 10% citric acid then extracted with EtOAc (2x250 mL), washed with water, brine, dried organics with MgSO₄, filtered and concentrated to yield title compound. (16 g, 61%) APCI

25 (AP-): 230.1 (M-H)⁻.

-203-

Step 2: (2R,4R)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester
5 (16 g, 69 mmol) was dissolved in 100 mL DMF, added imidazole (12 g, 173 mmol), dimethylaminopyridine (0.85 g, 6.9 mmol), and *tert*-butyldimethylsilyl chloride (12.5 g, 83 mmol). Stirred at ambient temperature overnight, dissolved in 350 mL EtOAc, washed with 10% citric acid (3x200 mL), water (2x200 mL), brine, dried with MgSO₄, filtered, and concentrated to yield title compound. (20
10 g, 84%) APCI (AP-): 344.2 (M-H)⁻.

Step 3: (2R,4R)-2-(4-bromo-2-fluoro-phenylcarbamoyl)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

(2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-pyrrolidine-1,2-dicarboxylic
15 acid 1-*tert*-butyl ester (4.45 g, 12.9 mmol) was dissolved in 50 mL dry CHCl₃ under an Ar atmosphere, added 4-bromo-2-fluoroaniline (2.45 g, 12.9 mmol), EEDQ (3.8 g, 15 mmol), and triethylamine (2.6 mL, 19.3 mmol). Reaction was refluxed for 6 hours, cooled and concentrated. Redissolved in EtOAc (250 mL), washed with 10% HCl (2x200 mL), 0.1M NaOH (2x200 mL), water, brine, dried
20 with MgSO₄, filtered and concentrated to yield title compound. (5.5 g, 82%)
APCI (AP-): 515.1, 517.1 (M-H)⁻.

**Step 4: (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-
25 butyl ester**

(2R,4R)-2-(4-Bromo-2-fluoro-phenylcarbamoyl)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (5.7 g, 10 mmol), 2-(methylthio)benzene boronic acid (2.1 g, 12.6 mmol), and tetrabutylammonium bromide (0.17 g, 0.52 mmol) were combined in 50 mL toluene, added 10 mL of a
30 2M aqueous Na₂CO₃ solution followed by tetrakis(triphenyl)phosphine palladium(0) (0.61 g, 0.52 mmol). Heated reaction at reflux for 4 hours, cooled,

-204-

dissolved in EtOAc (250 mL), washed with water (200 mL), brine, dried with MgSO₄, filtered, and concentrated in the presence of 15 g coarse silica. Slurried silica in mobile phase and loaded onto a silica gel column, eluted with 25% EtOAc in hexanes. Combined and concentrated fractions to yield title compound.

5 (4.1 g, 70%) APCI (AP-): 559.3 (M-H)⁻.

Step 5: (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

10 (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (4.1 g, 7.3 mmol) was dissolved in 20 mL EtOAc, added added *m*-chloroperoxybenzoic acid (7.2 g, 29 mmol) in one portion and stirred at ambient temperature for 3 hours. A 10% aqueous solution (50 mL) of Na₂S₂O₃ was added to the vigorously stirring
15 reaction in order to quench peroxide. After 20 minutes the solution was diluted with 100 mL EtOAc, separated layers, washed organics with sat. NaHCO₃ (3x100 mL), water (2x100 mL), brine, dried with MgSO₄, filtered and concentrated to yield title compound. (4.3 g, 99%) APCI (AP-): 591.3 (M-H)⁻.

20 **Step 6: (2R,4R)-4-hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chlorophenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]**

(2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (3.25 g, 5.5 mmol) was dissolved in 50 mL DCM, added 25 mL
25 trifluoroacetic acid and stirred reaction at ambient temperature for 90 minutes. Solution was concentrated. The resulting oil was dissolved in 75 mL THF, added triethylamine (3.8 mL, 5.4 mmol) followed by 4-chlorophenylisocyanate (0.84 g, 5.5 mmol). Stirred at ambient temperature for 2 hours. The reaction was concentrated and purified on a silica gel column eluted with straight ethyl acetate
30 initially moving to 5% MeOH in ethyl acetate. Combined and concentrated pure fractions to yield title compound. (2.5 g, 85%) APCI (AP-): 530.1, 532.1(M-H)⁻.

-205-


Step 7: (2R)-4-oxo-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]

(2R,4R)-4-hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] (0.78 g, 1.5 mmol) was dissolved in 10 mL dry DCM under an Argon atmosphere, added 0.5 g celite, 0.5 g of 4 angstrom crushed molecular sieves, and pyridinium chlorochromate (0.47 g, 2.2 mmol). Stirred at ambient temperature for 6 hours, reaction appears to have stalled. Added 1 g pyridinium chlorochromate and stirred at RT overnight. HPLC did not change. Added 40 mL of a 1:1 mixture of hexanes:Et₂O. Filtered through a filter funnel with silica gel. Washed with DCM. Compound did not elute. Washed with DCM:MeOH (90:10). Concentrated filtrates. Suspect pyridinium chlorochromate also eluted judged by dark color. Redissolved in DCM, washed with sat. NaHCO₃, dried DCM with MgSO₄, filtered and concentrated in vacuo. Purified on a silica gel column using an automated ISCO system with gradient of 50-100% EtOAc in hexanes over 30 minutes, then 100% EtOAc for 10 minutes. Combined and concentrated pure fractions, redissolved in acetonitrile and lyophilized to yield title compound. (0.12 g, 15%)

20

Example 23

(2R,4S)-4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]-2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]



25

-206-

Step 1: (2R,4R)-4-hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

Cis-4-hydroxy-D-proline (15 g, 114 mmol) was dissolved in 150 mL of THF/H₂O (2:1), added 2M aqueous NaOH solution (86 mL, 172 mmol),
5 followed by Boc₂O (27 g, 126 mmol). Reaction was stirred at ambient temperature overnight. The solution was acidified with 10% citric acid then extracted with EtOAc (2x250 mL), washed with water, brine, dried organics with MgSO₄, filtered and concentrated to yield title compound. (16 g, 61%) APCI (AP-): 230.1 (M-H)⁻.

10

Step 2: (2R,4R)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (16 g, 69 mmol) was dissolved in 100 mL DMF, added imidazole (12 g, 173
15 mmol), dimethylaminopyridine (0.85 g, 6.9 mmol), and *tert*-butyldimethylsilyl chloride (12.5 g, 83 mmol). Stirred at ambient temperature overnight, dissolved in 350 mL EtOAc, washed with 10% citric acid (3x200 mL), water (2x200 mL), brine, dried with MgSO₄, filtered, and concentrated to yield title compound. (20 g, 84%) APCI (AP-): 344.2 (M-H)⁻.

20

Step 3: (2R,4R)-2-(4-bromo-2-fluoro-phenylcarbamoyl)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

(2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (4.45 g, 12.9 mmol) was dissolved in 50 mL dry CHCl₃
25 under an Ar atmosphere, added 4-bromo-2-fluoroaniline (2.45 g, 12.9 mmol), EEDQ (3.8 g, 15 mmol), and triethylamine (2.6 mL, 19.3 mmol). Reaction was refluxed for 6 hours, cooled and concentrated. Redissolved in EtOAc (250 mL), washed with 10% HCl (2x200 mL), 0.1M NaOH (2x200 mL), water, brine, dried with MgSO₄, filtered and concentrated to yield title compound. (5.5 g, 82%)

30 APCI (AP-): 515.1, 517.1 (M-H)⁻.

-207-

Step 4: (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

- 5 (2R,4R)-2-(4-Bromo-2-fluoro-phenylcarbamoyl)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (5.7 g, 10 mmol), 2-(methylthio)benzene boronic acid (2.1 g, 12.6 mmol), and tetrabutylammonium bromide (0.17 g, 0.52 mmol) were combined in 50 mL toluene, added 10 mL of a 2M aqueous Na₂CO₃ solution followed by tetrakis(triphenyl)phosphine
- 10 palladium(0) (0.61 g, 0.52 mmol). Heated reaction at reflux for 4 hours, cooled, dissolved in EtOAc (250 mL), washed with water (200 mL), brine, dried with MgSO₄, filtered, and concentrated in the presence of 15 g coarse silica. Slurried silica in mobile phase and loaded onto a silica gel column, eluted with 25% EtOAc in hexanes. Combined and concentrated fractions to yield title compound.
- 15 (4.1 g, 70%) APCI (AP-): 559.3 (M-H)⁻.

Step 5: (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

- 20 (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (4.1 g, 7.3 mmol) was dissolved in 20 mL EtOAc, added added *m*-chloroperoxybenzoic acid (7.2 g, 29 mmol) in one portion and stirred at ambient temperature for 3 hours. A 10% aqueous solution (50 mL) of Na₂S₂O₃ was added to the vigorously stirring
- 25 reaction in order to quench peroxide. After 20 minutes the solution was diluted with 100 mL EtOAc, separated layers, washed organics with sat. NaHCO₃ (3x100 mL), water (2x100 mL), brine, dried with MgSO₄, filtered and concentrated to yield title compound. (4.3 g, 99%) APCI (AP-): 591.3 (M-H)⁻.

- 30 **Step 6: (2R,4R)-4-hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chlorophenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]**

-208-


- (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (3.25 g, 5.5 mmol) was dissolved in 50 mL DCM, added 25 mL trifluoroacetic acid and stirred reaction at ambient temperature for 90 minutes.
- 5 Solution was concentrated. The resulting oil was dissolved in 75 mL THF, added triethylamine (3.8 mL, 5.4 mmol) followed by 4-chlorophenylisocyanate (0.84 g, 5.5 mmol). Stirred at ambient temperature for 2 hours. The reaction was concentrated and purified on a silica gel column eluted with straight ethyl acetate initially moving to 5% MeOH in ethyl acetate. Combined and concentrated pure
- 10 fractions to yield title compound. (2.5 g, 85%) APCI (AP-): 530.1, 532.1(M-H)⁻.

- Step 7: (2R,4S)-4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chlorophenyl)-amide]-2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]**
- (2R,4R)-4-hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chlorophenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] (0.1 g, 0.19 mmol) in 3 mL dry DCM was added to a solution of [Bis(2-methoxyethyl)amino]-sulfur trifluoride (38 μ l, 0.21 mmol) in 3 mL dry DCM at -78°C under Argon. Stirred at -78°C for 30 minutes then allowed to warm to room temperature. Stirred for one hour then concentrated reaction. Purified by
- 15 preparatory HPLC, combined and lyophilized fractions to yield title compound. (0.020 g, 20%) APCI (AP-): 532.1, 534.1(M-H)⁻.
- 20

Example 24

- 25 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-p-tolylamide

-209-



Step 1: (2R,4R)-4-hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

Cis-4-hydroxy-D-proline (15 g, 114 mmol) was dissolved in 150 mL of
 5 THF/ H₂O (2:1), added 2M aqueous NaOH solution (86 mL, 172 mmol),
 followed by Boc₂O (27 g, 126 mmol). Reaction was stirred at ambient
 temperature overnight. The solution was acidified with 10% citric acid then
 extracted with EtOAc (2x250 mL), washed with water, brine, dried organics with
 MgSO₄, filtered and concentrated to yield title compound. (16 g, 61%) APCI
 10 (AP-): 230.1 (M-H)⁻.

Step 2: (2R,4R)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester
 15 (16 g, 69 mmol) was dissolved in 100 mL DMF, added imidazole (12 g, 173
 mmol), dimethylaminopyridine (0.85 g, 6.9 mmol), and *tert*-butyldimethylsilyl
 chloride (12.5 g, 83 mmol). Stirred at ambient temperature overnight, dissolved in
 350 mL EtOAc, washed with 10% citric acid (3x200 mL), water (2x200 mL),
 brine, dried with MgSO₄, filtered, and concentrated to yield title compound. (20
 20 g, 84%) APCI (AP-): 344.2 (M-H)⁻.

Step 3: (2R,4R)-2-(4-bromo-2-fluorophenylcarbamoyl)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

(2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-pyrrolidine-1,2-dicarboxylic
 25 acid 1-*tert*-butyl ester (4.45 g, 12.9 mmol) was dissolved in 50 mL dry CHCl₃

-210-

under an Ar atmosphere, added 4-bromo-2-fluoroaniline (2.45 g, 12.9 mmol), EEDQ (3.8 g, 15 mmol), and triethylamine (2.6 mL, 19.3 mmol). Reaction was refluxed for 6 hours, cooled and concentrated. Redissolved in EtOAc (250 mL), washed with 10% HCl (2x200 mL), 0.1 M NaOH (2x200 mL), water, brine, dried
5 with MgSO₄, filtered and concentrated to yield title compound. (5.5 g, 82%)
APCI (AP⁻): 515.1, 517.1 (M-H)⁻.

Step 4: (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester
10 (2R,4R)-2-(4-Bromo-2-fluoro-phenylcarbamoyl)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (5.7 g, 10 mmol), 2-(methylthio)benzene boronic acid (2.1 g, 12.6 mmol), and tetrabutylammonium bromide (0.17 g, 0.52 mmol) were combined in 50 mL toluene, added 10 mL of a
15 2M aqueous Na₂CO₃ solution followed by tetrakis(triphenyl)phosphine palladium(0) (0.61 g, 0.52 mmol). Heated reaction at reflux for 4 hours, cooled, dissolved in EtOAc (250 mL), washed with water (200 mL), brine, dried with MgSO₄, filtered, and concentrated in the presence of 15 g coarse silica. Slurried
silica in mobile phase and loaded onto a silica gel column, eluted with 25%
20 EtOAc in hexanes. Combined and concentrated fractions to yield title compound. (4.1 g, 70%) APCI (AP⁻): 559.3 (M-H)⁻.

Step 5: (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester
25 (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (4.1 g, 7.3 mmol) was dissolved in 20 mL EtOAc, added added *m*-chloroperoxybenzoic acid (7.2 g, 29 mmol) in one portion and stirred at ambient temperature for 3 hours. A
30 10% aqueous solution (50 mL) of Na₂S₂O₃ was added to the vigorously stirring reaction in order to quench peroxide. After 20 minutes the solution was diluted

-211-


with 100 mL EtOAc, separated layers, washed organics with sat. NaHCO_3 (3x100 mL), water (2x100 mL), brine, dried with MgSO_4 , filtered and concentrated to yield title compound. (4.3 g, 99%) APCI (AP-): 591.3 (M-H)⁻.

5 Step 6: (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-p-tolylamide

(2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (0.35 g, 0.59 mmol) was dissolved in 10 mL DCM, added 5 mL
 10 trifluoroacetic acid and stirred reaction at ambient temperature for 2 hours. Solution was concentrated. The resulting oil was dissolved in 20 mL THF, added triethylamine (0.41 mL, 3 mmol) followed by p-tolylisocyanate (0.079 g, 0.59 mmol). Stirred at ambient temperature for 2 hours. The reaction was concentrated and purified on a silica gel column eluted using an automated ISCO
 15 system with gradient of 0-10% MeOH in ethyl acetate over 40 minutes. Concentrated fractions then redissolved in acetonitrile/water and lyophilized to yield title compound. (0.17 g, 56%) APCI (AP-): 510.1(M-H)⁻.

Example 25

20 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(4-fluoro-phenyl)-amide]



Step 1: (2R,4R)-4-hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

25 Cis-4-hydroxy-D-proline (15 g, 114 mmol) was dissolved in 150 mL of THF/ H_2O (2:1), added 2M aqueous NaOH solution (86 mL, 172 mmol),

-212-

followed by Boc_2O (27 g, 126 mmol). Reaction was stirred at ambient temperature overnight. The solution was acidified with 10% citric acid then extracted with EtOAc (2x250 mL), washed with water, brine, dried organics with MgSO_4 , filtered and concentrated to yield title compound. (16 g, 61%) APCI

5 (AP-): 230.1 (M-H)⁻.

Step 2: (2R,4R)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

10 (16 g, 69 mmol) was dissolved in 100 mL DMF, added imidazole (12 g, 173 mmol), dimethylaminopyridine (0.85 g, 6.9 mmol), and *tert*-butyldimethylsilyl chloride (12.5 g, 83 mmol). Stirred at ambient temperature overnight, dissolved in 350 mL EtOAc, washed with 10% citric acid (3x200 mL), water (2x200 mL), brine, dried with MgSO_4 , filtered, and concentrated to yield title compound. (20
15 g, 84%) APCI (AP-): 344.2 (M-H)⁻.

Step 3: (2R,4R)-2-(4-bromo-2-fluoro-phenylcarbamoyl)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

(2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-pyrrolidine-1,2-dicarboxylic

20 acid 1-*tert*-butyl ester (4.45 g, 12.9 mmol) was dissolved in 50 mL dry CHCl_3 under an Ar atmosphere, added 4-bromo-2-fluoroaniline (2.45 g, 12.9 mmol), EEDQ (3.8 g, 15 mmol), and triethylamine (2.6 mL, 19.3 mmol). Reaction was refluxed for 6 hours, cooled and concentrated. Redissolved in EtOAc (250 mL), washed with 10% HCl (2x200 mL), 0.1M NaOH (2x200 mL), water, brine, dried
25 with MgSO_4 , filtered and concentrated to yield title compound. (5.5 g, 82%)
APCI (AP-): 515.1, 517.1 (M-H)⁻.

**Step 4: (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-
30 butyl ester**

-213-

(2R,4R)-2-(4-Bromo-2-fluoro-phenylcarbamoyl)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (5.7 g, 10 mmol), 2-(methylthio)benzene boronic acid (2.1 g, 12.6 mmol), and tetrabutylammonium bromide (0.17 g, 0.52 mmol) were combined in 50 mL toluene, added 10 mL of a
5 2M aqueous Na₂CO₃ solution followed by tetrakis(triphenyl)phosphine palladium(0) (0.61 g, 0.52 mmol). Heated reaction at reflux for 4 hours, cooled, dissolved in EtOAc (250 mL), washed with water (200 mL), brine, dried with MgSO₄, filtered, and concentrated in the presence of 15 g coarse silica. Slurried
10 EtOAc in hexanes. Combined and concentrated fractions to yield title compound. (4.1 g, 70%) APCI (AP-): 559.3 (M-H)⁻.

Step 5: (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester
15

(2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methylsulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (4.1 g, 7.3 mmol) was dissolved in 20 mL EtOAc, added added *m*-chloroperoxybenzoic acid (7.2 g, 29 mmol) in one portion and stirred at ambient temperature for 3 hours. A
20 10% aqueous solution (50 mL) of Na₂S₂O₃ was added to the vigorously stirring reaction in order to quench peroxide. After 20 minutes the solution was diluted with 100 mL EtOAc, separated layers, washed organics with sat. NaHCO₃ (3x100 mL), water (2x100 mL), brine, dried with MgSO₄, filtered and concentrated to yield title compound. (4.3 g, 99%) APCI (AP-): 591.3 (M-H)⁻.

25

Step 6: (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(4-fluoro-phenyl)-amide]

(2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl
30 ester (0.35 g, 0.59 mmol) was dissolved in 10 mL DCM, added 5 mL trifluoroacetic acid and stirred reaction at ambient temperature for 2 hours.


-214-

Solution was concentrated. The resulting oil was dissolved in 20 mL THF, added triethylamine (0.41 mL, 3 mmol) followed by 4-fluorophenylisocyanate (0.081 g, 0.59 mmol). Stirred at ambient temperature for 2 hours. The reaction was concentrated and purified on a silica gel column eluted using an automated ISCO system with gradient of 80-100% ethyl acetate in hexanes for 20 minutes then straight ethyl acetate for 20 minutes. Concentrated fractions then redissolved in acetonitrile/water and lyophilized to yield title compound. (0.175 g, 57%) APCI (AP⁻): 514.1(M-H)⁻.

10

Example 26

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]



15

Step 1: (2R,4R)-4-hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

Cis-4-hydroxy-D-proline (15 g, 114 mmol) was dissolved in 150 mL of THF/ H₂O (2:1), added 2M aqueous NaOH solution (86 mL, 172 mmol), followed by Boc₂O (27 g, 126 mmol). Reaction was stirred at ambient temperature overnight. The solution was acidified with 10% citric acid then extracted with EtOAc (2x250 mL), washed with water, brine, dried organics with MgSO₄, filtered and concentrated to yield title compound. (16 g, 61%) APCI (AP⁻): 230.1 (M-H)⁻.

Step 2: (2R,4R)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

-215-

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (16 g, 69 mmol) was dissolved in 100 mL DMF, added imidazole (12 g, 173 mmol), dimethylaminopyridine (0.85 g, 6.9 mmol), and *tert*-butyldimethylsilyl chloride (12.5 g, 83 mmol). Stirred at ambient temperature overnight, dissolved in 5 350 mL EtOAc, washed with 10% citric acid (3x200 mL), water (2x200 mL), brine, dried with MgSO₄, filtered, and concentrated to yield title compound. (20 g, 84%) APCI (AP-): 344.2 (M-H)⁻.

Step 3: (2R,4R)-2-(4-bromo-2-fluoro-phenylcarbamoyl)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1-carboxylic acid *tert*-butyl ester
10 (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (4.45 g, 12.9 mmol) was dissolved in 50 mL dry CHCl₃ under an Ar atmosphere, added 4-bromo-2-fluoroaniline (2.45 g, 12.9 mmol), EEDQ (3.8 g, 15 mmol), and triethylamine (2.6 mL, 19.3 mmol). Reaction was 15 refluxed for 6 hours, cooled and concentrated. Redissolved in EtOAc (250 mL), washed with 10% HCl (2x200 mL), 0.1M NaOH (2x200 mL), water, brine, dried with MgSO₄, filtered and concentrated to yield title compound. (5.5 g, 82%) APCI (AP-): 515.1, 517.1 (M-H)⁻.

Step 4: (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methylsulfanyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester
20 (2R,4R)-2-(4-Bromo-2-fluoro-phenylcarbamoyl)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (5.7 g, 10 mmol), 2- 25 (methylthio)benzene boronic acid (2.1 g, 12.6 mmol), and tetrabutylammonium bromide (0.17 g, 0.52 mmol) were combined in 50 mL toluene, added 10 mL of a 2M aqueous Na₂CO₃ solution followed by tetrakis(triphenyl)phosphine palladium(0) (0.61 g, 0.52 mmol). Heated reaction at reflux for 4 hours, cooled, dissolved in EtOAc (250 mL), washed with water (200 mL), brine, dried with 30 MgSO₄, filtered, and concentrated in the presence of 15 g coarse silica. Slurried silica in mobile phase and loaded onto a silica gel column, eluted with 25%

-216-

EtOAc in hexanes. Combined and concentrated fractions to yield title compound. (4.1 g, 70%) APCI (AP-): 559.3 (M-H)⁻.

Step 5: (2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-

5 **methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester**

(2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methylsulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (4.1 g, 7.3 mmol) was dissolved in 20 mL EtOAc, added added *m*-chloroperoxybenzoic acid (7.2 g, 29 mmol) in one portion and stirred at ambient temperature for 3 hours. A 10% aqueous solution (50 mL) of Na₂S₂O₃ was added to the vigorously stirring reaction in order to quench peroxide. After 20 minutes the solution was diluted with 100 mL EtOAc, separated layers, washed organics with sat. NaHCO₃ (3x100 mL), water (2x100 mL), brine, dried with MgSO₄, filtered and concentrated to 15 yield title compound. (4.3 g, 99%) APCI (AP-): 591.3 (M-H)⁻.

Step 6: (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]

(2R,4R)-4-(*tert*-Butyl-dimethyl-silanyloxy)-2-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (0.35 g, 0.59 mmol) was dissolved in 10 mL DCM, added 5 mL trifluoroacetic acid and stirred reaction at ambient temperature for 2 hours. Solution was concentrated. The resulting oil was dissolved in 15 mL DMF, added triethylamine (0.41 mL, 3 mmol) followed by (5-Chloro-pyridin-2-yl)-carbamic acid 4-nitro-phenyl ester (0.17 g, 0.059 mmol; see Example 3, step 3a). Stirred at 25 50°C for 2 hours. The reaction was cooled, dissolved in 100 mL ethyl acetate, washed with sat. NaHCO₃ (3x50 mL), 10% citric acid (2x50 mL), brine, dried with MgSO₄, filtered and concentrated. Purified on a silica gel column eluted using an automated ISCO system with gradient of 0-10% MeOH in ethyl acetate 30 over 40 minutes. Concentrated fractions then redissolved in acetonitrile/water and


-217-

lyophilized to yield title compound. (0.20 g, 64%) APCI (AP-): 531.1, 533.0(M-H)⁻.

Example 27

5

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3-trifluoromethyl-biphenyl-4-yl)-amide]



10 Step 1: Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester

DL-proline (10 g, 86.9 mmol) was dissolved in 120 mL of THF: H₂O (2:1), added 87 mL of a 2M NaOH solution followed by Boc₂O (24.6 g, 113 mmol). Stirred at ambient temperature overnight. Removed THF in vacuo, acidified water to pH 3 with citric acid, extracted twice with EtOAc, washed
15 organics with water, brine, dried with MgSO₄, filtered and concentrated to yield title compound as a white solid. (18.7 g, quant.) APCI (AP-): 214.1 (M-H)⁻.

Step 2: 2-(4-Bromo-2-trifluoromethyl-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

20 Pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (4 g, 18.6 mmol) was dissolved in 200 mL dry diethyl ether under Ar, cooled in an ice water bath, added dry pyridine (9 mL, 111 mmol) followed by the dropwise addition of oxalyl chloride (4.8 mL, 54 mmol). A precipitate forms immediately. The reaction was stirred vigorously at 0°C for one hour, then at ambient temperature for one hour.
25 Added 100 mL diethyl ether, filtered off solids washing with diethyl ether.

-218-

Concentrated the filtrates to an off white oil. Redissolved (2 g, 8.5 mmol) oil in 40 mL dry DCM under Ar, cooled to 0°C, added 2.8 mL pyridine followed by 4-bromo-2-trifluoromethylaniline (2.05 g, 8.6 mmol). The reaction was allowed to warm to ambient temperature, stirred overnight then concentrated. Redissolved in 100 mL EtOAc, washed with 1 N HCl (3x50 mL), brine (100 mL), dried with MgSO₄, filtered and concentrated to yield title compound. (1.98 g, 53%) APCI (AP-): 435.0, 437.0 (M-H)⁻.

Step 3: 2-(2'-Methylsulfanyl-3-trifluoromethyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

2-(4-Bromo-2-trifluoromethyl-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (1.92 g, 4.4 mmol), 2-(methylthio)benzene boronic acid (0.88 g, 5.2 mmol), and tetrabutylammonium bromide (70 mg, 0.22 mmol) were combined in 40 mL toluene, added 4.4 mL of a 2M aqueous Na₂CO₃ solution followed by tetrakis(triphenylphosphine) palladium(0) (0.253 g, 0.22 mmol). Heated reaction at reflux 4 hours, cooled, concentrated, redissolved in EtOAc (250 mL), washed with water (3x200 mL), brine (200 mL), dried with MgSO₄, filtered, and concentrated. Purified on a silica gel column eluted with 20% EtOAc in hexanes. Combined and concentrated pure fractions to yield title compound. (1.8 g, 85%) APCI (AP-): 479.1 (M-H)⁻.

Step 4: 2-(2'-Methanesulfonyl-3-trifluoromethyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester

2-(2'-Methylsulfanyl-3-trifluoromethyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (1.8 g, 3.7 mmol) was dissolved in 20 mL acetonitrile, and added oxone (4.6 g, 6.8 mmol) in one portion and stirred at ambient temperature 4 days and then concentrated. Redissolved in 100 mL EtOAc and 100 mL sat. NaHCO₃, separated layers, washed organics with brine, dried with MgSO₄, filtered and concentrated. Purified on a silica gel column eluted with 20% EtOAc in hexanes. Combined and concentrated pure fractions to yield title compound. (1.05 g, 55%) APCI (AP-): 511.1 (M-H)⁻.

-219-

Step 5: Pyrrolidine-2-carboxylic acid (2'-methanesulfonyl-3-trifluoromethyl-biphenyl-4-yl)-amide

2-(2'-Methanesulfonyl-3-trifluoromethyl-biphenyl-4-ylcarbamoyl)-

- 5 pyrrolidine-1-carboxylic acid *tert*-butyl ester (0.97 g, 1.9 mmol) was dissolved in 10 mL DCM and 5 mL trifluoroacetic acid and stirred at ambient temperature overnight and concentrated. Redissolved in 100 mL EtOAc and 100 mL sat. NaHCO₃, separated layers, washed organics with brine, dried with MgSO₄, filtered and concentrated to yield title compound as an oil (0.78 g, 100%) APCI
- 10 (AP-): 411.0 (M-H)⁻.

Step 6: Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3-trifluoromethyl-biphenyl-4-yl)-amide]


Pyrrolidine-2-carboxylic acid (2'-methanesulfonyl-3-trifluoromethyl-

- 15 biphenyl-4-yl)-amide (0.4 g, 0.97 mmol) was dissolved in 10 mL THF, added diisopropylethylamine (0.5 mL, 2.9 mmol), followed by 4-chlorophenylisocyanate (148 mg, 0.96 mmol). Stirred reaction at ambient temperature for one hour then concentrated. Purified on a silica gel column eluted with 20% moving to 50% EtOAc in hexanes. Combined and concentrated pure fractions to yield title
- 20 compound. (0.48 g, 88%) APCI (AP-): 564.0, 566.0 (M-H)⁻.


General Procedure for Preparation of Examples 28 *et. seq.*:

-220-


Step 1
Route 1



Route 2



Step 2



- Synthesis of 2 (Step 1):** As provided in Route 1, a mixture of 1 (where the structure of YAZ is as provided in the Scheme), B (where B is a nitrogen in a heteroaryl or heterocycloalkyl ring, an acyclic alkyl primary or secondary amine, or an acyclic primary or secondary amide), K_3PO_4 , CuI, and *trans*-diaminocyclohexane is heated in dioxane at reflux to obtain compound 2, wherein the point of attachment of YA to B is the nitrogen of the heteroaryl or heterocycloalkyl ring, or the nitrogen of the acyclic primary or secondary alkyl amine, or the nitrogen of the acyclic primary or secondary amide. In the case where Y is a nitro group, the moiety is reduced to the amino group with RaNi and EtOH under a hydrogen atmosphere.

-221-


Alternatively, as provided in Route 2, CuI, K₂CO₃, and 8-hydroxyquinoline in DMSO can be used to carry out the coupling reaction.

Synthesis of 4 (Step 2): The appropriate aniline (2= YAB), N-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline (EEDQ), triethylamine and carboxylic acid (3) are heated at reflux in chloroform to produce 4.

Synthesis of 5: A solution of compound 4, TFA and DCM is stirred at room temperature for 1 h and then concentrated under reduced pressure. The resulting oil is dissolved in THF and cooled to 0 °C followed by the addition of triethylamine and the appropriate isocyanate to produce compound 5.

Example 28

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]



Step 1: (2R,4R)-4-Hydroxy-pyrrolidine-2-carboxylic acid methyl ester.

Cis-4-Hydroxy-D-proline (15g, 115mmol) was suspended in 150mL anhydrous methanol under an argon atmosphere, then cooled to 0°C before bubbling in HCl gas for 15 minutes. The solution gradually became homogenous. The argon and HCl gas lines were removed and the solution was refluxed for 4h. The solution was cooled and then concentrated under reduced pressure. The crude material was redissolved in 100mL methanol and diethyl ether was added until a

-222-

precipitate formed. The precipitate was filtered off, washed with diethyl ether, and dried in vacuo overnight to reveal 1 (20 g, 95%) as a white solid. MS: APCI (AP+): 146 (M)+.

5 **Step 2: (2R,4R)-4-Hydroxy-1-trityl-pyrrolidine-2-carboxylic acid methyl ester (2).**

Into a solution of 1 (10g, 55mmol) in anhydrous CHCl_3 (100mL) was added triethylamine (19mL, 138mmol) and triphenylmethyl chloride (14.5g, 52mmol). The mixture was stirred at RT for 3d before concentrating and
10 redissolving in EtOAc. The solution was washed sequentially with 10% aq. citric acid, water, and brine before drying over MgSO_4 and concentrating under reduced pressure to reveal 2 (20g, 100%) as a yellow solid.

15 **Step 3: (2R,4R)-4-Methoxy-1-trityl-pyrrolidine-2-carboxylic acid methyl ester (3).**

Into a solution of 2 (5.71g, 14.7mmol) in anhydrous DMF (20mL) and anhydrous THF (20mL) was added MeI (3.67mL, 58.9mmol). The solution was cooled to 0°C in an ice bath and NaH (0.766g, 19.2mmol) was added in one portion. The mixture was stirred at RT for 26h before adding EtOAc and washing
20 sequentially with water and brine. The organic layer was dried over MgSO_4 and concentrated under reduced pressure. Purification of the crude material by flash chromatography revealed 3 (4.67g, 79%) as a white solid.

Step 4: (2R,4R)-4-Methoxy-pyrrolidine-2-carboxylic acid methyl ester (4).

25 To a flask containing 3 (4.67g, 11.6mmol) was added a solution of CH_2Cl_2 (27mL), water (0.3mL), and TFA (3.0mL, 38.9mmol). The solution was stirred at RT for 3h before concentrating under reduced pressure to reveal impure 4. MS: APCI (AP+): 160.1 (M)+.

30 **Step 5: (2R,4R)- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester 2-methyl ester (5).**

-223-

Into a solution of 4 (1.85g, 11.6mmol) in CH₂Cl₂ (20mL) was added triethylamine (6.49mL, 46.5mmol), di-*tert*-butyl dicarbonate (5.08g, 23.3mmol), and dimethylaminopyridine (0.142g, 1.16mmol). The mixture was stirred at RT for 22h before concentrating under reduced pressure and redissolving in EtOAc.

- 5 The solution was washed sequentially with 10% aq. citric acid and brine before drying over MgSO₄ and concentrating under reduced pressure. The crude material was purified by flash chromatography to reveal 5 (2.54g, 84% over two steps) as a yellow solid. MS: APCI (AP+): 260.1 (M)+.

10 **Step 6: (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (6).**

- Into a solution of 5 (2.54g, 9.80mmol) in acetonitrile (20mL) was added water (20mL) and LiOH·H₂O (1.64g, 39.2mmol). The mixture was stirred at RT for 28h before removing the acetonitrile under reduced pressure. EtOAc was
15 added to the residue that was then washed with 1N HCl. The aqueous layer was extracted with additional EtOAc and the combined organic layers were washed with brine and dried over MgSO₄. The solution was concentrated under reduced pressure to reveal 6 (2.16g, 90%) as a white solid. MS: APCI (AP-): 244.1 (M)-.

20 **Step 7: 1-(4-Amino-3-fluoro-phenyl)-piperidin-2-one (7).**

- 2-Fluoro-4-iodoaniline (10.0g, 42.2mmol) was combined with δ-valerolactam (6.27g, 63.3mmol), CuI (0.804g, 4.22mmol), and K₃PO₄ (22.4g, 105mmol). 1,4-Dioxane (60mL) was added followed by *trans*-1,2-diaminocyclohexane (1.01mL, 8.44mmol). The mixture was heated to reflux for
25 22h before cooling and diluting with EtOAc. The mixture was filtered through a plug of silica, eluting with EtOAc, and the filtrate concentrated under reduced pressure. Purification of the crude product by flash chromatography revealed 7 (3.40g, 39%) as a brown solid. MS: APCI (AP+): 209.1 (M)+.

30 **Step 8: (2R,4R)-2-[2-Fluoro-4-(2-oxo-piperidin-1-yl)-phenylcarbamoyl]-4-methoxy-pyrrolidine-1-carboxylic acid *tert*-butyl ester (8).**

-224-

Into a solution of 6 (0.250g, 1.02mmol) in CHCl_3 (10mL) was added 7 (0.212g, 1.02mmol), EEDQ (0.302g, 1.22mmol), and triethylamine (0.213mL, 1.53mmol). The solution was stirred at reflux for 19h before cooling to RT and adding EtOAc. The solution was washed sequentially with 10% aq. citric acid, 5 1N NaOH, water, and brine, before drying over MgSO_4 and concentrating under reduced pressure. The crude material was purified by flash chromatography to reveal 8 (0.329g, 74%) as a tan foam. MS: APCI (AP+): 436.1 (M)+, (AP-): 434.1 (M)-.

10 **Step 9: (2R, 4R)-4-Methoxy-pyrrolidine-2-carboxylic acid [2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide (9).**

Into a solution of 8 (0.329g, 0.761mmol) in anhydrous CH_2Cl_2 (5mL) was added TFA (5mL). The solution was stirred at RT for 0.5h before concentrating under reduced pressure to reveal 9 (0.255g, 100%) as a tan oil.

15


Step 10: (2R,4R)- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide](10).

Into a solution of 9 (0.128g, 0.380mmol) in anhydrous THF (5mL) at 0°C was added triethylamine (0.265mL, 1.90mmol) and 4-chlorophenyl isocyanate 20 (0.058g, 0.380mmol). The solution was stirred at RT for 3.5h before concentrating under reduced pressure. The crude material was purified by flash chromatography, followed by filtration through a fritted funnel and concentration under reduced pressure. Remaining EtOAc was azeotroped off with CHCl_3 . The resulting solid was lyophilized from acetonitrile / water to reveal 10 (0.186g, 25 100%) as a white solid. MS: APCI (AP+): 489.1 (M)+, (AP-): 487.1 (M)-; CHN calc'd for $\text{C}_{24}\text{H}_{26}\text{Cl}_2\text{F}_2\text{N}_4\text{O}_4$: %C 50.75; %H 4.58; %N 9.07. Found: %C 50.62; %H 4.19; %N 9.02.

-225-

Example 29

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2-oxo-piperidin-1-yl)-phenyl]-amide]




5

The compound was prepared as generally provided in Example 28. MS:
APCI (AP+): 457.1 (M)+.

10

Example 30

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]




15

The compound was prepared as generally provided in Example 28. MS:
APCI (AP+): 475.1 (M)+.

-226-

Example 31

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]




5

The compound was prepared as generally provided in Example 28. MS:
APCI (AP+): 476.1 (M)+.

10

Example 32

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]




15

The compound was prepared as generally provided in Example 28. MS:
APCI (AP+): 490.1 (M)+.

-227-

Example 33

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]




5

The compound was prepared as generally provided in Example 28. MS: APCI (AP+): 504.1 (M)+.

10

Example 34

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]




15

The compound was prepared as generally provided in Example 28. MS: APCI (AP+): 503.1 (M)+.

-228-

Example 35

(2R,4R)-4-Propoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]




5

The compound was prepared as generally provided in Example 28. MS:
APCI (AP+): 517.2 (M)+.

10

Example 36

(2R,4R)-4-Propoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]




15

The compound was prepared as generally provided in Example 28. MS:
APCI (AP+): 518.2 (M)+.

-229-

Example 37

(2R,4R)-4-Cyano-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]




5

The compound was prepared as generally provided in Example 28. MS: APCI (AP+): 484.2 (M)+.

10

Example 38

(2R,4R)-4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]




15

The compound was prepared as generally provided in Example 28. MS: APCI (AP+): 477.1 (M)+.

-230-

Example 39

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-azetidin-1-yl)-phenyl]-amide]




5

The compound was prepared as generally provided in Example 28. MS:
APCI (AP+): 461.1 (M)+.

10

Example 40

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-azetidin-1-yl)-phenyl]-amide]




15

The compound was prepared as generally provided in Example 28. MS:
APCI (AP+): 462.1 (M)+.

-231-

Example 41

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-pyrrolidin-1-yl)-phenyl]-amide]




5

The compound was prepared as generally provided in Example 28. MS:
APCI (AP+): 475.2 (M)+.

10

Example 42

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-pyrrolidin-1-yl)-phenyl]-amide]




15

The compound was prepared as generally provided in Example 28. MS:
APCI (AP+): 476.2 (M)+.

-232-

Example 43

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-azepan-1-yl)-phenyl]-amide]




5

The compound was prepared as generally provided in Example 28. MS:
APCI (AP+): 503.2 (M)+.

10

Example 44

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-azepan-1-yl)-phenyl]-amide]




15

The compound was prepared as generally provided in Example 28. MS:
APCI (AP+): 504.2 (M)+.

-233-

Example 45

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide]




5

The compound was prepared as generally provided in Example 28. MS:
APCI (AP+): 468.2 (M)+.

10

Example 46

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide]




15

The compound was prepared as generally provided in Example 28. MS:
APCI (AP+): 467.2 (M)+.

-234-

Example 47

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide]




5

The compound was prepared as generally provided in Example 28. MS:
APCI (AP+): 499.2 (M)+.

10

Example 48

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide]



15


The compound was prepared as generally provided in Example 28. MS:
APCI (AP+): 500.2 (M)+.

-235-

Example 49

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-{{2-fluoro-4-(2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-phenyl}-
amide}

5




The compound was prepared as generally provided in Example 28. MS:
APCI (AP+): 505.2 (M)+.

10

Example 50

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-{{4-(2,3-dimethyl-5-oxo-2,5-dihydro-pyrazol-1-yl)-phenyl}-amide}



15


The compound was prepared as generally provided in Example 28. MS:
APCI (AP+): 470.2 (M)+.

20

-236-

Example 51

**4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-
{[4-(2-hydroxymethyl-pyrrolidin-1-yl)-phenyl]-amide}**




5

The compound was prepared as generally provided in Example 28. APCI
(AP+): 459.2 (M+H)+.

10

Example 52

**(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[(4-pyrrolidin-1-yl-phenyl)-amide]**




15

The compound was prepared as generally provided in Example 28. APCI
(AP+): 429.1 (M)+.

-237-

Example 53

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-pyrazol-1-yl-phenyl)-amide]




5

The compound was prepared as generally provided in Example 28. APCI (AP+): 426.1 (M)+.

10

Example 54

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-pyrazol-1-yl-phenyl)-amide]



15


The compound was prepared as generally provided in Example 28. APCI (AP+): 444.1 (M)+.

Example 55

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-[1,2,4]triazol-1-yl-phenyl)-amide]

20

-238-




The compound was prepared as generally provided in Example 28. APCI (AP+): 427.1 (M)+.

5

Example 56

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-[1,2,3]triazol-2-yl-phenyl)-amide]



10


The compound was prepared as generally provided in Example 28. APCI (AP+): 427.1 (M)+.

15

Example 57

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-[1,2,3]triazol-1-yl-phenyl)-amide]

-239-




The compound was prepared as generally provided in Example 28. APCI (AP+): 427.1 (M)+.

5

Example 58

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(4-acetylaminophenyl)-amide] 1-[(4-chloro-phenyl)-amide]



10


The compound was prepared as generally provided in Example 28. APCI (AP+): 417.1 (M)+.

15

Example 59

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(cyclopentanecarbonyl-amino)-phenyl]-amide]

-240-




The compound was prepared as generally provided in Example 28. APCI (AP+): 471.2 (M)+.

5

Example 60

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-pyrimidin-5-yl-phenyl)-amide]



10


The compound was prepared as generally provided in Example 28. APCI (AP+): 484.2 (M)+.

15

Example 61

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-pyrazol-1-yl-phenyl)-amide]

-241-




The compound was prepared as generally provided in Example 28. APCI (AP+): 440.2 (M)+.

5

Example 62

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-pyrazol-1-yl-phenyl)-amide]



10


The compound was prepared as generally provided in Example 28. APCI (AP+): 458.1 (M)+.

15

Example 63

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-pyrazol-1-yl-phenyl)-amide]

-242-




The compound was prepared as generally provided in Example 28. APCI (AP+): 472.1 (M)+.

5

Example 64

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-pyrazol-1-yl-phenyl)-amide]



10


The compound was prepared as generally provided in Example 28. APCI (AP+): 454.2 (M)+.

Example 65

15

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-methyl-pyrazol-1-yl)-phenyl]-amide]

-243-



Step 1: 2-Fluoro-4-(3-methyl-pyrazol-1-yl)phenylamine (1)

2-Fluoro-4-iodoaniline (3.0 g, 12.6 mmol), 3-methylpyrazole (1.22 mL, 15.18 mmol), Cs_2CO_3 (8.66 g, 26.58 mmol) and CuI (0.072 g, 0.379 mmol) were placed in a flask under nitrogen. The flask was evacuated under vacuum and refilled with nitrogen (5 times). 1,4-Dioxane (15 mL) was added followed by *trans*-1,2-diaminocyclohexane (152 μL , 1.266 mmol). The reaction mixture was heated under reflux for 36 h. Mixture cooled and filtered. Solvents removed.

Residue purified by silica gel flash chromatography eluting with a gradient of EtOAc in hexanes (0 to 30%). This purification afforded **1** as a pale brown solid (0.917 g, 38%). MS: APCI (AP+): 192.0 (M)+.

From the same purification, 2-Fluoro-4-(5-methyl-pyrazol-1-yl)phenylamine (**2**) was obtained as pale brown solid (0.265 g, 11%). MS: APCI (AP+): 192.0 (M)+.

Step 2: (2R,4R)-2-[2-Fluoro-4-(3-methyl-pyrazol-1-yl)-phenylcarbamoyl]-4-methoxy-pyrrolidine-1-carboxylic acid *tert*-butyl ester (3).

(2R,4R)-4-methoxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (0.350 g, 1.427 mmol), aniline **1** (0.273 g, 1.427 mmol) and EEDQ (0.529 g, 2.140 mmol) were dissolved in CHCl_3 (30 mL). Et_3N (0.6 mL, 4.281 mmol) was added and the reaction mixture was heated at reflux for 18 h. The mixture was allowed to cool and diluted with CHCl_3 (100 mL) and washed with 5% HCl, 0.5 N NaOH, water and brine (40 mL each), the organic extract was dried over MgSO_4 , filtered and the solvent removed to give a pale tan foam. The crude product was

-244-

used for the next step without purification. MS: APCI (AP+): 419.2 (M)+, (AP-): 417.2 (M)-.

Step 3: (2R, 4R)-4-Methoxy-pyrrolidine-2-carboxylic acid [2-fluoro-4-(3-methyl-pyrazol-1-yl)-phenyl]-amide (4)

Boc-amine 3 (~0.6 g, 1.427 mmol) was dissolved in 25% TFA in dichloromethane. The mixture was stirred at ambient temperature for 2 h. The solvents were removed and the residue dried under vacuum to give 4 as a tan oil (~0.450 g).

10

Step 4: (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-methyl-pyrazol-1-yl)-phenyl]-amide] (5)


A solution of amine 4 (~0.450 g, 1.427 mmol) in dichloromethane (35 mL) was cooled at 0°C. Triethylamine (1.5 mL, 10.7 mmol) was added and the mixture stirred for 15 min. Then, 4-chlorophenyl isocyanate (0.24 g, 1.569 mmol) was incorporated. The solution was stirred at ambient temperature for 18 h. The mixture was diluted with CHCl₃ (100 mL) and washed with 5% HCl, water, brine (40 mL each). The organic extracts were dried over MgSO₄, filtered and the solvent removed. The residue was purified by flash chromatography on silica gel eluting with a gradient of EtOAc in hexanes (0 to 65%) to provide 5 as white solid (0.56 g, 83% in 3 steps). MS: APCI (AP+): 472.2 (M)+, (AP-): 470.2 (M)-. Anal. calcd for C₂₃H₂₃ClFN₅O₃: C, 58.54; H, 4.91; N, 14.84. Found: C, 58.35; H, 4.59; N, 14.49.

25

Example 66

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-methyl-pyrazol-1-yl)-phenyl]-amide]

-245-




The compound was prepared as generally provided for Example 65. APCI (AP+): 458.2 (M)+.

5

Example 67

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-methyl-pyrazol-1-yl)-phenyl]-amide]



10


The compound was prepared as generally provided for Example 65. APCI (AP+): 472.2 (M)+.

15

Example 68

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(5-methyl-pyrazol-1-yl)-phenyl]-amide]

-246-




The compound was prepared as generally provided for Example 65. APCI (AP+): 454.3 (M)+.

5

Example 69

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3,5-dimethyl-pyrazol-1-yl)-phenyl]-amide]



10


The compound was prepared as generally provided for Example 65. APCI (AP+): 454.2 (M)+.

15

Example 70

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3,5-dimethyl-pyrazol-1-yl)-phenyl]-amide]

-247-




The compound was prepared as generally provided for Example 65. APCI (AP+): 468.2 (M)+.

5

Example 71

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3,5-dimethyl-pyrazol-1-yl)-2-fluoro-phenyl]-amide]




10

The compound was prepared as generally provided for Example 65. APCI (AP+): 486.2 (M)+.

15

Example 72

1-(4-Amino-3-fluoro-phenyl)-1H-pyridin-2-one



According to Route 2 for the preparation of YAB as provided in the
20 General Procedure for the Preparation of Examples 28-60, 2-Fluoro-4-iodoaniline

-248-


(10.0g, 42.2mmol) was combined with 1*H*-pyridin-2-one (1.0g, 4.22mmol), CuI (0.120g, 0.633mmol), K₂CO₃ (0.641g, 4.64mmol), and 8-hydroxyquinoline (0.092g, 0.633mmol) in DMSO (3mL). The mixture was degassed with a stream of argon and then heated at reflux for 20h before cooling to ambient temperature.

- 5 10% aq. NH₄OH and EtOAc were added, and the mixture was filtered through a plug of layered celite and decolorizing charcoal, eluting with EtOAc. The filtrate was concentrated under reduced pressure. Purification of the crude product by mpic revealed the title compound (0.53g, 62%) as a yellow solid. MS: APCI (AP+): 205.1 (M)+.

10

Example 73

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(4-tert-butyl-phenyl)-amide] 1-[(4-chloro-phenyl)-amide]




15

The compound was prepared as generally provided for Example 28. MS: APCI (AP+): 414.2 (M)+.

Example 74

- 20 (2*R*,4*R*)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2[(3,5'-difluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]

-249-

**Step 1: *N*-tert-Butyl-4-fluoro-benzenesulfonamide**

Into a DCM solution (50 mL) of *tert*-butylamine (5.9 mL, 56.5 mmol) and triethylamine (7.16 mL, 51.3 mmol) at 0°C was slowly added a DCM solution (20 mL) of 4-fluoro-benzenesulfonyl chloride (10 g, 51.3 mmol), and the reaction stirred at RT for 19h. The reaction was concentrated and the resulting solid recrystallized from ether/hexanes to get the product (9.5 g, 80%) as white crystals. MS: APCI (AP+): 230.1 (M)+.

10

Step 2: *N*-tert-Butyl-4-fluoro-benzenesulfonamide-3-boronic acid

Butyllithium (2.2 M, 4.3 mL, 9.51 mmol) was added drop-wise to a 0°C solution of *N*-tert-butyl-4-fluoro-benzenesulfonamide (1.0 g, 4.32 mmol), and the solution stirred for 15 min. at 0°C and then RT for 1.5 h. Triisopropyl borate (1.19 mL, 5.18 mmol) was added, and the solution stirred at RT of 2.5 h. The mixture was cooled to 0°C, 10% aq. HCl was added slowly, and the reaction was stirred for 1.5h. The reaction was then extracted with EtOAc three times and the organics combined and concentrated. The resulting oil was dissolved in ether and extracted with 1N NaOH three times, and the extracts combined and acidified at 0°C with 6N HCl. The solution was extracted with ether three times and the organics were combined and dried over MgSO₄. Filtration and concentration of the organics revealed the product as an oil which formed a solid upon addition of hexanes and concentration under reduced pressure. (0.641g, 54%) APCI (AP-): 274.1 (M)-.

25

-250-

Step 3: (2R, 4R)-4-(tert-Butyl-dimethyl-silanyloxy)-2-(2-fluoro-4-iodo-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid tert-butyl ester.

4-(tert-Butyl-dimethyl-silanyloxy)-pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl ester (0.673 g, 1.94 mmol), 2-fluoro-4-iodoaniline (0.46 g, 1.94 mmol),
5 EEDQ (0.575 g, 2.32 mmol) and triethylamine (0.408 mL, 2.91 mmol) were heated at reflux in CHCl₃ (10 mL) for 22 h. The solution was concentrated and resulting oil partitioned between EtOAc and water. The organic layer was washed with brine and dried over MgSO₄. The product was purified by mplc to produce the product as a clear oil (1.0g, 90%). MS: APCI (AP-): 563.2 (M)-.

10

Step 4: (2R,4R)-4-(tert-Butyl-dimethyl-silanyloxy)-2-(2'-tert-butylsulfamoyl-3,5'-difluoro-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid tert-butyl ester

A mixture of 4-(tert-butyl-dimethyl-silanyloxy)-2-(2-fluoro-4-iodo-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid tert-butyl ester (0.56g, 0.99 mol),
15 Pd(PPh₃)₄ (0.11 g, 0.1 mmol), *N*-tert-butyl-4-fluoro-benzenesulfonamide-3-boronic acid (0.33 g, 1.19 mmol), tetrabutylammonium bromide (0.016 g, 0.05 mmol), sodium carbonate (0.210 g, 1.98 mmol), water (2.0 mL) and toluene (10 mL) was degassed with a stream of argon gas and then heated at reflux 21.5 h.
20 The mixture was then allowed to cool to RT, concentrated and partitioned between water and EtOAc. The organic layer was washed with brine, dried over MgSO₄ and concentrated. The resulting oil was purified by mplc to reveal the product as an oil (0.119g, 18%). MS: APCI (AP-): 666.4 (M)-.

Step 5: (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(2'-tert-butylsulfamoyl-3,5'-difluoro-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide]

A solution of 4-(tert-butyl-dimethyl-silanyloxy)-2-(2'-tert-butylsulfamoyl-3,5'-difluoro-biphenyl-4-ylcarbamoyl)-pyrrolidine-1-carboxylic acid tert-butyl
30 ester (0.228g, 0.342 mmol) and trifluoroacetic acid (1.5 mL) in CHCl₃ (1.5 mL) was stirred at ambient temperature for 1h. The solution was concentrated, hexanes added and then the solution concentrated again. The resulting oil was

-251-


dissolved in THF and cooled 0°C. Triethylamine (0.190 mL, 1.36 mmol) was added followed by 4-chlorophenyl isocyanate (0.052 g, 0.342 mmol) and the solution stirred at RT for 5.5 h. The solution was concentrated and the oil purified by mpic to reveal the product as a solid (0.078g, 38%). MS: APCI (AP-): 605.2 (M)-.

Step 6: (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3,5'-difluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(2'-*tert*-butylsulfamoyl-3,5'-difluoro-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide] (0.078g, 0.128 mmol) was dissolved in TFA (3 mL) and stirred at RT for 1 h. The reaction was concentrated and dissolved in MeCN and water and purified by reverse phase HPLC to reveal the product as a white solid (0.05g, 77%) after lyophilization. MS: APCI (AP+): 551.1 (M)+.


Example 75

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)-amide]



The compound was prepared as generally provided for Example 28. MS: APCI (AP+): 472.2 (M)+.

-252-

Example 75a. Alternative Procedure.**Synthesis of 2:**

- 5 A solution of 1 where Y is a nitro group is treated with a halo-alkylchloride in the presence of base and refluxing toluene to produce an intermediate halo-amide that is then cyclized in the presence of potassium *tert*-butoxide in THF to produce a cyclic lactam B.

10 Synthesis of 3:

A solution of 2 where Y is a nitro group is reduced under RaNi mediated hydrogenation to produce 3 where Y is an amino group.


Synthesis of 5 and 6:

- 15 Intermediates 5 and 6 were prepared as provided in Example 28.

Example 76

- 20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide]

-253-



Step 1: 5'-Nitro-3,4,5,6-tetrahydro-[1,2']bipyridinyl-2-one.

See Example 75a. 2-Amino-5-nitro-pyridine (2.0 g, 14.3 mmol), 5-bromo-pentanoyl chloride (2.86 g, 14.3 mmol), triethylamine (2.39 g, 17.2 mmol), catalytic DMAP were heated at reflux in toluene (70 mL) 22 h. The reaction was allowed to cool to RT, concentrated and the resulting oil portioned between EtOAc and water. The organic layer was separated, washed with brine, dried over MgSO₄ and filtered. Concentration of the filtrate produced an oil which was dissolved in THF (140 mL). Potassium tert-Butoxide (1.68g, 15.0 mmol) was added portion-wise and the reaction stirred at RT for 2h. Water was then added, and the reaction extracted with EtOAc. The organics were then washed with brine and dried over MgSO₄. Purification of the crude material by mpc revealed the product as an orange solid. MS: APCI (AP+): 222.1 (M)+.

Step 2: 5'-Amino-3,4,5,6-tetrahydro-[1,2']bipyridinyl-2-one.

5'-Nitro-3,4,5,6-tetrahydro-[1,2']bipyridinyl-2-one (0.66 g, 2.98 mmol) was dissolved in 1:1 MeOH/THF (50 mL) and hydrogenated over RaNi (1g) at 4285 psi/mole for 0.8 h. The catalyst was filtered, and the filtrate concentrated. The crude solid was purified by recrystallization from THF and hexanes to give the product as a yellow solid (0.275g, 48%). MS: APCI (AP+): 192.1 (M)+.

Step 3: (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chlorophenyl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide].


-254-

The compound was prepared as generally provided for Example 28. MS:
APCI (AP+): 472.2 (M)+.

Example 77

5

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(2,3-dimethyl-5-oxo-2,5-dihydro-pyrazol-1-yl)-phenyl]-amide]




10

The compound was prepared as generally provided for Example 28.
Formation of the 5-chloro-pyridin-2-yl moiety can be found in Example 26. MS:
APCI (AP+): 485.3 (M)+.

15

Example 78

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,3-dimethyl-5-oxo-2,5-dihydro-pyrazol-1-yl)-phenyl]-amide]



20


-255-

The compound was prepared as generally provided for Example 28. MS:
APCI (AP+): 484.3 (M)+.

Example 79

5

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[(2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide]



10 **Step 1. 6'-Amino-[1,3']bipyridinyl-2-one.**

2-Amino-5-iodopyridine (5.0g, 22.7mmol) was combined with 1H-pyridin-2-one (2.59g, 27.27 mmol), CuI (0.65g, 3.41 mmol), K₂CO₃ (3.45 g, 24.9mmol), and 8-hydroxyquinoline (0.49 g, 3.40 mmol) in DMSO (20 mL). The mixture was degassed with a stream of argon and then heated at reflux for 18h
15 before cooling to RT. 10% aq. NH₄OH and EtOAc were added, and the mixture was filtered through a plug of layered celite and decolorizing charcoal, eluting with EtOAc. The filtrate was concentrated under reduced pressure. Purification of the crude product by mpic revealed A (0.40g, 10%) as a yellow solid. MS:
APCI (AP+): 188.1 (M)+.

20

Step 2: (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide].

The compound was prepared as generally provided for Example 28. MS:
APCI (AP+): 468.3 (M)+.


25

-256-

Example 80

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide]-

TFA Salt




The compound was prepared as generally provided for in Examples 28 and 75a. Formation of the 5-chloro-pyridin-2-yl moiety can be found in Example 26.

10 Purification was accomplished via reverse phase HPLC. MS: APCI (AP+): 473.2 (M)+.

Example 81

15 4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide]




The compound was prepared as generally provided for Example 28. MS:
20 APCI (AP+): 486.3 (M)+.

-257-

Example 82


(2R,4R)-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[(2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide]



The compound was prepared as generally provided for Examples 28 and 79. MS: APCI (AP+): 482.3 (M)+.

Example 83

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[[2-fluoro-4-(6-oxo-6H-pyridazin-1-yl)-phenyl]-amide]



Step 1: 2-(4-Amino-3-fluoro-phenyl)-2H-pyridazin-3-one.

2-Fluoro-4-iodoaniline (0.5 g, 2.10 mmol) was combined with 2H-Pyridazin-3-one (0.243 g, 2.53 mmol), CuI (0.06 g, 0.315 mmol), K₂CO₃ (0.32 g, 2.31 mmol), and 8-hydroxyquinoline (0.046 g, 0.315 mmol) in DMSO (3mL).

-258-


The mixture was degassed with a stream of argon and then heated at reflux for 23 h before cooling to RT. 10% aq. NH_4OH and EtOAc were added, and the mixture was filtered through a plug of layered celite and decolorizing charcoal, eluting with EtOAc. The filtrate was concentrated under reduced pressure. Purification of the crude product by mpls revealed A (0.16 g, 37%) as a yellow solid. MS: APCI (AP+): 206.1 (M)+.

Step 2: (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(6-oxo-6H-pyridazin-1-yl)-phenyl]-amide].

The compound was prepared as generally provided for Example 28. MS: APCI (AP+): 486.1 (M)+.

Example 84

(2R,4S)-4-Amino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]



Step 1: (2R,4R)-toluene-4-sulfonic acid 1-(4-chloro-phenylcarbamoyl)-5-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidin-3-yl ester.

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] (see example 17, 0.41 g, 0.77 mmol) was dissolved in 5ml dry DCM, added Et3N (0.21 mL, 1.54 mmol), DMAP (10 mg, 0.8 mmol), and TsCl (0.16 g, 0.85 mmol). Stirred at ambient temperature overnight. The solution was dissolved in 100ml EtOAc, washed with 10% citric acid (3x50ml), sat NaHCO_3 (3x50ml), brine

-259-

(100ml), dried with MgSO₄, filtered and concentrated to yield title compound. (0.54 g, 97%) MS: APCI (AP-): 684.1 (M)-.

Step 2: (2R,4S)-4-Azido-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide].

(2R,4R)-toluene-4-sulfonic acid 1-(4-chloro-phenylcarbamoyl)-5-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidin-3-yl ester (0.2 g, 0.29 mmol) was dissolved in 4ml dry DMF under Ar, added sodium azide (0.38 mg, 0.58 mmol) and stirred at 60°C for 6 hours. The reaction was cooled, taken up in 100ml EtOAc, washed with water, sat. NaHCO₃, brine, dried with MgSO₄, filtered and concentrated to yield title compound. (0.155 g, 95%) MS: APCI (AP+): 557.0 (M)+.


Step 3: (2R,4S)-4-Amino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide].

(2R,4S)-4-Azido-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] (0.145 g, 0.26 mmol) was shaken with Raney Nickel (300mg) in 16 mL tetrahydrofuran in a hydrogen atmosphere under pressure for 3 days. Reaction was filtered and concentrated. Purified by HPLC and lyophilized pure fractions to yield title compound. (105 mg, 76%) MS: APCI (AP-): 529.1 (M)-.

Example 85

(2R,4R)-4-Amino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]

-260-




The compound was prepared as generally provided for Example 85 substituting (2R,4S)-toluene-4-sulfonic acid 1-(4-chloro-phenylcarbamoyl)-5-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidin-3-yl ester for (2R,4R)-toluene-4-sulfonic acid 1-(4-chloro-phenylcarbamoyl)-5-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidin-3-yl ester. MS: APCI (AP+): 531.1 (M)+.

10

Example 86

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]



15


The compound was prepared as generally provided for Example 17, substituting 2-(N-*tert*-butyl)phenylsulfonamide boronic acid for 2-(methylthio)benzene boronic acid. MS: APCI (AP+): 533.0 (M)+.

20

-261-

Example 87

(2R)-4-Hydroxyimino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]




5

The compound was prepared as generally provided for Example 22. MS: APCI (AP+): 545.0 (M)+.

10

Example 88

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methylsulfamoyl-biphenyl-4-yl)-amide]




15

The compound was prepared as generally provided for Example 86. MS: APCI (AP+): 547.2 (M)+.

-262-

Example 89

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-dimethylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide]




5

The compound was prepared as generally provided for Example 86. MS: APCI (AP+): 561.2 (M)+.

10

Example 90

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]



15


The compound was prepared as generally provided for Example 28, substituting 3-fluoro-2'-methanesulfonyl-biphenyl-4-ylamine for 1-(4-amino-3-fluoro-phenyl)-piperidin-2-one. MS: APCI (AP+): 547.1 (M)+.

20

-263-

Example 91

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]




5

The compound was prepared as generally provided for Example 28, substituting 4'-amino-3'-fluoro-biphenyl-2-sulfonic acid tert-butylamide for 1-(4-amino-3-fluoro-phenyl)-piperidin-2-one. MS: APCI (AP+): 549.0 (M)+.

10

Example 92

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]



15


The compound was prepared as generally provided for Example 90, substituting (2R,4R)-4-ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester for (2R,4R)-4-methoxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester. MS:

20 APCI (AP+): 561.1 (M)+.

-264-

Example 93


5 (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]



The compound was prepared as generally provided for Example 28, substituting (2R,4R)-4-ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester for (2R,4R)-4-methoxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester. MS:
10 APCI (AP+): 560.2 (M)+.

Example 94

15 (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]



The compound was prepared as generally provided for Example 91,
20 substituting (2R,4R)-4-ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester


-265-

for (2R,4R)-4-methoxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester. MS:
APCI (AP+): 561.1 (M)+.

Example 95

5

(2R,4S)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]




10

The compound was prepared as generally provided for Example 28,
substituting (2R,4S)-4-methoxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl
ester for (2R,4R)-4-methoxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester.
MS: APCI (AP+): 546.0 (M)+.

15

Example 96

(2R,4R)-4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]



20


-266-

The compound was prepared as generally provided for Example 28, substituting (2R,4R)-4-ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester for (2R,4R)-4-methoxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester MS: APCI (AP+): 534.0 (M)+.

5

Example 97

(2R)-4,4-Difluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]



10

The compound was prepared as generally provided for Example 28, substituting (2R)-4-difluoro-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester for (2R,4R)-4-methoxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester. MS: APCI (AP+): 552.0 (M)+.


15

Example 98

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(4-fluoro-phenyl)-amide]

20


-267-



The compound was prepared as generally provided for Example 28,
substituting 4-fluorophenyl isocyanate for 4-chlorophenyl isocyanate. MS: APCI
5 (AP+): 530.2 (M)+.

Example 99

(2R,4R)-4-Acetylamino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
10 phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]




The compound was prepared as generally provided for Example 84. MS:
APCI (AP+): 573.2 (M)+.

15

-268-

Example 100

(2R,4R)-4-Methanesulfonylamino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]




5

The compound was prepared as generally provided for Example 84. MS: APCI (AP+): 609.2 (M)+.

10

Example 101

(2R,4S)-4-(1H-Tetrazol-5-yl)-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]




15

The compound was prepared as generally provided for Example 28. MS: APCI (AP+): 527.2 (M)+.

-269-

Example 102

(2R,4S)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]




The compound was prepared as generally provided for Example 28, substituting (2R,4S)-4-methoxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester for (2R,4R)-4-methoxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester.

10 MS: APCI (AP+): 489.2 (M)+.

Example 103

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-methyl-imidazol-1-yl)-phenyl]-amide]




The compound was prepared as generally provided for Example 28, substituting 2-Fluoro-4-(2-methyl-imidazol-1-yl)-phenylamine for 1-(4-amino-3-

20 fluoro-phenyl)-piperidin-2-one. MS: APCI (AP+): 472.2 (M)+.

-270-

Example 104

(2R,4R)-4-Cyano-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]



5


The compound was prepared as generally provided for Example 28 substituting substituting (2R,4R)-4-cyano-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester for (2R,4R)-4-methoxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester. MS: APCI (AP+): 484.2 (M)+.

10

Example 105

(2R,4R)-4-(1H-Tetrazol-5-yl)-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]

15




The compound was prepared as generally provided for Example 104. MS: APCI (AP-): 525.3 (M)-.

20

-271-

Example 106

(2R,4R)-4-Trifluoromethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]



5


The compound was prepared as generally provided for Example 28, (2R,4R)-4-trifluoromethyl-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester for (2R,4R)-4-methoxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester.. MS:

10 APCI (AP+): 527.2 (M)+.

Example 107

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-cyano-3-fluoro-biphenyl-4-yl)-amide]

15




The compound was prepared as generally provided for Example 1, substituting 2-cyanobenzene boronic acid for 2-(methylthio)benzene boronic acid.

20 MS: APCI (AP+): 493.3 (M)+.

-272-

Example 108

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[(2'-aminomethyl-3-fluoro-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide]




The compound was prepared as generally provided for Example 107. MS:
APCI (AP+): 497.3 (M)+.

10

Example 109

(2R,4R)-4'--{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3'-fluoro-biphenyl-2-carboxylic acid methyl ester




15

The compound was prepared as generally provided for Example
107. MS: APCI (AP+): 526.2 (M)+.

20 General Procedure for Example 110

-273-



5 **Synthesis of 3:** The carboxylic acid (1), aniline or amine (2), EEDQ, and triethylamine are heated at reflux in chloroform to produce 3.

Synthesis of 4: A solution of compound 3, TFA, and DCM is stirred at room temperature for 1 h and then concentrated under reduced pressure. The resulting
10 oil is dissolved in THF and cooled to 0 °C followed by the addition of triethylamine or *N,N*-diisopropylethylamine (DIEA) and the appropriate isocyanate to produce compound 4.

Synthesis of 5: To solution of compound 4 in THF is added potassium
15 trimethylsilanolate and is stirred at room temperature overnight to produce 5.


Synthesis of 7: To solution of compound 5 in DMF is added *N,N*-diisopropylethylamine, PyBOP, and NHG₁G₂ and is stirred at room temperature overnight to produce 7.

20

Example 110

-274-

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2- [[4-(2,5-dihydro-pyrrole-1-carbonyl)-phenyl]-amide]



5 **Step 1: (2R,4R)-4-Hydroxy-pyrrolidine-2-carboxylic acid methyl ester (1).**

Cis-4-Hydroxy-D-proline (15 g, 115 mmol) was suspended in 150 mL anhydrous methanol under an argon atmosphere, then cooled to 0°C before bubbling in HCl gas for 15 minutes. The solution gradually became homogenous. The argon and HCl gas lines were removed and the solution was refluxed for 4h.

10 The solution was cooled and then concentrated under reduced pressure. The crude material was redissolved in 100 mL methanol and diethyl ether was added until a precipitate formed. The precipitate was filtered off, washed with diethyl ether, and dried in vacuo overnight to give **1** (20 g, 95%) as a white solid. MS: APCI (AP+): 146 (M)+.

15

Step 2: (2R,4R)-4-Hydroxy-1-trityl-pyrrolidine-2-carboxylic acid methyl ester (2).

Into a solution of **1** (10 g, 55 mmol) in anhydrous CHCl₃ (100 mL) was added triethylamine (19 mL, 138 mmol) and triphenylmethyl chloride (14.5 g, 52 mmol). The mixture was stirred at RT for 3d before concentrating and redissolving in EtOAc. The solution was washed sequentially with 10% aq. citric acid, water, and brine before drying over MgSO₄ and concentrating under reduced pressure to give **2** (20 g, 100%) as a yellow solid.

25 **Step 3: (2R,4R)-4-Methoxy-1-trityl-pyrrolidine-2-carboxylic acid methyl ester (3).**

-275-

Into a solution of **2** (5.71 g, 14.7 mmol) in anhydrous DMF (20 mL) and anhydrous THF (20 mL) was added MeI (3.67 mL, 58.9 mmol). The solution was cooled to 0°C in an ice bath and NaH (0.766 g, 19.2 mmol) was added in one portion. The mixture was stirred at RT for 26h before adding EtOAc and washing sequentially with water and brine. The organic layer was dried over MgSO₄ and concentrated under reduced pressure. Purification of the crude material by flash chromatography revealed **3** (4.67 g, 79%) as a white solid.

Step 4: (2R,4R)-4-Methoxy-pyrrolidine-2-carboxylic acid methyl ester (4).

To a flask containing **3** (4.67 g, 11.6 mmol) was added a solution of CH₂Cl₂ (27 mL), water (0.3 mL), and TFA (3.0 mL, 38.9 mmol). The solution was stirred at RT for 3h before concentrating under reduced pressure to reveal impure **4**. MS: APCI (AP+): 160.1 (M)+.

Step 5: (2R,4R)- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl ester 2-methyl ester (5).

Into a solution of **4** (1.85 g, 11.6 mmol) in CH₂Cl₂ (20 mL) was added triethylamine (6.49 mL, 46.5 mmol), di-tert-butyl dicarbonate (5.08 g, 23.3 mmol), and dimethylaminopyridine (0.142 g, 1.16 mmol). The mixture was stirred at RT for 22h before concentrating under reduced pressure and redissolving in EtOAc. The solution was washed sequentially with 10% aq. citric acid and brine before drying over MgSO₄ and concentrating under reduced pressure. The crude material was purified by flash chromatography to reveal **5** (2.54 g, 84% over two steps) as a yellow solid. MS: APCI (AP+): 260.1 (M)+.

Step 6: (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl ester (6).

Into a solution of **5** (2.54 g, 9.80 mmol) in acetonitrile (20 mL) was added water (20 mL) and LiOH·H₂O (1.64 g, 39.2 mmol). The mixture was stirred at RT for 28h before removing the acetonitrile under reduced pressure. EtOAc was added to the residue that was then washed with 1N HCl. The aqueous layer was extracted with additional EtOAc and the combined organic layers were washed

-276-

with brine and dried over MgSO_4 . The solution was concentrated under reduced pressure to reveal 6 (2.16 g, 90%) as a white solid. MS: APCI (AP-): 244.1 (M)-.

Step 7: (2R,4R)-4-Methoxy-2-(4-methoxycarbonyl-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (7).

Into a solution of 6 (8.11 g, 33.0 mmol) in CHCl_3 (200 mL) was added 4-amino-benzoic acid methyl ester (5.00 g, 33.0 mmol), EEDQ (9.81 g, 39.6 mmol), and triethylamine (6.9 mL, 49.6 mmol). The solution was stirred at reflux for 19h before cooling to RT and adding EtOAc. The solution was washed sequentially with 1N HCl, 0.1N NaOH, water, and brine, before drying over MgSO_4 and concentrating under reduced pressure to give 7 (12.50 g, 100%). MS: APCI (AP+): 379.3 (M)+, (AP-): 377.3 (M)-.

Step 8: (2R,4R)-4-[(4-Methoxy-pyrrolidine-2-carbonyl)-amino]-benzoic acid methyl ester (8).

Into a solution of 7 (12.5 g, 33.0 mmol) in anhydrous CH_2Cl_2 (70 mL) was added TFA (20 mL). The solution was stirred at RT overnight before concentrating under reduced pressure to give 8 (9.19 g, 100%).

Step 9: (2R,4R)-4-[[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-benzoic acid methyl ester (9).

Into a solution of 8 (9.19 g, 33.0 mmol) in anhydrous THF (200 mL) at 0°C was added *N,N*-diisopropylethylamine (DIEA) (28.7 mL, 165 mmol) and 4-chlorophenyl isocyanate (5.07 g, 33.0 mmol). The solution was stirred at RT overnight before concentrating under reduced pressure. The crude material was purified on a silica gel column eluted with 10% moving to 50% EtOAc in hexanes. Combined and concentrated pure fractions. Remaining EtOAc was azeotroped off with acetonitrile and was lyophilized from acetonitrile / water to give 9 (13.37 g, 94%) as a solid. MS: APCI (AP+): 432.1 (M)+, (AP-): 430.1 (M)-.

-277-

Step 10: (2R,4R)-4-[[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-benzoic acid (10).


Into a solution of **9** (3.0 g, 6.94 mmol) in THF (50 mL) was added potassium trimethylsilanolate (3.85 g, 27.3 mmol). The solution was stirred at RT
5 for 48h before concentrating under reduced pressure and adding EtOAc. The solution was washed sequentially with 1N HCl, and brine, before drying over MgSO₄ and concentrating under reduced pressure to give **10** (2.9 g, 100%). MS: APCI (AP+): 418.1 (M)+, (AP-): 416.1 (M)-.

10 Step 11: (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,5-dihydro-pyrrole-1-carbonyl)-phenyl]-amide] (11).

Into a solution of **10** (0.25 g, 0.589 mmol) in DMF (3 mL) was added *N,N*-diisopropylethylamine (DIEA) (0.4 mL, 2.32 mmol), PyBOP (0.34 g, 0.653 mmol), and 3-pyrroline (0.09 mL, 1.19 mmol). The solution was stirred at RT for
15 19h and EtOAc added. The solution was washed sequentially with 1N HCl, saturated NaHCO₃, water, and brine, before drying over MgSO₄ and concentrating under reduced pressure. The crude material was purified on a silica gel column eluted with 20% EtOAc in hexanes moving to 1% MeOH in EtOAc. Combined and concentrated pure fractions. Remaining EtOAc was azeotroped off with
20 acetonitrile and was lyophilized from acetonitrile / water to give **11** (0.116 g, 41%) as a solid. MS: APCI (AP+): 469.2 (M)+, (AP-): 468.1 (M)-.

-278-

General Procedure for Example 111



5

Synthesis of 2: The carboxylic acid (1) is converted to the acid chloride (2) upon treatment with oxalyl chloride and DMF (catalytic) in dichloromethane.

Synthesis of 3: To the acid chloride (2) in dichloromethane is then added

- 10 NHG₁G₂ and triethylamine in dichloromethane. Alternatively NHG₁G₂ can be coupled directly to the carboxylic acid with addition of PyBOP and *N,N*-diisopropylethylamine in DMF. Reduction of the nitro group to the corresponding aniline is readily achieved with a transition metal such as palladium on carbon or Raney nickel and hydrogen to give 3.

15

Synthesis of 5: The carboxylic acid (4), aniline (3), EEDQ, and triethylamine are heated at reflux in chloroform to produce 5.

Synthesis of 6: A solution of compound 5, TFA, and DCM is stirred at room

- 20 temperature for 1h and then concentrated under reduced pressure. The resulting


-279-

oil is dissolved in THF and cooled to 0 °C followed by the addition of triethylamine or *N,N*-diisopropylethylamine (DIEA) and the appropriate isocyanate to produce compound 6.

5

Example 111

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-2-fluoro-phenyl)-amide]



10 **Step 1: (2R,4R)-4-hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (1).**

Cis-4-hydroxy-D-proline (15 g, 114 mmol) was dissolved in 150 mL of THF/ H₂O (2:1), added 2M aqueous NaOH solution (86 mL, 172 mmol), followed by Boc₂O (27 g, 126 mmol). Reaction was stirred at ambient

15 temperature overnight. The solution was acidified with 10% citric acid then extracted with EtOAc (2x250 mL), washed with water, brine, dried organics with MgSO₄, filtered and concentrated to give **1** (16 g, 61%) APCI (AP-): 230.1 (M-H)⁻.

20 **Step 2: (2R,4R)-4-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (2).**

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester (16 g, 69 mmol) was dissolved in 100 mL DMF, added imidazole (12 g, 173 mmol), dimethylaminopyridine (0.85 g, 6.9 mmol), and *tert*-butyldimethylsilyl

25 chloride (12.5 g, 83 mmol). Stirred at ambient temperature overnight, dissolved in

-280-

350 mL EtOAc, washed with 10% citric acid (3x200 mL), water (2x200 mL), brine, dried with MgSO₄, filtered, and concentrated to give 2 (20 g, 84%) APCI (AP⁻): 344.2 (M-H)⁻.

5 **Step 3: 3-Fluoro-4-nitro-benzoyl chloride (3).**

Into a solution of 3-fluoro-4-nitro-benzoic acid (4.0 g, 21.6 mmol) in anhydrous CH₂Cl₂ (200 mL) at 0°C was slowly added oxalyl chloride (2.45 mL, 28.0 mmol) and a drop of DMF. The solution was stirred and allowed to warm to RT overnight before concentrating under reduced pressure to give 3 (4.39 g,
10 100%).

Step 4: 3-Fluoro-N,N-dimethyl-4-nitro-benzamide (4).

Into a solution of 3 (1.50 g, 7.36 mmol) in CH₂Cl₂ (30 mL) at 0°C was added triethylamine (1.5 mL, 10.9 mmol) and dimethylamine 40% in water (0.83
15 mL, 7.36 mmol) dropwise. The solution was stirred and allowed to warm to RT overnight before concentrating under reduced pressure and adding EtOAc. The solution was washed sequentially with saturated NaHCO₃, water, and brine, before drying over MgSO₄ and concentrating under reduced pressure to give impure 4 (0.78 g, 50%). MS: APCI (AP⁺): 213.1 (M)⁺, (AP⁻): 212.1 (M)⁻.

20

Step 5: 4-Amino-3-fluoro-N,N-dimethyl-benzamide (5).

To a Parr apparatus was added 4 (0.78 g, 3.67 mmol), Raney nickel (1.2 g), and THF (50 mL). The vessel was sealed under a hydrogen atmosphere and shaken under pressure at room temperature for 15h. The vessel was then
25 depressurized and the solids filtered off, washing with THF. The filtrate was concentrated under reduced pressure to give impure 5 (0.66 g, 100%). MS: APCI (AP⁺): 183.1 (M)⁺, (AP⁻): 181.0 (M)⁻.

30 **Step 6: (2R,4R)-4-(tert-Butyl-dimethyl-silanyloxy)-2-(4-dimethylcarbamoyl-2-fluoro-phenylcarbamoyl)-pyrrolidine-1-carboxylic acid tert-butyl ester (6).**

Into a solution of 2 (1.25 g, 3.61 mmol) in CHCl₃ (25 mL) was added 5 (0.66 g, 3.61 mmol), EEDQ (1.07 g, 4.32 mmol), and triethylamine (0.74 mL,

-281-

5.33 mmol). The solution was stirred at reflux for 19h before cooling to RT and adding EtOAc. The solution was washed sequentially with 1N HCl, 0.1N NaOH, water, and brine, before drying over MgSO₄ and concentrating under reduced pressure. The crude material was purified on a silica gel column eluted with 20% moving to 50% EtOAc in hexanes. Combined and concentrated pure fractions. to give 6 (0.33 g, 18%). MS: APCI (AP+): 510.3 (M)+, (AP-): 508.4 (M)-.

Step 7: (2R,4R)-4-Hydroxy-pyrrolidine-2-carboxylic acid (4-dimethylcarbamoyl-2-fluoro-phenyl)-amide (7).

10 Into a solution of 6 (0.33 g, 0.647 mmol) in anhydrous CH₂Cl₂ (10 mL) was added TFA (2 mL). The solution was stirred at RT for 4h before concentrating under reduced pressure to give 7 (0.19 g, 100%).


Step 8: (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-2-fluoro-phenyl)-amide] (8).

15 Into a solution of 7 (0.19 g, 0.643 mmol) in anhydrous THF (10 mL) at 0°C was added *N,N*-diisopropylethylamine (DIEA) (0.56 mL, 3.21 mmol) and 4-chlorophenyl isocyanate (0.099 g, 0.643 mmol). The solution was stirred at RT overnight before concentrating under reduced pressure. Crystallized from EtOAc in hexanes and filtered to give 8 (0.12 g, 43%) as a solid. MS: APCI (AP+): 449.2 (M)+, (AP-): 447.1 (M)-.

Example 112

25 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-phenyl]-amide]

-282-




The compound was prepared as generally provided for Example 111. MS:
APCI (AP+): 457.1 (M)+, (AP-): 455.1 (M)-.

5

Example 113

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-phenyl]-amide]



10


The compound was prepared as generally provided for Example 111. MS:
APCI (AP+): 441.1 (M)+, (AP-): 439.1 (M)-.

15

Example 114

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2-methyl-pyrrolidine-1-carbonyl)-phenyl]-amide]

-283-




The compound was prepared as generally provided for Example 111. MS:
APCI (AP+): 471.2 (M)+, (AP-): 469.2 (M)-.

5

Example 115

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[[4-(ethyl-methyl-carbamoyl)-phenyl]-amide]

10




The compound was prepared as generally provided for Example 111. MS:
APCI (AP+): 445.2 (M)+, (AP-): 443.1 (M)-.

15

Example 116

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[[4-(dimethylcarbamoyl)-phenyl]-amide]

-284-




The compound was prepared as generally provided for Example 111. MS:
APCI (AP+): 431.1 (M)+, (AP-): 429.1 (M)-.

5

Example 117

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2R-methyl-pyrrolidine-1-carbonyl)-phenyl]-amide}



10


The compound was prepared as generally provided for Example 111. MS:
APCI (AP+): 471.2 (M)+, (AP-): 469.2 (M)-.

15

Example 118

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2S-methyl-pyrrolidine-1-carbonyl)-phenyl]-amide}

-285-




The compound was prepared as generally provided for Example 111. MS:
APCI (AP+): 471.2 (M)+, (AP-): 469.2 (M)-.

5

Example 119

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[[2-fluoro-4-(pyrrolidine-1-carbonyl)-phenyl]-amide]



10


The compound was prepared as generally provided for Example 111. MS:
APCI (AP+): 475.2 (M)+, (AP-): 473.1 (M)-.

15

Example 120

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[[4-(pyrrolidine-1-carbonyl)-2-pyrrolidin-1-yl-phenyl]-amide]

-286-




The compound was prepared as generally provided for Example 111. MS:
APCI (AP+): 526.2 (M)+, (AP-): 524.3 (M)-.

5

Example 121

(2R,4R)-4-[[1-(4-Chloro-phenylcarbamoyl)-4-hydroxy-pyrrolidine-2-carbonyl]-amino]-3-pyrrolidin-1-yl-benzoic acid methyl ester



10


The compound was prepared as generally provided for Example 111. MS:
APCI (AP+): 487.2 (M)+, (AP-): 485.2 (M)-.

Example 122

15

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(azetidine-1-carbonyl)-phenyl]-amide] 1-[(4-chloro-phenyl)-amide]

-287-




The compound was prepared as generally provided for Example 111o.

MS: APCI (AP+): 443.2 (M)+, (AP-): 441.1 (M)-.

5

Example 123

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(pyrrolidine-1-carbonyl)-phenyl]-amide]



10

The compound was prepared as generally provided for Example 111.

Formation of the 5-chloro-pyridin-2-yl moiety can be found in Example 26. MS:


APCI (AP+): 476.2 (M)+, (AP-): 474.2 (M)-.

15

Example 124

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-phenyl]-amide]

-288-



The compound was prepared as generally provided for Example 111.


Formation of the 5-chloro-pyridin-2-yl moiety can be found in Example 26. MS:

5 APCI (AP+): 458.2 (M)+, (AP-): 456.2 (M)-.

Example 125

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(4-dimethylcarbamoyl-phenyl)-amide]

10



The compound was prepared as generally provided for Example 111.

Formation of the 5-chloro-pyridin-2-yl moiety can be found in Example 26. MS:


15 APCI (AP+): 432.1 (M)+, (AP-): 430.1 (M)-.

Example 126

(2R,4R)-4-[[1-(4-Chloro-phenylcarbamoyl)-4-hydroxy-pyrrolidine-2-carbonyl]-amino]-3-dimethylamino-benzoic acid methyl ester

20

-289-




The compound was prepared as generally provided for Example 111. MS:
APCI (AP+): 461.2 (M)+, (AP-): 459.1 (M)-.

5

Example 127

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-phenyl]-amide]



10


The compound was prepared as generally provided for Example 110
and/or DD-2. MS: APCI (AP+): 471.2 (M)+, (AP-): 469.2 (M)-.

15

Example 128

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(dimethylcarbamoyl)-phenyl]-amide]

-290-




The compound was prepared as generally provided for Example 110 and/or 111. MS: APCI (AP+): 445.2 (M)+, (AP-): 443.2 (M)-.

5

Example 129

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-phenyl]-amide]



10


The compound was prepared as generally provided for Example 111. MS: APCI (AP+): 485.3 (M)+, (AP-): 483.3 (M)-.

15

Example 130

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-phenyl)-amide]

-291-




The compound was prepared as generally provided for Example 111. MS: APCI (AP+): 459.2 (M)+, (AP-): 457.2 (M)-.

5

Example 131

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(pyrrolidine-1-carbonyl)-phenyl]-amide]



10


The compound was prepared as generally provided for Example 110 and/or 111. MS: APCI (AP+): 489.3 (M)+, (AP-): 487.2 (M)-.

15

Example 132

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-2-fluoro-phenyl)-amide]

-292-




The compound was prepared as generally provided for Example 110 and/or 111. MS: APCI (AP+): 463.2 (M)+, (AP-): 461.2 (M)-.

5

Example 133

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(pyrrolidine-1-carbonyl)-phenyl]-amide]



10


The compound was prepared as generally provided for Example 111. MS: APCI (AP+): 503.3 (M)+, (AP-): 501.3 (M)-.

Example 134

15

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-2-fluoro-phenyl)-amide]

-293-




The compound was prepared as generally provided for Example 111. MS:
APCI (AP+): 477.3 (M)+, (AP-): 475.2 (M)-.

5

Example 135

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-methyl-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]



10


The compound was prepared as generally provided for Example 28. MS:
APCI (AP+): 485.3 (M)+, (AP-): 483.3 (M)-.

15

Example 136

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-methyl-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]

-294-




The compound was prepared as generally provided for Example 28. MS:
APCI (AP+): 499.2 (M)+, (AP-): 497.2 (M)-.

5

Example 137

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-quinolin-8-yl-phenyl)-amide]



10


The compound was prepared as generally provided for Example 19. MS:
APCI (AP+): 489.1 (M)+, (AP-): 488.1 (M)-.

15

Example 138

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3,5-difluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]

-295-




The compound was prepared as generally provided for Example 19. MS:
APCI (AP+): 534.1 (M)+, (AP-): 532.0 (M)-.

5

Example 139

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-2-methyl-biphenyl-4-yl)-amide]



10


The compound was prepared as generally provided for Example 19. MS:
APCI (AP+): 512.1 (M)+, (AP-): 510.1 (M)-.

15

Example 140

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-biphenyl-4-yl)-amide]

-296-




The compound was prepared as generally provided for Example 17. MS:
APCI (AP+): 514.0 (M)+, (AP-): 512.0 (M)-.

5

Example 141

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[(2'-methanesulfinyl-biphenyl-4-yl)-amide]



10


The compound was prepared as generally provided for Example 17. MS:
APCI (AP+): 498.2 (M)+, (AP-): 496.2 (M)-.

15

Example 142

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[(3-methyl-2'-methylsulfanyl-biphenyl-4-yl)-amide]

-297-




The compound was prepared as generally provided for Example 17. MS:
APCI (AP+): 496.1 (M)+, (AP-): 494.1 (M)-.

5

Example 143

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[(2-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]

10




The compound was prepared as generally provided for Example 17. MS:
APCI (AP+): 532.0 (M)+, (AP-): 530.0 (M)-.

15

Example 144

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[(2'-methanesulfonyl-3-methyl-biphenyl-4-yl)-amide]

-298-




The compound was prepared as generally provided for Example 17. MS: APCI (AP+): 528.1 (M)+, (AP-): 526.1 (M)-.

5

Example 145

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3-trifluoromethyl-biphenyl-4-yl)-amide]



10


The compound was prepared as generally provided for Example 17. MS: APCI (AP+): 582.1 (M)+, (AP-): 580.1 (M)-.

15

Example 146

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methoxy-biphenyl-4-yl)-amide]

-299-

**Step 1: 2-Methoxy-4'-nitro-biphenyl (146a).**

1-Bromo-4-nitrobenzene (1.00g, 4.95mmol), 2-methoxyphenylboronic
5 acid (1.13g, 7.43mmol), K_3PO_4 (1.58g, 7.43mmol), and anhydrous DMF (5mL)
were combined in a flask and degassed with argon. Tetrakis(triphenylphosphine)
palladium(0) (0.858g, 0.743mmol) was added to the mixture which was again
degassed with argon. The mixture was heated at 110°C for 3h before cooling,
diluting with EtOAc, and filtering through a plug of silica gel. The solution was
10 washed sequentially with water and brine before drying over $MgSO_4$. The
solution was concentrated under reduced pressure and purified by flash
chromatography to provide **146-a** (1.05g, 93%) as a yellow solid. MS: APCI
(AP+): 230.1 (M)+.

Step 2: 2'-Methoxy-biphenyl-4-ylamine (146-b).


To a Parr apparatus was added **146-a** (1.05g, 4.58mmol), Raney nickel
(0.66g), and a 4:1 mixture of THF:MeOH (50mL). The vessel was sealed under a
hydrogen atmosphere and shaken under pressure at RT for 18h. The vessel was
then depressurized and the solids filtered off, washing with THF. The filtrate was
20 concentrated under reduced pressure to reveal an unknown amount of **146-b** as a
tan oil that was carried on without further purification. MS: APCI (AP+): 200.0
(M)+.

The title compound (**146**) was prepared from **146b** as generally provided
25 for in Examples 26-28. MS: APCI (AP+): 466.1 (M)+.

-300-

Example 147

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-hydroxy-biphenyl-4-yl)-amide]



5


In a flask was dissolved 146a (0.100g, 0.215mmol) in anhydrous DCM (5mL). At 0°C, boron tribromide (0.203mL, 2.15mmol) was added dropwise. The solution was stirred at RT for 25min before concentrating under reduced pressure. The crude material was purified by silica gel flash chromatography followed by lyophilization from acetonitrile / water to provide the title compound (0.039g, 40%) as a white solid. MS: APCI (AP+): 452.1 (M)+.

10

Example 148

15

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-iodo-phenyl)-amide]




20

The compound was prepared as generally provided for Example 28. MS: APCI (AP+): 518.0 (M)+.

-301-

Example 149


(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-{{2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl}-amide}



The compound was prepared as generally provided for Example 28. MS:
APCI (AP+): 485.2 (M)+.

Example 150

(2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-
yl)-amide] 2-{{2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl}-amide}




The compound was prepared as generally provided for Example 28 and
Example 3, step 3a. MS: APCI (AP+): 486.2 (M)+.

Example 151

-302-


(2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-hydroxy-2-oxo-piperidin-1-yl)-phenyl]-amide]



- 5 The compound was prepared as generally provided for Example 28. MS:
APCI (AP+): 505.3 (M)+.

Example 152

- 10 (2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-tetrahydro-pyrimidin-1-yl)-phenyl]-amide]




- 15 The compound was prepared as generally provided for Example 28. MS:
APCI (AP+): 490.3 (M)+.

Example 153

- 20 (2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-imidazolidin-1-yl)-phenyl]-amide]

-303-




The compound was prepared as generally provided for Example 28. MS:
APCI (AP+): 476.2 (M)+.

5

Example 154

(2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-oxazolidin-3-yl)-phenyl]-amide]



10


The compound was prepared as generally provided for Example 28. MS:
APCI (AP+): 477.2 (M)+.

15

Example 155

(2R, 4R)-1-(4-[[1-(4-Chloro-phenyl)carbamoyl]-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-2-oxo-piperidine-3-carboxylic acid ethyl ester

-304-




The compound was prepared as generally provided for Example 28. MS:
APCI (AP+): 561.3 (M)+.

5

Example 156

(2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-{[2-fluoro-4-(3-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide}



10


The compound was prepared as generally provided for Example 28. MS:
APCI (AP+): 515.2 (M)+.

15

Example 157

(2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-
yl)-amide] 2-{[2-fluoro-4-(3-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide}

-305-




The compound was prepared as generally provided for Example 28 and Example 23, step 3a. MS: APCI (AP+): 516.2 (M)+.

5

Example 158

(2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide]



10

Step 1: 3-Methyl-1-(4-nitro-phenyl)-1H-pyridin-2-one (158-a).

1-Iodo-4-nitrobenzene (1.61g, 6.45mmol) was combined with 3-methyl-2-pyridone (0.845g, 7.74mmol), CuI (0.246g, 1.29mmol), and K₃PO₄ (2.74g, 12.9mmol). 1,4-Dioxane (6mL) was added followed by *trans*-1,2-diaminocyclohexane (0.194mL, 1.61mmol). The mixture was heated to reflux for 19h before cooling and diluting with EtOAc. The mixture was filtered through a plug of silica, eluting with EtOAc, and the filtrate concentrated under reduced pressure. Purification of the crude product by flash chromatography provided 158-a (0.528g, 36%) as a brown solid. MS: APCI (AP+): 231.1 (M)+.

20

-306-


Step 2: 1-(4-Amino-phenyl)-3-methyl-1H-pyridin-2-one (158-b).

To a flask was added 158a (0.523g, 2.27mmol), glacial acetic acid (7.6mL), conc. HCl (3.8mL, 45.4mmol), and mossy tin (0.539g, 4.54mmol). The mixture was heated to reflux for 90min before cooling and concentrating under reduced pressure. The residue was dissolved in EtOAc and washed with aq. 1N NaOH until shown basic with pH paper. The aqueous layer was extracted twice with EtOAc and the combined organics washed with brine and dried over MgSO₄. After placing under vacuum, 158-b (0.407g, 89%) was revealed as a tan solid that was carried on without further purification. MS: APCI (AP+): 201.1 (M)+.

The title compound (158) was prepared from 158-b as generally provided for Example 28. MS: APCI (AP+): 481.3 (M)+.

Example 159

(2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(3-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide]




The compound was prepared as generally provided for Example 158 and Example 3, step 3a. MS: APCI (AP+): 482.3 (M)+.

Example 160

(2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(morpholin-4-yl)-phenyl]-amide]

-307-

**Step 1: 4-(4-Nitro-phenyl)-morpholine (160-a).**


1-Fluoro-4-nitrobenzene (3.00mL, 28.3mmol) was combined in a flask
 5 with anhydrous 2-propanol (28mL), triethylamine (4.34mL, 31.1mmol), and
 morpholine (2.47mL, 28.3mmol). The mixture was refluxed for 3.5h before
 adding additional morpholine (2.47mL, 28.3mmol) and refluxing for 20h. The
 mixture was then cooled and concentrated under reduced pressure. Purification by
 silica gel flash chromatography provided **160-a** (6.09g, 98%) as a 95% pure solid.
 10 MS: APCI (AP-): 207.1 (M)-.

The title compound (**160**) was prepared from **160-a** as generally provided
 for **146** and Example 28. MS: APCI (AP+): 459.3 (M)+.

15

Example 161

**(2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
 amide] 2-[[4-(2-methyl-5-oxo-pyrrolidin-1-yl)-phenyl]-amide]**




20

The compound was prepared as generally provided for in Example 28 and
 146. MS: APCI (AP+): 471.3 (M)+.

-308-

Example 162


(2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(2-methyl-5-oxo-pyrrolidin-1-yl)-phenyl]-amide]



The compound was prepared as generally provided for in Example 3, step 3a, and Example 161. MS: APCI (AP+): 472.2 (M)+.

Example 163

(2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(4-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide]




The compound was prepared as generally provided for Example 28. MS: APCI (AP+): 499.2 (M)+.

Example 164


-309-

(2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(4-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide]]



- 5 The compound was prepared as generally provided for Example 28 and Example 3, step 3a. MS: APCI (AP+): 500.2 (M)+.

General Procedure for Examples 165-173:



10

Synthesis of 2: A mixture of 1 (where Z is a halide), B (where B is a nitrogen in a ring), K_2CO_3 , CuI , and 8-hydroxyquinoline is heated in DMSO at 130°C to obtain compound 2.

15

Synthesis of 4: The appropriate aniline (2), EEDQ , triethylamine and carboxylic acid (3) are heated at reflux in chloroform to produce 4.


-310-

Synthesis of 5: A solution of compound 4, TFA and DCM is stirred at room temperature for 1h and then concentrated under reduced pressure. The resulting oil is dissolved in THF and cooled to 0 °C followed by the addition of triethylamine and the appropriate isocyanate to produce compound 5.

5

Example 166

(2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide]



10

Step 1: 1-(4-Amino-3-fluoro-phenyl)-5-methyl-1H-pyridin-2-one (167-a).

2-Fluoro-4-iodoaniline (0.905g, 42.2mmol) was combined with 5-methyl-1H-pyridin-2-one (0.500g, 4.58mmol), 8-hydroxyquinoline (0.083g, 0.573mmol),
15 CuI (0.109g, 0.0573mmol), and K₂CO₃ (0.580g, 4.20mmol), in DMSO (3mL). The mixture was degassed with a stream of argon and then heated at reflux for 16h before cooling to RT. 10% aq. NH₄OH and EtOAc were added, and the mixture was filtered through a plug of layered celite and decolorizing charcoal, eluting with EtOAc. The filtrate was concentrated under reduced pressure.
20 Purification of the crude product by silica gel flash chromatography revealed **167-a** (0.673g, 81%) as a white solid. MS: APCI (AP+): 219.1 (M)+.


The title compound (**166**) was prepared from **166-a** as generally provided for in Example 28. MS: APCI (AP+): 499.2 (M)+.

25

Example 167

-311-

(2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(5-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide]




5

The compound was prepared as generally provided for in Example 166 and Example 3, step 3a. MS: APCI (AP+): 500.2 (M)+.

Example 168

10

(2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(4-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide]



15


The compound was prepared as generally provided for in Example 166. MS: APCI (AP+): 515.2 (M)+.

Example 169

20

(2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(4-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide]

-312-




The compound was prepared as generally provided for in Example 167.

MS: APCI (AP+): 516.2 (M)+.

5

Example 170

(2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide}



10

The compound was prepared as generally provided for in Example 166.


MS: APCI (AP+): 499.2 (M)+.

15

Example 171

(2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(3-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide}

-313-




The compound was prepared as generally provided for in Example 166.

MS: APCI (AP+): 500.2 (M)+.

5

Example 172

(2R, 4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(6-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide}



10

The compound was prepared as generally provided for in Example 166.


MS: APCI (AP+): 499.2 (M)+.

15

Example 173

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(5-chloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide]

-314-




The compound was prepared as generally provided for in Example 166.
 MS: APCI (AP+): 519.1 (M)+.

5

General Procedure for the Preparation of Examples 174-180

Route 1



10 Synthesis of 2:


Aniline **1** is converted to an acrylamide **2** by the addition of acryoyl chloride and a base, such as saturated sodium bicarbonate, in a solvent such as ethyl acetate at room temperature. Alternatively, the aniline may be converted to **2** by the addition of an acrylic acid and adding a coupling reagent such as

15 dicyclohexylcarbodi-imide (DCC), O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU), or EEDQ.

-315-

Synthesis of 3 and 4:

To the acrylamide **2** is added an excess of trimethylsilyl diazomethane in a solvent such as ethyl acetate or dichloromethane. The resulting dihydropyrazole **3** may then be treated with an isocyanate, in the presence of base such as pyridine or triethylamine, in a solvent such as dichloromethane, to afford compound **4**.

Route 2

10

Synthesis of 5-8:


The chemistry for the preparation of compounds **5-8** is similar to that described in **Route 1**.

Synthesis of 9:

Compound **8** may be converted to **9** by a Suzuki coupling with a boronic acid, although other coupling conditions commonly known to the skilled practitioner may also be used. This route is particularly useful for compounds containing a biaryl A-B group. In situations where a tert-butylsulphonamide is present in B, which occurs in some of the particularly preferred compounds, the sulphonamide may be formed by stirring with trifluoroacetic acid for 16 h.

20

-316-

Route 3**5 Synthesis of 10:**

The synthetic routes commence with formation of acrylamide **10** as described for **Route 1**.

Synthesis of 11:

- 10 Ethyl diazoacetate is mixed with acrylamide **10** to afford the appropriately substituted dihydropyrazole **11**.

Synthesis of 12 and 13:


- 15 Similar chemistry to that described in **Route 1** allows conversion of **11** to **12**. Reduction of the ester moiety in **12** is then achieved using a reducing agent such as super-hydride to afford the alcohol **13**.

Route 4

- 20 Alternatively compounds may be prepared according to scheme 2d. Here the acrylic ester (**14**) is reacted with a diazo compound to form the dihydropyrazole (**15**). Addition of a chloroformate affords compounds of formula (**16**). Deprotection of the ester is achieved by acid, such as TFA or hydrogen

-317-


chloride to afford compounds of formula (17). Addition of an aniline with a coupling reagent such as EEDQ and base affords compounds of formula (18). Deprotection of the carbamate with Pd/C and hydrogen affords compounds of formula (19) which may then be reacted with isocyanates to afford compounds of formula (20)



Example 174

5-Methyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]

10



-318-

Step 1: N-(2'-tert-Butylsulfamoyl-3-fluoro-biphenyl-4-yl)-2-methyl-acrylamide

To 4'-Amino-3'-fluoro-biphenyl-2-sulfonic acid tert-butylamide (5.000 g, 15.51 mmoles) at room temperature in ethyl acetate (50 ml) was added saturated sodium bicarbonate, aqueous, 50 ml and then 2-methyl-acryloyl chloride (1.25 equiv.). Stirred for 2 hours then added a further 1 ml of the acid chloride. After 2 hours an aliquot showed product by NMR. Product seems to co-run with starting material. Extracted into ethyl acetate, washed with brine, dried MgSO_4 , and then concentrated in vacuo. Collected some crystalline product from ethyl acetate hexane, 0.48 g. Columned mother liquor to afford product. Combined with previously crystallized fraction to afford N-(2'-tert-Butylsulfamoyl-3-fluoro-biphenyl-4-yl)-2-methyl-acrylamide (4.520 g). APCI (AP+): 391 (M+H)⁺.

Step 2: 3-Methyl-3,4-dihydro-2H-pyrazole-3-carboxylic acid (2'-tert-butylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide

To N-(2'-tert-Butylsulfamoyl-3-fluoro-biphenyl-4-yl)-2-methyl-acrylamide (0.525g) in ethyl acetate (3.5ml) was added 2M trimethylsilyl diazomethane (3.5ml) and the mixture stirred at room temperature overnight. NMR indicated complete reaction. Evaporated and then dissolved in acetonitrile (20ml) and treated with aq. HF (1ml). Stirred for 1h. Added sat. NaHCO_3 (5ml), extracted into EtOAc (25ml). Washed with brine, dried MgSO_4 , evaporated in vacuo from methylene chloride to afford a white foam of the product: 3-Methyl-3,4-dihydro-2H-pyrazole-3-carboxylic acid (2'-tert-butylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide (0.592g). APCI (AP+): 433 (M+H)⁺.

25

Step 3: 5-Methyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 5-[(2'-tert-butylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide]

To a solution of 3-methyl-3,4-dihydro-2H-pyrazole-3-carboxylic acid (2'-tert-butylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide (0.416g) in methylene chloride (4ml) was added pyridine (1ml) and 4-chlorophenyl isocyanate (1.25g). Stirred at room temperature for 16 hours. A further amount of 4-chlorophenyl isocyanate (0.200g) and pyridine (0.2ml) was added. Stirred for 48 hours. Columned, silica

30

-319-

gel - eluant 10-100 EtOAc in hexane over 25mins. Affords: 5-methyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 5-[(2'-tert-butylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide]; 0.365g


5 **Step 4: 5-Methyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]**

To 5-methyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 5-[(2'-tert-butylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide] (0.300g) was added trifluoroacetic acid (10.0g). Stirred at room temperature
10 overnight and then evaporated the TFA. Redissolved in hot methanol and added water until cloudy white. Lyophilized to afford 5-methyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 0.256 g. APCI (AP+): 530 (M+H)⁺

15

Example 175

3-Hydroxymethyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]



20

Step 1: N-(3-Fluoro-2'-methanesulfonyl-biphenyl-4-yl)-acrylamide

To 3-Fluoro-2'-methanesulfonyl-biphenyl-4-ylamine (8.510 g) in ethyl acetate (400 ml) was added sat. NaHCO₃ (100 ml) and then acryloyl chloride (1.3 equiv.). Stirred at room temperature for 3 hours. Washed EtOAc extract with
25 brine, dried over MgSO₄ and evaporated in vacuo. Triturated with ether and then

-320-

filtered to afford N-(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-acrylamide (9.670 g, 94%).

Step 2: 5-(3-Fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-4,5-dihydro-1H-pyrazole-3-carboxylic acid ethyl ester

To N-(3-Fluoro-2'-methanesulfonyl-biphenyl-4-yl)-acrylamide (4.47g) in ethyl acetate (150ml) was added ethyl diazoacetate (1.5 equiv.) and the mixture heated at 60 °C for 24 hours. Evaporated and then columned: eluant 10-100 EtOAc in hexane. Collected 5-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-4,5-dihydro-1H-pyrazole-3-carboxylic acid ethyl ester (1.67g)

Step 3: 1-(4-Chloro-phenylcarbamoyl)-5-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-4,5-dihydro-1H-pyrazole-3-carboxylic acid ethyl ester

To a solution of 5-(3-Fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-4,5-dihydro-1H-pyrazole-3-carboxylic acid ethyl ester (1.520g) in methylene chloride (12ml) was added pyridine (4 equiv.) and 4-chlorophenyl isocyanate (1.25 equiv.). Stirred at room temperature for 16 hours. Evaporated and recorded crude 1H: 80156x134RM1 - revealed expected product and relatively clean. Purified by silica gel chromatography, eluant 10-100% EtOAc in hexane over 25 mins. Collected 1-(4-chloro-phenylcarbamoyl)-5-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-4,5-dihydro-1H-pyrazole-3-carboxylic acid ethyl ester (1.480g).

Step 4: 3-Hydroxymethyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]

To a solution of 1-(4-Chloro-phenylcarbamoyl)-5-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-4,5-dihydro-1H-pyrazole-3-carboxylic acid ethyl ester (0.228g) in THF (3ml) at 0 °C was added a 1M solution of superhydride in THF (1.17ml). Stirred for 30 mins and then quenched with sat. NH₄Cl. Extracted into EtOAc and washed with brine. Evaporated in vacuo. Crude 1H indicated clean reaction to product. Column - eluant 0-5% methanol in ethyl


-321-

acetate. Affords 3-Hydroxymethyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] after lyophilization from MeCN/H₂O: 0.230 g; MS (APCI) +ve: 545

5

Example 176


4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]



This compound was prepared using the same procedures as found for
 10 Example 175 with acryloyl chloride substituted for 2-Methyl-acryloyl chloride in
 step 1. APCI (AP-): 514 (M-H)⁻.

Example 177

15 **(5R) 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide]**



This compound was prepared using the same procedures as found for
 Example 176 and the enantiomers separated by chiral HPLC Column: Chiral Pak


-322-

AD 250x4.6 mm; 254 nm, mobile phase: Hexane (20); Ethanol (20); Methanol (60) Affords: (R) enantiomer @ 11.038 mins APCI (AP-): 514 (M-H)⁻ and (S) enantiomer @ 36.084 mins APCI (AP-): 514 (M-H)⁻.

5

Example 178

4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide]




This compound was prepared using the same procedures as found for
10 Example 175 with acryloyl chloride substituted for 2-Methyl-acryloyl chloride in step 1 and 1-(4-Amino-3-fluoro-phenyl)-piperidin-2-one substituted for 4'-Amino-3'-fluoro-biphenyl-2-sulfonic acid tert-butylamide. APCI (AP-): 456 (M-H)⁻.

15

Example 179

4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]




-323-

This compound was prepared using the same procedures as found for Example 176 with acryloyl chloride substituted for 2-Methyl-acryloyl chloride in step 1 and 3-Fluoro-2'-methanesulfonyl-biphenyl-4-ylamine substituted for 4'-Amino-3'-fluoro-biphenyl-2-sulfonic acid tert-butylamide. APCI (AP-): 513 (M-H)⁻

Example 180

1-(4-Chloro-phenylcarbamoyl)-5-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-4,5-dihydro-1H-pyrazole-3-carboxylic acid ethyl ester



10

This compound was prepared using the same procedures as found for Example 175 wherein step 4 was omitted. APCI (AP-): 585 (M-H)⁻.

Example 181

FORMULATIONS

15

The compounds of the present invention can be administered alone or in combination with one or more therapeutic agents. These include, for example, other anticoagulant, antiplatelet, or platelet inhibitory agents which include non-steroidal anti-inflammatory agents such as aspirin, ibuprofen, naproxen sodium, indomethacin, piroxica, and ticlopidine; thrombin inhibitors such as argatroban, efegatran, inogatran, factor VIIa inhibitors, thrombolytic or fibrinolytic agents such as tissue plasminogen activator, urokinase or streptokinase; GP IIb-IIIa antagonists, and P2Y12 antagonists.

20

-324-

The compounds are thus well suited to formulation for convenient administration to mammals for the prevention and treatment of such disorders.

The following examples further illustrate typical formulations provided by
5 the invention.

Formulation 1

Ingredient	Amount
compound of Formulas I-V	0.5 to 800 mg
sodium benzoate	5 mg
isotonic saline	1000 mL

The above ingredients are mixed and dissolved in the saline for IV
10 administration to a human suffering from, for example, arterial thrombosis.

Formulation 2

Ingredient	Amount
compound of Formulas I-V	0.5 to 800 mg
cellulose, microcrystalline	400 mg
stearic acid	5 mg
silicon dioxide	10 mg
sugar, confectionery	50 mg

The ingredients are blended to uniformity and pressed into a tablet that is
15 well suited for oral administration to a human for preventing, for example, cerebral infarction.

-325-

Formulation 3

Ingredient	Amount
compound of Formulas I-I	0.5 to 800 mg
starch, dried	250 mg
magnesium stearate	10 mg

The ingredients are combined and milled to afford material suitable for ,
filling hard gelatin capsules administered to humans suffering from, for example,
5 venous thrombosis.

Formulation 4

Ingredient	Amount	% wt./(total wt.)
compound of Formulas I-I	1 to 50	
Polyethylene glycol 1000	32 to 75	
Polyethylene glycol 4000	16 to 25	


The ingredients are combined via melting and then poured into molds
10 containing 2.5 g total weight.

While embodiments of the invention have been illustrated and described, it
is not intended that these embodiments illustrate and describe all possible forms of
the invention. Rather, the words used in the specification are words of description
15 rather than limitation, and it is understood that various changes may be made
without departing from the spirit and scope of the invention.

-326-

What is claimed is:

1. A compound of Formula I:



or a pharmaceutically acceptable salt thereof wherein:

- A is aryl or substituted aryl or monocyclic heteroaryl or substituted
10 monocyclic heteroaryl;

B is $\text{-NH}-\overset{\text{O}}{\parallel}{\text{C}}-(\text{C}_1\text{-C}_6)\text{alkyl}$, $\text{-NH}-\overset{\text{O}}{\parallel}{\text{C}}-(\text{C}_3\text{-C}_7)\text{cycloalkyl}$, -

$\text{NH}-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}(\text{C}_1\text{-C}_6\text{ alkyl})$, $\overset{\text{O}}{\parallel}{\text{C}}-\text{R}^1$, $(\text{C}_3\text{-C}_7)\text{cycloalkyl}$, $(\text{C}_3\text{-C}_7)\text{heterocyclo}$, $(\text{C}_4\text{-C}_7)\text{cycloalkenyl}$, unsaturated $(\text{C}_4\text{-C}_7)\text{heterocyclo}$, aryl, or

- heteroaryl, any of which may be optionally substituted by halo, $(\text{C}_1\text{-C}_6)\text{alkyl}$, or
15 halo $(\text{C}_1\text{-C}_6)\text{alkyl}$, $\text{O}-(\text{C}_1\text{-C}_6)$, -CN , haloalkyl, amino, alkylamino, amidino, amido,
or sulfonamido;

- C is phenyl or heteroaryl, wherein phenyl or heteroaryl is optionally
substituted with one or more substituents selected from aryl, heteroaryl, halogen,
hydroxy, $\text{-CO}_2\text{R}^2$, -COR^2 , $\text{-CONR}^2\text{R}^{2'}$, alkoxy, alkyl, -CN , haloalkyl, amino,
20 alkylamino, amidino, amido, or sulfonamido;

G is H, halo, $(\text{C}_1\text{-C}_6)\text{alkyl}$, halo $(\text{C}_1\text{-C}_6)\text{alkyl}$, hydroxy $(\text{C}_1\text{-C}_6)\text{alkyl}$,
 $\text{-CH}_2\text{O}-(\text{C}_1\text{-C}_6)\text{alkyl}$, $\text{-CH}_2\text{CO}_2(\text{C}_1\text{-C}_6)\text{alkyl}$, $\text{-CH}_2\text{-NR}^2\text{R}^{2'}$, or $\text{-CH}_2\text{-CONH}(\text{C}_1\text{-C}_6)\text{alkyl}$;

W^1 is a saturated or unsaturated, substituted or unsubstituted hydrocarbon

-327-


chain or hydrocarbon-heteroatom chain having from 2 to 6 atoms, wherein W¹ connects the nitrogen atom at position 1 to the carbon atom at position 2 to form a four to eight membered ring;

R¹ is (C₁-C₆)alkoxy, (C₃-C₇)cycloalkyl, (C₃-C₇)heterocycloalkyl, (C₄-C₇)cycloalkenyl, (C₄-C₇)heterocycloalkenyl, aryl, monocyclic heteroaryl, or -NR³R⁴;

R² and R^{2'} are each independently H or (C₁-C₆)alkyl; and

R³ and R⁴ are each independently H, (C₁-C₆)alkyl, aralkyl, aryl, monocyclic heteroaryl, alkoxycarbonyl, aralkoxycarbonyl, -SO₂alkyl, or joined together to form a saturated or unsaturated 3 to 7 membered ring.

2. A compound of formula IV:



15

or a pharmaceutically acceptable salt thereof wherein

“---” is a bond or is absent;

Z is C-H, C-halo, C-(C₁-C₆)alkyl, C-halo(C₁-C₆)alkyl, C-(C₁-C₆)alkoxy, or N;

20 X^{II} is CH₂, CH, NH or N;

R¹⁰ and R¹¹ are each independently H, -OH, halo, alkyl, haloalkyl, -NR⁸R⁹, -OR², -CN, -CH₂OH, -CH₂-NR³R⁴, aryl, monocyclic heteroaryl,

-328-

alkylaryl, $-\text{CH}=\text{O}$, $-\text{CH}_2\text{OR}^2$, $-\text{COR}^2$, $-\text{CO}_2\text{R}^2$, or $-\text{CONR}^3\text{R}^4$ or are taken together to form $=\text{O}$, $=\text{NOR}^2$, $=\text{C}(\text{C}_1\text{-C}_6\text{alkyl})_2$, or $=\text{CR}^2\text{H}$, with the proviso that when “---” is a bond, R^{11} is absent, and X^{II} is CH or N, and R^{10} is H, -OH, halo, alkyl, haloalkyl, $-\text{NR}^8\text{R}^9$, $-\text{OR}^2$, $-\text{CN}$, $-\text{CH}_2\text{OH}$, $-\text{CH}_2\text{-NR}^3\text{R}^4$, aryl, monocyclic

5 heteroaryl, alkylaryl, $-\text{CH}=\text{O}$, $-\text{CH}_2\text{OR}^2$, $-\text{COR}^2$, $-\text{CO}_2\text{R}^2$, or $-\text{CONR}^3\text{R}^4$;

B is $-\text{NH}-\overset{\text{O}}{\parallel}{\text{C}}-(\text{C}_1\text{-C}_6)\text{alkyl}$, $-\text{NH}-\overset{\text{O}}{\parallel}{\text{C}}-(\text{C}_3\text{-C}_7)\text{cycloalkyl}$ $-\text{NH}(\text{C}_1\text{-}$

$\text{C}_6)\text{alkanoyl}$, $\text{---}\overset{\text{O}}{\parallel}{\text{C}}\text{---}\text{R}^1$, $(\text{C}_3\text{-C}_7)\text{cycloalkyl}$, $(\text{C}_3\text{-C}_7)\text{heterocycloalkyl}$, $(\text{C}_4\text{-C}_7)\text{cycloalkenyl}$, $(\text{C}_4\text{-C}_7)\text{heterocycloalkenyl}$, aryl, or heteroaryl, any of which may be optionally substituted, wherein R^1 is $(\text{C}_1\text{-C}_6)\text{alkoxy}$, $(\text{C}_3\text{-C}_7)\text{cycloalkyl}$, $(\text{C}_3\text{-C}_7)\text{heterocycloalkyl}$, $(\text{C}_4\text{-C}_7)\text{cycloalkenyl}$, $(\text{C}_4\text{-C}_7)\text{heterocycloalkenyl}$, aryl, 10 monocyclic heteroaryl, or $-\text{NR}^3\text{R}^4$;

G is H, halo, $(\text{C}_1\text{-C}_6)\text{alkyl}$, halo $(\text{C}_1\text{-C}_6)\text{alkyl}$, hydroxy $(\text{C}_1\text{-C}_6)\text{alkyl}$, $-\text{CH}_2\text{O}-(\text{C}_1\text{-C}_6)\text{alkyl}$, $-\text{CH}_2\text{-CO}_2(\text{C}_1\text{-C}_6)\text{alkyl}$, $-\text{CH}_2\text{-CONH}_2$, or $-\text{CH}_2\text{-CONH}(\text{C}_1\text{-C}_6)\text{alkyl}$;

15 R^2 and $\text{R}^{2'}$ are each independently H or $(\text{C}_1\text{-C}_6)\text{alkyl}$; and

R^3 and R^4 are independently H, $(\text{C}_1\text{-C}_6)\text{alkyl}$, aralkyl, aryl, monocyclic heteroaryl, alkoxycarbonyl, aralkoxycarbonyl, $-\text{SO}_2\text{alkyl}$, or joined together to form a saturated or unsaturated 5 to 7 membered ring;

R^{12} and R^{13} are each independently H, halo, $(\text{C}_1\text{-C}_6)\text{alkyl}$, or halo $(\text{C}_1\text{-C}_6)\text{alkyl}$, or $(\text{C}_1\text{-C}_6)\text{alkoxy}$; 20


R^{14} and R^{15} are each independently H, halo, alkyl, or haloalkyl, or NR^8R^9 wherein R^8 and R^9 are as defined for R^3 and R^4 ;

X and Y are each C, or one of X and Y is C and the other is N, provided that when one of X or Y is N, R^{14} or R^{15} is absent at that position.

25


3. The compound of claim 2, wherein B is

-329-




Q is NH or is absent; and wherein $R^{1'}$ is OH, $-O-(C_1-C_6)$, or $NR^{3'}R^{4'}$, wherein $R^{3'}$ and $R^{4'}$ are independently H, (C_1-C_6) alkyl, or are joined together to form a optionally substituted saturated or unsaturated 5 to 7 membered ring;

5




, wherein “~” indicates the point of attachment, or B is optionally substituted phenyl or pyridyl with formyl, H_2NSO_2 , $MeSO_2$, $MeSO$, MeS , (C_1-C_6) alkoxycarbonyl, cyano, aminomethyl, methoxy, or hydroxy;



10

, wherein “~” indicates the point of attachment and which is an optionally substituted 4, 5, 6 or 7-membered ring; wherein R^x and R^y are H, halo, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, or (C_1-C_6) carboxyalkyl, or taken together are $O=$; X^{III} is CH_2 , $CH-OH$, $CH-CO_2(C_1-C_6)$ alkyl, O, S, NH, or $N(C_1-C_6)$ alkyl, provided that when R^x and R^y taken together are $O=$, X^{III} is CH_2 .

15 provided that when R^x and R^y taken together are $O=$, X^{III} is CH_2 ;




R^{18} or a tautomer thereof, wherein Y^{II} is CH_2 , $CH(C_1-C_6)$ alkyl, or N and R^{18} is H, (C_1-C_6) alkyl, hydroxymethyl, $CH_2O-(C_1-C_6)$ alkyl, $CONR^2R^{2'}$, or $CH_2NR^2R^{2'}$.


20

-330-

a 5-membered heteroaryl ring which contains at least one nitrogen atom, which is selected from imidazolyl, pyrazolyl, triazolyl, tetrazolyl, oxadiazolyl, thiadiazolyl, oxazolyl, isoxazolyl, thiazolyl, or isothiazolyl, any of which may be optionally substituted with halo, (C₁-C₆)alkyl, hydroxy(C₁-C₆)alkyl, CH₂O-(C₁-



5 C₆)alkyl, or halo(C₁-C₆)alkyl, or is $R^{18}-\text{---}-Z^a$, wherein "---" indicates the point of attachment; X^a, Y^a, and Z^a are each independently CR^c or N, wherein R^c is H or (C₁-C₆)alkyl; and R¹⁸ is H, (C₁-C₆)alkyl, hydroxymethyl, CH₂O-(C₁-C₆)alkyl, or NR³R⁴;



10 $\overset{1}{R}^{20}$, wherein “~” indicates the point of attachment;
J₁, J₂, J₃, and J₄ are each C, or any of J₁, J₂, J₃, or J₄ is N; and R¹⁹, R²⁰, R²¹, and R²²
are each independently H, halo, hydroxy, NH₂, NR²³R²⁴, NO₂, SH, (C₁-C₆)alkyl,
halo(C₁-C₆)alkyl, (C₁-C₆)alkanoyl, (C₁-C₆)alkoxycarbonyl, or (C₁-C₆)alkoxy,
wherein R²³ and R²⁴ are each independently H, (C₁-C₆)alkyl, aralkyl, aryl,
15 monocyclic heteroaryl, alkoxycarbonyl, aralkoxycarbonyl, -SO₂alkyl, or joined
together to form a saturated or unsaturated 3 to 7 membered ring, or R¹⁹ and R²⁰,
R²⁰ and R²¹, or R²¹ and R²², together with the carbons to which they are attached,
form a 5, 6, or seven membered saturated or unsaturated cycloalkyl or
heterocycloalkyl ring, or an aryl or heteroaryl ring; provided that when any of J₁,
20 J₂, J₃, or J₄ is N, R¹⁹, R²⁰, R²¹ or R²², respectively, is absent at that position.

4. A compound which is:

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

-331-

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

5 Pyrrolidine-1,2-dicarboxylic acid 1-[(2,4-difluoro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-p-tolylamide;

10 Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-[(4-methoxy-phenyl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-bromo-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-[(4-isopropyl-phenyl)-amide];

15 Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-phenylamide;

Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-[(4-trifluoromethyl-phenyl)-amide];

20 Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-[(3-methoxy-phenyl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

25 Pyrrolidine-1,2-dicarboxylic acid 1-[(4-ethyl-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(3,4-difluoro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

30 Pyrrolidine-1,2-dicarboxylic acid 1-[(3-fluoro-4-methyl-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(5-(2-sulfamoyl-phenyl)-pyridin-2-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 2-[(3-chloro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

5 Pyrrolidine-1,2-dicarboxylic acid 2-[(3-chloro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(5-(2-sulfamoyl-phenyl)-pyridin-2-yl)-amide];

10 Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-(1H-tetrazol-5-yl)-phenyl)-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

15 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Oxo-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

20 4-Methylene-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Methyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4,4-Difluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

25 4-Amino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Hydroxymethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

30 1-(4-Chloro-phenylcarbamoyl)-5-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-3-carboxylic acid;

1-(4-Chloro-phenylcarbamoyl)-5-(3-fluoro-2'-methanesulfonyl-biphenyl-4-ylcarbamoyl)-pyrrolidine-3-carboxylic acid methyl ester;

-333-

- Pyrrolidine-1,2,4-tricarboxylic acid 4-amide 1-[(4-chloro-phenyl)-amide]
2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- 4-Aminomethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- 5 4-Methoxymethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- 4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-
[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- 4-Methylamino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
10 amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- 4-Acetylamino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- 4-Methanesulfonylamino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- 15 5-Oxo-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-
[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- Piperidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-
2'-methanesulfonyl-biphenyl-4-yl)-amide];
- Morpholine-3,4-dicarboxylic acid 4-[(4-chloro-phenyl)-amide] 3-[(3-
20 fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- Piperazine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-
2'-methanesulfonyl-biphenyl-4-yl)-amide];
- 2,5-Dihydro-pyrrole-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-
[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
- 25 Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-
methanesulfonyl-3-methyl-biphenyl-4-yl)-amide];
- 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[(2'-methanesulfonyl-3-methyl-biphenyl-4-yl)-amide];
- Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-
30 methanesulfonyl-biphenyl-4-yl)-amide];
- 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[(2'-methanesulfonyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[5-(2-methanesulfonyl-phenyl)-pyrimidin-2-yl]-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[5-(2-methanesulfonyl-phenyl)-pyrimidin-2-yl]-amide];

5 Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[5-(2-methanesulfonyl-phenyl)-3-methyl-pyridin-2-yl]-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[5-(2-methanesulfonyl-phenyl)-3-methyl-pyridin-2-yl]-amide];

10 Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3-trifluoromethyl-biphenyl-4-yl)-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3-trifluoromethyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[5-(2-methanesulfonyl-phenyl)-6-methyl-pyridin-2-yl]-amide];

15 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[5-(2-methanesulfonyl-phenyl)-6-methyl-pyridin-2-yl]-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-ethyl-2'-methanesulfonyl-biphenyl-4-yl)-amide];

20 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-ethyl-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3,5-difluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3,5-dichloro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

25 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3,5-dichloro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3,5-dimethyl-biphenyl-4-yl)-amide];

30 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3,5-dimethyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 2-[(3-chloro-2'-methanesulfonyl-5-methyl-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide];

-335-

- 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(3-chloro-2'-methanesulfonyl-5-methyl-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide];
Pyrrolidine-1,2-dicarboxylic acid 2-[(2-chloro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide];
- 5 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(2-chloro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide];
Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-2-methyl-biphenyl-4-yl)-amide];
4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
10 2-[(2'-methanesulfonyl-2-methyl-biphenyl-4-yl)-amide];
Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[(2-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- 15 4-{[1-(4-Chloro-phenylcarbamoyl)-pyrrolidine-2-carbonyl]-amino}-2'-methanesulfonyl-biphenyl-3-carboxylic acid methyl ester;
4-{[1-(4-Chloro-phenylcarbamoyl)-4-hydroxy-pyrrolidine-2-carbonyl]-amino}-2'-methanesulfonyl-biphenyl-3-carboxylic acid methyl ester;
Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-[(4-trifluoromethyl-phenyl)-amide];
20 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-m-tolylamide;
Pyrrolidine-1,2-dicarboxylic acid 1-[(3-acetyl-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
- 25 Pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-2-methyl-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
Pyrrolidine-1,2-dicarboxylic acid 1-[(2-fluoro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
Pyrrolidine-1,2-dicarboxylic acid 1-[(3-chloro-4-fluoro-phenyl)-amide] 2-
30 [(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
Pyrrolidine-1,2-dicarboxylic acid 1-[(3-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

-336-

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-cyano-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-2-methyl-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

5 Pyrrolidine-1,2-dicarboxylic acid 1-[(3,4-dichloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide] 1-[(3-trifluoromethyl-phenyl)-amide];

10 Pyrrolidine-1,2-dicarboxylic acid 1-[(4-dimethylamino-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

(2S)-Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-cyano-phenyl)-amide];

15 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(4-fluoro-phenyl)-amide];

20 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-p-tolylamide;

4-Oxo-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]; and

25 4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]; or

a pharmaceutically acceptable salt thereof.

5. A compound which is:

30 (R)-Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

-337-

(S)-Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

5 (2R,4S)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

10 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(4-fluoro-phenyl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-p-tolylamide;

15 (2R)-4-Oxo-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]; and

(2R,4S)-4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide]; or a pharmaceutically acceptable salt thereof.

20

6. A compound which is:

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

25 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide};

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide};

5 (2R,4R)-4-Propoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide};

(2R,4R)-4-Propoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide};

10 (2R,4R)-4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-oxo-azetidin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(2-oxo-azetidin-1-yl)-phenyl]-amide};

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-oxo-pyrrolidin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(2-oxo-pyrrolidin-1-yl)-phenyl]-amide};

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-oxo-azepan-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(2-oxo-azepan-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-phenyl]-amide};

-339-

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{4-(2,3-dimethyl-5-oxo-2,5-dihydro-pyrazol-1-yl)-phenyl}-amide};

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
5 2-{{4-(2-hydroxymethyl-pyrrolidin-1-yl)-phenyl}-amide};

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-pyrrolidin-1-yl-phenyl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-pyrazol-1-yl-phenyl)-amide];

10 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-pyrazol-1-yl-phenyl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-[1,2,4]triazol-1-yl-phenyl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-[1,2,3]triazol-2-yl-phenyl)-amide];
15

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-[1,2,3]triazol-1-yl-phenyl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(4-acetylamino-phenyl)-amide] 1-[(4-chloro-phenyl)-amide];

20 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{4-(cyclopentanecarbonyl-amino)-phenyl}-amide}; and

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[(4-pyrimidin-5-yl-phenyl)-amide]; or

a pharmaceutically acceptable salt thereof.

25

7. A compound which is:

1-(4-Chloro-phenylcarbamoyl)-5-[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenylcarbamoyl]-pyrrolidine-3-carboxylic acid;

1-(4-Chloro-phenylcarbamoyl)-5-[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenylcarbamoyl]-pyrrolidine-3-carboxylic acid methyl ester;
30

1-(4-Chloro-phenylcarbamoyl)-5-[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenylcarbamoyl]-pyrrolidine-3-carboxylic acid ethyl ester;

-340-

- Pyrrolidine-1,2,4-tricarboxylic acid 4-amide 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 4-Cyano-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 5 4-(1H-Tetrazol-5-yl)-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 4-Aminomethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 4-Methylaminomethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 10 4-Dimethylaminomethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 4-Acetyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 15 Pyrrolidine-1,2,4-tricarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide} 4-methylamide;
- Pyrrolidine-1,2,4-tricarboxylic acid 1-[(4-chloro-phenyl)-amide] 4-dimethylamide 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 4-Trifluoromethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 20 4-Methyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 4-Isopropoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 25 4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 4,4-Difluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 1-(4-{[1-(4-Chloro-phenyl)carbamoyl]-4-methoxy-pyrrolidine-2-carbonyl}-amino)-phenyl)-pyrrolidine-2-carboxylic acid;
- 30 1-(4-{[1-(4-Chloro-phenyl)carbamoyl]-4-hydroxy-pyrrolidine-2-carbonyl}-amino)-phenyl)-pyrrolidine-2-carboxylic acid;

-341-

- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)-amide];
- 5 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[5-(2-oxo-piperidin-1-yl)-pyrimidin-2-yl]-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-(2-oxo-piperidin-1-yl)-pyrimidin-5-yl]-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[5-(2-oxo-piperidin-1-yl)-pyrimidin-2-yl]-amide];
- 10 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-(2-oxo-piperidin-1-yl)-pyrimidin-5-yl]-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide] 1-[(5-fluoro-pyridin-2-yl)-amide];
- 15 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-fluoro-pyridin-2-yl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-fluoro-pyridin-2-yl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)-amide];
- 4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 20 4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(5-fluoro-pyridin-2-yl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 25 4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)-amide];
- 4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)-amide];
- 4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(5-fluoro-pyridin-2-yl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)-amide];
- 30 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-phenyl]-amide];

- 1-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-5-oxo-pyrrolidine-2-carboxylic acid;
- 1-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-5-oxo-pyrrolidine-2-carboxylic acid;
- 5 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(4,4-dimethyl-2,5-dioxo-imidazolidin-1-yl)-2-fluoro-phenyl]-amide};
- [1-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-2,5-dioxo-imidazolidin-4-yl]-acetic acid;
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
- 10 2-{[4-(5,5-dimethyl-2,4-dioxo-oxazolidin-3-yl)-2-fluoro-phenyl]-amide};
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(2,5-dioxo-pyrrolidin-1-yl)-2-fluoro-phenyl]-amide};
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-oxo-oxazolidin-3-yl)-phenyl]-amide};
- 15 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(2,5-dioxo-oxazolidin-3-yl)-2-fluoro-phenyl]-amide};
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-oxo-imidazolidin-1-yl)-phenyl]-amide};
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
- 20 2-{[4-(2,5-dioxo-imidazolidin-1-yl)-2-fluoro-phenyl]-amide};
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(3-methyl-2,5-dioxo-imidazolidin-1-yl)-phenyl]-amide};
- 3-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-2-oxo-imidazolidine-4-carboxylic acid;
- 25 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2,4,5-trioxo-imidazolidin-1-yl)-phenyl]-amide};
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(2,5-dioxo-2,5-dihydro-pyrrol-1-yl)-2-fluoro-phenyl]-amide};
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(2-aminomethyl-5-oxo-pyrrolidin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];
- 30 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(2-aminomethyl-5-oxo-pyrrolidin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];

- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(2-carbamoyl-5-oxo-pyrrolidin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(2-carbamoyl-5-oxo-pyrrolidin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];
- 5 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(2,2-bis-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];
- 3-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-2-oxo-imidazolidine-4-carboxylic acid;
- 10 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(3,3-dimethyl-2,5-dioxo-pyrrolidin-1-yl)-2-fluoro-phenyl]-amide};
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(3,3,4-trimethyl-2,5-dioxo-pyrrolidin-1-yl)-phenyl]-amide};
- 1-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-5-oxo-pyrrolidine-2,2-dicarboxylic acid;
- 15 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-methyl-5-oxo-pyrrolidin-1-yl)-phenyl]-amide};
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-methyl-6-oxo-piperidin-1-yl)-phenyl]-amide};
- 20 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(3-hydroxy-2-oxo-piperidin-1-yl)-phenyl]-amide};
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(5-methyl-2,6-dioxo-tetrahydro-pyrimidin-1-yl)-phenyl]-amide};
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-oxo-tetrahydro-pyrimidin-1-yl)-phenyl]-amide};
- 25 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(3-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(2,5-dioxo-piperazin-1-yl)-2-fluoro-phenyl]-amide};
- 30 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(3-hydroxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[[2-fluoro-4-(3-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[[2-fluoro-4-(6-methyl-2-oxo-4-trifluoromethyl-2H-pyridin-1-yl)-phenyl]-
5 amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[[4-(2,6-dioxo-piperidin-1-yl)-2-fluoro-phenyl]-amide];
- 1-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-
amino}-3-fluoro-phenyl)-2-oxo-piperidine-3-carboxylic acid;
- 10 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[[4-(5,5-dihydroxy-2,4,6-trioxo-tetrahydro-pyrimidin-1-yl)-2-fluoro-phenyl]-
amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[[2-fluoro-4-(2,4,6-trioxo-tetrahydro-pyrimidin-1-yl)-phenyl]-amide];
- 15 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[[4-(4,4-dimethyl-2,6-dioxo-piperidin-1-yl)-2-fluoro-phenyl]-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[[2-fluoro-4-(2-hydroxy-4-oxo-azetidin-1-yl)-phenyl]-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(acetyl-methyl-amino)-
20 2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];
- 4-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-
amino}-3-fluoro-phenyl)-1H-pyrrole-3-carboxylic acid;
- 2-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-
amino}-3-fluoro-phenyl)-2H-pyrazole-3-carboxylic acid;
- 25 1-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-
amino}-3-fluoro-phenyl)-1H-imidazole-2-carboxylic acid;
- 1-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-
amino}-3-fluoro-phenyl)-1H-pyrrole-2-carboxylic acid;
- 3-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-
30 amino}-3-fluoro-phenyl)-1H-pyrrole-2-carboxylic acid;
- 3-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-
amino}-3-fluoro-phenyl)-furan-2-carboxylic acid;

- 3-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-thiophene-2-carboxylic acid;
- 4-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-furan-3-carboxylic acid;
- 5 4-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-thiophene-3-carboxylic acid;
- 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- 3-Oxo-pyrazolidine-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
- 10 3-Oxo-pyrazolidine-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[[4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 15 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 5-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 5-[[4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 20 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 5-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3,5'-difluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
- 25 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(5'-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3,5'-difluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
- 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3,5'-difluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
- 30 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(5'-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

-346-

4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[(3,5'-difluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-
amide] 2-[(3,5'-difluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

5 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[(3,5'-difluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
2-[(3,5'-difluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide], or
a pharmaceutically acceptable salt thereof.

10

8. A compound which is:

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[(4-pyrazol-1-yl-phenyl)-amide];

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[(2-fluoro-4-pyrazol-1-yl-phenyl)-amide];

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[(2-fluoro-4-pyrazol-1-yl-phenyl)-amide];

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-
amide] 2-[(4-pyrazol-1-yl-phenyl)-amide];

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[[2-fluoro-4-(3-methyl-pyrazol-1-yl)-phenyl]-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[[2-fluoro-4-(5-methyl-pyrazol-1-yl)-phenyl]-amide];

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[[4-(5-methyl-pyrazol-1-yl)-phenyl]-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[[4-(3,5-dimethyl-pyrazol-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[[4-(3,5-dimethyl-pyrazol-1-yl)-phenyl]-amide];

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[[4-(3,5-dimethyl-pyrazol-1-yl)-2-fluoro-phenyl]-amide];

-347-

- (2R,4S)-4-Amino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- (2R,4R)-4-Amino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- 5 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
- (2R)-4-Hydroxyimino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methylsulfamoyl-biphenyl-4-yl)-amide];
- 10 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-dimethylsulfamoyl-3-fluoro-biphenyl-4-yl)-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- 15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
- (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- 20 (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
- (2R,4S)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- 25 (2R,4R)-4-Fluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- (2R)-4,4-Difluoro-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide] 1-[(4-fluoro-phenyl)-amide];
- 30 (2R,4R)-4-Acetylamino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

- (2R,4R)-4-Methanesulfonylamino-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];
- (2R,4S)-4-(1H-Tetrazol-5-yl)-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 5 (2R,4S)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Cyano-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- (2R,4R)-4-(1H-Tetrazol-5-yl)-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 10 (2R,4R)-4-Trifluoromethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-cyano-3-fluoro-biphenyl-4-yl)-amide];
- 15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[(2'-aminomethyl-3-fluoro-biphenyl-4-yl)-amide] 1-[(4-chloro-phenyl)-amide];
- (2R,4R)-4'-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3'-fluoro-biphenyl-2-carboxylic acid methyl ester;
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,5-dihydro-pyrrole-1-carbonyl)-phenyl]-amide];
- 20 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-2-fluoro-phenyl)-amide];
- (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-phenyl]-amide];
- 25 Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-phenyl]-amide];
- (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2-methyl-pyrrolidine-1-carbonyl)-phenyl]-amide];
- (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(ethyl-methyl-carbamoyl)-phenyl]-amide];
- 30 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-phenyl)-amide];

- (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2R-methyl-pyrrolidine-1-carbonyl)-phenyl]-amide];
- (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2S-methyl-pyrrolidine-1-carbonyl)-phenyl]-amide];
- 5 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(pyrrolidine-1-carbonyl)-phenyl]-amide];
- (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-2-pyrrolidin-1-yl-phenyl]-amide];
- (2R,4R)-4-[[1-(4-Chloro-phenylcarbamoyl)-4-hydroxy-pyrrolidine-2-carbonyl]-amino]-3-pyrrolidin-1-yl-benzoic acid methyl ester;
- 10 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(azetidine-1-carbonyl)-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];
- (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(pyrrolidine-1-carbonyl)-phenyl]-amide];
- 15 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-phenyl]-amide];
- (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(dimethylcarbamoyl-phenyl)-amide];
- (2R,4R)-4-[[1-(4-Chloro-phenylcarbamoyl)-4-hydroxy-pyrrolidine-2-carbonyl]-amino]-3-dimethylamino-benzoic acid methyl ester;
- 20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(dimethylcarbamoyl-phenyl)-amide];
- 25 (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(pyrrolidine-1-carbonyl)-phenyl]-amide];
- (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(dimethylcarbamoyl-phenyl)-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(pyrrolidine-1-carbonyl)-phenyl]-amide];
- 30 (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(pyrrolidine-1-carbonyl)-phenyl]-amide];

-350-

(2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-2-fluoro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-methyl-4-(2-oxo-piperidin-1-yl)-phenyl)-amide];

5 (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-methyl-4-(2-oxo-piperidin-1-yl)-phenyl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-quinolin-8-yl-phenyl)-amide];

Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3,5-difluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

10 Pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-2-methyl-biphenyl-4-yl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-biphenyl-4-yl)-amide];

15 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-biphenyl-4-yl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3-methyl-2'-methylsulfonyl-biphenyl-4-yl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

20 (2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3-methyl-biphenyl-4-yl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methanesulfonyl-3-trifluoromethyl-biphenyl-4-yl)-amide];

25 5-Methyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

3-Hydroxymethyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

30 (R) 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

-351-

4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
5-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide]
5-[(3-fluoro-2'-methanesulfonyl-biphenyl-4-yl)-amide];

5 1-(4-Chloro-phenylcarbamoyl)-5-(3-fluoro-2'-methanesulfonyl-biphenyl-
4-ylcarbamoyl)-4,5-dihydro-1H-pyrazole-3-carboxylic acid ethyl ester;

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[(2'-methoxy-biphenyl-4-yl)-amide];

(2R,4R)-4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
10 phenyl)-amide] 2-[(2'-hydroxy-biphenyl-4-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[(2-fluoro-4-iodo-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[[2-fluoro-4-(3-hydroxy-2-oxo-piperidin-1-yl)-phenyl]-amide];

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-tetrahydro-pyrimidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-imidazolidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
20 phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-oxazolidin-3-yl)-phenyl]-amide];

(2R,4R)-1-(4-[[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-
carbonyl]-amino]-3-fluoro-phenyl)-2-oxo-piperidine-3-carboxylic acid ethyl
ester;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-
25 phenyl)-amide] 2-[[2-fluoro-4-(3-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl]-
amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-
pyridin-2-yl)-amide] 2-[[2-fluoro-4-(3-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl]-
amide];

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-
pyridin-2-yl)-amide] 2-[[4-(3-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

-352-

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-morpholin-4-yl-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(2-methyl-5-oxo-pyrrolidin-1-yl)-phenyl]-amide};

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[4-(2-methyl-5-oxo-pyrrolidin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(4-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(4-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(5-chloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];

15 4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 2-[(4-tert-butyl-phenyl)-amide] 1-[(4-chloro-phenyl)-amide];

4-Hydroxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3,5'-difluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];

4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)-amide];

20 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide];

4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[4-(2,3-dimethyl-5-oxo-2,5-dihydro-pyrazol-1-yl)-phenyl]-amide};

25 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(2,3-dimethyl-5-oxo-2,5-dihydro-pyrazol-1-yl)-phenyl]-amide};

4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];

4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide]-TFA Salt;

4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl)-amide];

4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide]; or

4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(6-oxo-6H-pyridazin-1-yl)-phenyl]-amide];

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(6-methyl-5-oxo-5H-[1,2,4]triazin-4-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-chloro-5-methyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3-methyl-5-oxo-4,5-dihydro-pyrazol-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(5-chloro-3-methyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3,5-dimethoxy-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-3-pyrrol-1-yl-2H-pyridin-1-yl)-phenyl]-amide];

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-3-phenyl-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-methyl-2-oxo-pyrrolidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-methyl-2-oxo-imidazolidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(3,5-dimethyl-2-oxo-imidazolidin-1-yl)-2-fluoro-phenyl]-amide};

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(3,5,5-trimethyl-2-oxo-imidazolidin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(3-isopropyl-5,5-dimethyl-2-oxo-imidazolidin-1-yl)-phenyl]-amide};

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(5-methyl-2-oxo-imidazolidin-1-yl)-phenyl]-amide};

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(5,5-dimethyl-2-oxo-imidazolidin-1-yl)-2-fluoro-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(3,4-dimethyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide};

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(4-chloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(3-methyl-2-oxo-pyrrolidin-1-yl)-phenyl]-amide};

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(3-methyl-2-oxo-imidazolidin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[4-(3,5-dimethyl-2-oxo-imidazolidin-1-yl)-2-fluoro-phenyl]-amide};

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(3,5,5-trimethyl-2-oxo-imidazolidin-1-yl)-phenyl]-amide};

-355-

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(3-isopropyl-5,5-dimethyl-2-oxo-imidazolidin-1-yl)-phenyl]-amide];

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(5-methyl-2-oxo-imidazolidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(5,5-dimethyl-2-oxo-imidazolidin-1-yl)-2-fluoro-phenyl]-amide];

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(3,4-dimethyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(4-chloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

15 or a pharmaceutically acceptable salt thereof.

9. A compound which is:

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-quinolin-1-yl)-phenyl]-amide];

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-oxo-3H-isoquinolin-2-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(1-oxo-1H-isoquinolin-2-yl)-phenyl]-amide];

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-oxo-5H-[1,2,4]triazin-4-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(6-oxo-6H-pyrimidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyrazin-1-yl)-phenyl]-amide];

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyrimidin-1-yl)-phenyl]-amide];

- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(2-oxo-2H-[1,3,5]triazin-1-yl)-phenyl]-amide}};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[6-methyl-2-oxo-2H-[1,3']bipyridinyl-6'-yl]-amide}};
- 5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(1-methyl-1H-imidazol-2-yl)-phenyl]-amide}};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(1-methyl-1H-pyrrol-2-yl)-phenyl]-amide}};
- (2R,4R)-5-Aza-spiro[2.4]heptane-5,6-dicarboxylic acid 5-[(4-chloro-phenyl)-amide] 6-{{[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide}};
- 10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(2-oxo-2H-pyridine-1-carbonyl)-phenyl]-amide}};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(3-methyl-3H-imidazol-4-yl)-phenyl]-amide}};
- 15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(2-methyl-2H-pyrazol-3-yl)-phenyl]-amide}};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[5-(2,5-dihydro-pyrrole-1-carbonyl)-pyridin-2-yl]-amide}};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(3-methyl-5-oxo-2,5-dihydro-pyrazol-1-yl)-phenyl]-amide}};
- 20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(5-methoxymethyl-3-methyl-pyrazol-1-yl)-phenyl]-amide}};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(3-methoxymethyl-5-methyl-pyrazol-1-yl)-phenyl]-amide}};
- 25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(5-ethyl-3-methyl-pyrazol-1-yl)-phenyl]-amide}};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(5-isobutyl-3-methyl-pyrazol-1-yl)-phenyl]-amide}};
- 30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(5-isobutyl-3-methyl-pyrazol-1-yl)-phenyl]-amide}};

- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(5-isopropyl-pyrazol-1-yl)-phenyl]-amide}};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(5-ethyl-pyrazol-1-yl)-phenyl]-amide}};
- 5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(5-isobutyl-pyrazol-1-yl)-phenyl]-amide}};
- (2R,4R)-2-(4-{{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolodine-2-carbonyl]-amino}}-3-fluoro-phenyl)-2*H*-pyrazole-3-carboxylic acid methyl ester;
- (2R,4R)-2-(4-{{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolodine-2-carbonyl]-amino}}-3-fluoro-phenyl)-2*H*-pyrazole-3-carboxylic acid;
- 10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(5-methoxymethyl-pyrazol-1-yl)-phenyl]-amide}};
- (2R,4R)-5-(4-{{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}}-3-fluoro-phenyl)-oxazole-4-carboxylic acid ethyl ester;
- 15 (2R,4R)-5-(4-{{[1-(4-Chloro-phenylcarbamoyl)-4-hydroxy-pyrrolidine-2-carbonyl]-amino}}-3-fluoro-phenyl)-oxazole-4-carboxylic acid ethyl ester;
- (2R,4R)-5-(4-{{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}}-3-fluoro-phenyl)-oxazole-4-carboxylic acid;
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(4-methoxymethyl-oxazol-5-yl)-phenyl]-amide}};
- 20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[4-(4-ethyl-oxazol-5-yl)-2-fluoro-phenyl]-amide}};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(4-isopropyl-oxazol-5-yl)-phenyl]-amide}};
- 25 (2R,4R)-5-(6-{{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}}-pyridin-3-yl)-oxazole-4-carboxylic acid ethyl ester;
- (2R,4R)-5-(6-{{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}}-pyridin-3-yl)-oxazole-4-carboxylic acid;
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[5-(4-methoxymethyl-oxazol-5-yl)-pyridin-2-yl]-amide}};
- 30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[5-(4-ethyl-oxazol-5-yl)-pyridin-2-yl]-amide}};

- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[5-(4-vinyl-oxazol-5-yl)-pyridin-2-yl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[5-(4-isobutyl-oxazol-5-yl)-pyridin-2-yl]-amide];
- 5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(4-methyl-isoxazol-3-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(4-methyl-isoxazol-3-yl)-phenyl]-amide];
- (2R,4R)-3-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-isoxazole-4-carboxylic acid methyl ester;
- 10 (2R,4R)-3-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-isoxazole-4-carboxylic acid;
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(4-methoxymethyl-isoxazol-3-yl)-phenyl]-amide];
- 15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(4-isopropyl-isoxazol-3-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(4-cyclopropyl-isoxazol-3-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(4-cyclopropyl-isoxazol-3-yl)-phenyl]-amide];
- 20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-methyl-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-methyl-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- 25 (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-methyl-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[(4-chloro-phenyl)-amide] 1-[[2,6-difluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[(5-chloro-pyridin-2-yl)-amide] 1-[[2,6-difluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- 30 (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 2-[(4-chloro-phenyl)-amide] 1-[[2,6-difluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

- (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 2-[(5-chloro-pyridin-2-yl)-amide] 1-[[2,6-difluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(5'-methyl-2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(5'-methyl-2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];
- (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(5'-methyl-2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];
- (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(5'-methyl-2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methyl-2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(2'-methyl-2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 15 (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2'-methyl-2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];
- (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[(2'-methyl-2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(ethyl-isopropyl-carbamoyl)-phenyl]-amide];
- 20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(ethyl-isopropyl-carbamoyl)-2-fluoro-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3-ethyl-pyrrolidine-1-carbonyl)-phenyl]-amide];
- 25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3-ethyl-pyrrolidine-1-carbonyl)-2-fluoro-phenyl]-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3R-amino-pyrrolidine-1-carbonyl)-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3R-amino-pyrrolidine-1-carbonyl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];
- 30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3S-amino-pyrrolidine-1-carbonyl)-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

-360-

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(3S-amino-pyrrolidine-1-carbonyl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(3R-methylamino-pyrrolidine-1-carbonyl)-phenyl]-amide};

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(3R-methylamino-pyrrolidine-1-carbonyl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(3S-methylamino-pyrrolidine-1-carbonyl)-phenyl]-amide};

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(3S-methylamino-pyrrolidine-1-carbonyl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(3R-ethylamino-pyrrolidine-1-carbonyl)-phenyl]-amide};

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(3R-ethylamino-pyrrolidine-1-carbonyl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(3S-ethylamino-pyrrolidine-1-carbonyl)-phenyl]-amide};

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(3S-ethylamino-pyrrolidine-1-carbonyl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[4-(2,5-dihydro-pyrrole-1-carbonyl)-phenyl]-amide};

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(2,5-dihydro-pyrrole-1-carbonyl)-2-fluoro-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[4-(2,5-dihydro-pyrrole-1-carbonyl)-2-fluoro-phenyl]-amide};

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(pyrrole-1-carbonyl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(pyrrole-1-carbonyl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[4-(pyrrole-1-carbonyl)-phenyl]-amide};

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(pyrrole-1-carbonyl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(5-fluoro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

10 (2R,4R)-1-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(6-hydroxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

15 (2R,4R)-1-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid;

(2R,4R)-1-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-6-oxo-1,6-dihydro-pyridine-2-carboxylic acid methyl ester;

20 (2R,4R)-1-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-phenyl)-2,5-dioxo-imidazolidin-4-yl]-acetic acid;

(2R,4R)-1-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid methyl ester;

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-oxo-3-trifluoromethyl-2H-pyridin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(4-hydroxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

30 (2R,4R)-1-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-6-oxo-1,6-dihydro-pyridine-2-carboxylic acid;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-5-trifluoromethyl-2H-pyridin-1-yl)-phenyl]-amide];

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(6-cyclopropyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide];

(2R,4R)-1-(4-{[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid ethyl ester;

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3,3-dimethyl-2-oxo-azetidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(7-oxo-7H-thieno[2,3-c]pyridin-6-yl)-phenyl]-amide];

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-1-aza-spiro[4.5]dec-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-benzoyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-acetylsulfamoyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-thiazolidin-3-yl)-phenyl]-amide];

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-methyl-2-oxo-piperidin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-3-sulfamoyl-2H-pyridin-1-yl)-phenyl]-amide];

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,5-dioxo-2,5-dihydro-pyrrol-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-amino-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

-363-

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-3-phenoxy-2H-pyridin-1-yl)-phenyl]-amide];

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,2-dimethyl-5-oxo-pyrrolidin-1-yl)-2-fluoro-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-nitro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

10 (2R,4R)-1-(4-[[1-(4-Chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid tert-butyl ester;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(2,5-dioxo-pyrrolidin-1-yl)-phenyl]-amide];

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-benzothiazol-2-yl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-amino-4,6-dimethyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3-diethylaminomethyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-aminomethyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-methyl-5-nitro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

30 (2R,4R)-Acetic acid 1-(4-[[1-(4-chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridin-3-yl ester;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-({4-[3-(cyclopentanecarbonyl-amino)-2-oxo-2H-pyridin-1-yl]-2-fluoro-phenyl}-amide);

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(5-amino-3-methyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide} 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-({2-fluoro-4-[2-oxo-3-(pyridin-4-ylcarbamoyl)-2H-pyridin-1-yl]-phenyl}-amide);

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{4-(3,5-dichloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(3-amino-5-bromo-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide} 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{4-(3-cyano-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{4-(3-cyano-2-oxo-5-phenyl-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{4-(3-ethyl-6-methyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{4-(3-cyano-4-dimethylamino-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(4-methyl-3-nitro-2-oxo-2H-pyridin-1-yl)-phenyl}-amide};

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(5-nitro-2-oxo-3-trifluoromethyl-2H-pyridin-1-yl)-phenyl}-amide};

-365-

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(5-amino-6-oxo-6H-[3,4']bipyridinyl-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(5-chloro-3-nitro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(3-carbamoyl-2-oxo-5-trifluoromethyl-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];

10 (2R,4R)-5-Bromo-1-(4-{[1-(4-chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid isobutyl ester;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(3-chloro-2-oxo-5-trifluoromethyl-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(3-bromo-5-chloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(5-bromo-3-nitro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(3-nitro-2-oxo-5-trifluoromethyl-2H-pyridin-1-yl)-phenyl]-amide};

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(5-methyl-3-nitro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

(2R,4R)-5-Chloro-1-(4-{[1-(4-chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid methyl ester;

30

-366-

(2R,4R)-5-Chloro-1-(4-{[1-(4-chloro-phenylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(4-benzyloxy-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(4-amino-5-fluoro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(6-methyl-3-oxo-3H-isoquinolin-2-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(4-methyl-5-nitro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(5-acetylamino-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(4-amino-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(6-oxo-6H-[3,4']bipyridinyl-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(4-bromo-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(4-oxo-4H-furo[3,2-c]pyridin-5-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(1-oxo-1H-[2,6]naphthyridin-2-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(5-bromo-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];

-367-

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(4-oxo-4H-thieno[3,2-c]pyridin-5-yl)-phenyl}-amide};

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{2-fluoro-4-(4-oxo-1,4-dihydro-imidazo[4,5-c]pyridin-5-yl)-phenyl}-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(5-chloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide} 1-[(5-chloro-pyridin-2-yl)-amide];

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{2-fluoro-4-(5-fluoro-2-oxo-2H-pyridin-1-yl)-phenyl}-amide};

(2R,4R)-1-(4-{{1-(5-Chloro-pyridin-2-ylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl}-amino}-3-fluoro-phenyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid;

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{2-fluoro-4-(6-hydroxy-2-oxo-2H-pyridin-1-yl)-phenyl}-amide};

(2R,4R)-1-(4-{{1-(5-Chloro-pyridin-2-ylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl}-amino}-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid;

20

(2R,4R)-1-(4-{{1-(5-Chloro-pyridin-2-ylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl}-amino}-3-fluoro-phenyl)-6-oxo-1,6-dihydro-pyridine-2-carboxylic acid methyl ester;

25 (2R,4R)-[1-(4-{{1-(5-Chloro-pyridin-2-ylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl}-amino}-phenyl)-2,5-dioxo-imidazolidin-4-yl]-acetic acid;

(2R,4R)-1-(4-{{1-(5-Chloro-pyridin-2-ylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl}-amino}-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid methyl ester;

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{2-fluoro-4-(2-oxo-3-trifluoromethyl-2H-pyridin-1-yl)-phenyl}-amide};

-368-

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(4-hydroxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

5 (2R,4R)-1-(4-{[1-(5-Chloro-pyridin-2-ylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-6-oxo-1,6-dihydro-pyridine-2-carboxylic acid;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(2-oxo-5-trifluoromethyl-2H-pyridin-1-yl)-phenyl]-amide};

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[4-(6-cyclopropyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide};

(2R,4R)-1-(4-{[1-(5-Chloro-pyridin-2-ylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-6-oxo-1,6-dihydro-pyridine-3-
15 carboxylic acid ethyl ester;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(3-hydroxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[4-(3,3-dimethyl-2-oxo-azetidin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(7-oxo-7H-thieno[2,3-c]pyridin-6-yl)-phenyl]-amide};

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(2-oxo-1-aza-spiro[4.5]dec-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(3-benzoyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(5-chloro-pyridin-2-yl)-amide];

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(2-methyl-5-oxo-pyrrolidin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(2-methyl-6-oxo-piperidin-1-yl)-phenyl]-amide};

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(3-acetylsulfamoyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(2-oxo-thiazolidin-3-yl)-phenyl]-amide};

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(3-methyl-2-oxo-piperidin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(2-oxo-3-sulfamoyl-2H-pyridin-1-yl)-phenyl]-amide};

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[4-(2,5-dioxo-2,5-dihydro-pyrrol-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[4-(3-amino-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide} 1-[(5-chloro-pyridin-2-yl)-amide];

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(2-oxo-3-phenoxy-2H-pyridin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[4-(2,2-dimethyl-5-oxo-pyrrolidin-1-yl)-2-fluoro-phenyl]-amide};

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(3-nitro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

(2R,4R)-1-(4-{[1-(5-Chloro-pyridin-2-yl)carbamoyl]-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid tert-butyl ester;

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[4-(2,5-dioxo-pyrrolidin-1-yl)-phenyl]-amide};

-370-

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-benzothiazol-2-yl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-amino-4,6-dimethyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(3-diethylaminomethyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide];

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-aminomethyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(3-methyl-5-nitro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

15

(2R,4R)-Acetic acid 1-(4-{[1-(5-chloro-pyridin-2-ylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino}-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridin-3-yl ester;

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-([4-[3-(cyclopentanecarbonyl-amino)-2-oxo-2H-pyridin-1-yl]-2-fluoro-phenyl]-amide);

20

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(5-amino-3-methyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-([2-fluoro-4-[2-oxo-3-(pyridin-4-ylcarbamoyl)-2H-pyridin-1-yl]-phenyl]-amide);

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(3,5-dichloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide];

30

-371-

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(3-amino-5-bromo-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide} 1-[(5-chloro-pyridin-2-yl)-amide];

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{4-(3-cyano-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{4-(3-cyano-2-oxo-5-phenyl-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide};

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{4-(3-ethyl-6-methyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{4-(3-cyano-4-dimethylamino-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide};

15

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{2-fluoro-4-(4-methyl-3-nitro-2-oxo-2H-pyridin-1-yl)-phenyl}-amide};

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{2-fluoro-4-(5-nitro-2-oxo-3-trifluoromethyl-2H-pyridin-1-yl)-phenyl}-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(5-amino-6-oxo-6H-[3,4']bipyridinyl-1-yl)-2-fluoro-phenyl}-amide} 1-[(5-chloro-pyridin-2-yl)-amide];

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(5-chloro-3-nitro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide} 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{4-(3-carbamoyl-2-oxo-5-trifluoromethyl-2H-pyridin-1-yl)-2-fluoro-phenyl}-amide} 1-[(5-chloro-pyridin-2-yl)-amide];

30

-372-

(2R,4R)-5-Bromo-1-(4-[[1-(5-chloro-pyridin-2-ylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid isobutyl ester;

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-chloro-2-oxo-5-trifluoromethyl-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-bromo-5-chloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(5-bromo-3-nitro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(3-nitro-2-oxo-5-trifluoromethyl-2H-pyridin-1-yl)-phenyl]-amide];

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(5-methyl-3-nitro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-5-Chloro-1-(4-[[1-(5-chloro-pyridin-2-ylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid methyl ester;

20 (2R,4R)-5-Chloro-1-(4-[[1-(5-chloro-pyridin-2-ylcarbamoyl)-4-methoxy-pyrrolidine-2-carbonyl]-amino]-3-fluoro-phenyl)-2-oxo-1,2-dihydro-pyridine-3-carboxylic acid;

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(4-benzyloxy-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(4-amino-5-fluoro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

30

-373-

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(5-nitro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(6-methyl-3-oxo-3H-isoquinolin-2-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(4-methyl-5-nitro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(5-acetylamino-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(1-oxo-1H-isoquinolin-2-yl)-phenyl]-amide];

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(4-amino-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(6-oxo-6H-[3,4']bipyridinyl-1-yl)-phenyl]-amide];

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(3-oxo-3H-isoquinolin-2-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(4-bromo-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(4-oxo-4H-furo[3,2-c]pyridin-5-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(1-oxo-1H-[2,6]naphthyridin-2-yl)-phenyl]-amide];

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(5-bromo-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(4-oxo-4H-thieno[3,2-c]pyridin-5-yl)-phenyl]-amide};

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(4-oxo-1,4-dihydro-imidazo[4,5-c]pyridin-5-yl)-phenyl]-amide};

4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-{[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

10 3-Hydroxymethyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-{[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

5-Methyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-{[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

3-Methoxymethyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-{[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

15 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 5-{[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

3-Hydroxymethyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 5-{[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

20 5-Methyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 5-{[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

3-Methoxymethyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 5-{[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

25 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-{[4-(2-methyl-thiazol-4-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-2-methyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide};

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(3,4,5,6-tetrahydro-2H-[1,4']bipyridinyl-4-ylmethyl)-amide];

-375-

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[4-(2-dimethylaminomethyl-imidazol-1-yl)-2-fluoro-phenyl]-amide}};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{[2-chloro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{{[2-chloro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide} 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide}};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(2-methylamino-imidazol-1-yl)-phenyl]-amide}};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{[2-fluoro-4-(2-isopropyl-imidazol-1-yl)-phenyl]-amide}};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[4-(2-ethyl-imidazol-1-yl)-2-fluoro-phenyl]-amide}};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{{[4-(2-ethyl-imidazol-1-yl)-2-fluoro-phenyl]-amide}};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(2-isopropyl-imidazol-1-yl)-phenyl]-amide}};

(2R,4R)-2-Ethyl-4-methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide}};

4-Methoxy-2-methoxymethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide}};

2-Hydroxymethyl-4-methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide}};

{1-(4-Chloro-phenylcarbamoyl)-2-[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenylcarbamoyl]-4-methoxy-pyrrolidin-2-yl}-acetic acid methyl ester;

{1-(4-Chloro-phenylcarbamoyl)-2-[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenylcarbamoyl]-4-methoxy-pyrrolidin-2-yl}-acetic acid;

4-Hydroxymethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{{[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide}};

-376-

{ 1-(4-Chloro-phenylcarbamoyl)-5-[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenylcarbamoyl]-pyrrolidin-3-yl}-acetic acid;

{ 1-(4-Chloro-phenylcarbamoyl)-5-[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenylcarbamoyl]-pyrrolidin-3-yl}-acetic acid methyl ester;

5 (2R,4R)-4-Ethyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Isopropyl-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-fluoro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(3-fluoro-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-chloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-chloro-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-bromo-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(4-chloro-phenyl)-amide];

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-[[4-(3-bromo-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide] 1-[(5-chloro-pyridin-2-yl)-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-iodo-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(3-iodo-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[4-(3-ethyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide]; and

30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3-ethyl-2-oxo-2H-pyridin-1-yl)-2-fluoro-phenyl]-amide;

a pharmaceutically acceptable salt thereof.

-377-

10. A compound which is:
- (2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];
- 5 (2R,4R) 4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- (2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- (2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-[[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- 10 (2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];
- (2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(3-methoxy-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- 15 (2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[4-(3-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide];
- (2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-methyl-imidazol-1-yl)-phenyl]-amide];
- 20 (2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-2-fluoro-phenyl)-amide];
- (2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(5-methyl-pyrazol-1-yl)-phenyl]-amide];
- (2R,4R) 4,5-Dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
- 25 (2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-fluoro-4-pyrazol-1-yl)-phenyl]-amide];
- (2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(pyrrolidine-1-carbonyl)-phenyl]-amide];
- 30 (2R,4R) 4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[[2-fluoro-4-(2-oxo-piperidin-1-yl)-phenyl]-amide];

- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide};
- 5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(3-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-methyl-imidazol-1-yl)-phenyl]-amide};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(2-oxo-2H-[1,3']bipyridinyl-6'-yl)-amide];
- 10 (2R,4R)-4-Ethoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-[(4-dimethylcarbamoyl-2-fluoro-phenyl)-amide];
- 15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(5-methyl-pyrazol-1-yl)-phenyl]-amide};
- 5-Methyl-4,5-dihydro-pyrazole-1,5-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 5-[(3-fluoro-2'-sulfamoyl-biphenyl-4-yl)-amide];
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(4-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};
- 20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(3-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 2-{[2-chloro-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide} 1-[(4-chloro-phenyl)-amide];
- 25 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-methyl-4-(2-oxo-2H-pyridin-1-yl)-phenyl]-amide};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(6-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};
- (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(3-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};
- 30 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-

-379-

pyridin-2-yl)-amide] 2-{[2-fluoro-4-(5-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(5-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

5 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(5-chloro-pyridin-2-yl)-amide] 2-{[2-fluoro-4-(4-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(2,5-dihydro-pyrrole-1-carbonyl)-2-fluoro-phenyl]-amide};

10 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(2,5-dihydro-pyrrole-1-carbonyl)-phenyl]-amide}

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(2-methyl-5-oxo-pyrrolidin-1-yl)-phenyl]-amide};


15 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(2,5-dihydro-pyrrole-1-carbonyl)-phenyl]-amide};

(2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[2-fluoro-4-(4-methyl-2-oxo-2H-pyridin-1-yl)-phenyl]-amide};

20 (2R,4R)-4-Methoxy-pyrrolidine-1,2-dicarboxylic acid 1-[(4-chloro-phenyl)-amide] 2-{[4-(2,5-dihydro-pyrrole-1-carbonyl)-2-fluoro-phenyl]-amide};

or a pharmaceutically acceptable salt thereof.

11. A pharmaceutical composition comprising a compound of formula I



I

25

or a pharmaceutically acceptable salt thereof wherein:

-380-

A is aryl or substituted aryl or monocyclic heteroaryl or substituted monocyclic heteroaryl;

B is $\text{-NH}-\overset{\text{O}}{\parallel}{\text{C}}-(\text{C}_1\text{-C}_6)\text{alkyl}$, $\text{-NH}-\overset{\text{O}}{\parallel}{\text{C}}-(\text{C}_3\text{-C}_7)\text{cycloalkyl}$ $\text{-NH}(\text{C}_1\text{-}$

$\text{C}_6)\text{alkanoyl}$, $\text{-C}(=\text{O})\text{R}^1$, $(\text{C}_3\text{-C}_7)\text{cycloalkyl}$, $(\text{C}_3\text{-C}_7)\text{heterocycloalkyl}$, $(\text{C}_4\text{-C}_7)\text{cycloalkenyl}$, $(\text{C}_4\text{-C}_7)\text{heterocycloalkenyl}$, aryl, or heteroaryl, any of which may be optionally substituted;

C is phenyl or heteroaryl, wherein phenyl or heteroaryl is optionally substituted with one or more substituents selected from halogen, hydroxy, $\text{-CO}_2\text{R}^2$, -COR^2 , $\text{-CONR}^2\text{R}^{2'}$, alkoxy, alkyl, -CN , haloalkyl, amino, alkylamino, amidino, amido, or sulfonamido;

G is H, halo, $(\text{C}_1\text{-C}_6)\text{alkyl}$, halo $(\text{C}_1\text{-C}_6)\text{alkyl}$, hydroxy $(\text{C}_1\text{-C}_6)\text{alkyl}$, $\text{-CH}_2\text{O}-(\text{C}_1\text{-C}_6)\text{alkyl}$, $\text{-CH}_2\text{-CO}_2(\text{C}_1\text{-C}_6)\text{alkyl}$, $\text{-CH}_2\text{-CONH}_2$, or $\text{-CH}_2\text{-CONH}(\text{C}_1\text{-C}_6)\text{alkyl}$;

W^1 is a saturated or unsaturated, substituted or unsubstituted hydrocarbon chain or hydrocarbon-heteroatom chain having from 2 to 6 atoms, wherein W^1 connects the nitrogen atom at position 1 to the carbon atom at position 2 to form a four to eight membered ring;

R^1 is $(\text{C}_1\text{-C}_6)\text{alkoxy}$, $(\text{C}_3\text{-C}_7)\text{cycloalkyl}$, $(\text{C}_3\text{-C}_7)\text{heterocycloalkyl}$, $(\text{C}_4\text{-C}_7)\text{cycloalkenyl}$, $(\text{C}_4\text{-C}_7)\text{heterocycloalkenyl}$, aryl, monocyclic heteroaryl, or $\text{-NR}^3\text{R}^4$;

R^2 and $\text{R}^{2'}$ are each independently H or $(\text{C}_1\text{-C}_6)\text{alkyl}$; and

R^3 and R^4 are each independently H, $(\text{C}_1\text{-C}_6)\text{alkyl}$, aralkyl, aryl, monocyclic heteroaryl, alkoxycarbonyl, aralkoxycarbonyl, $\text{-SO}_2\text{alkyl}$, or joined together to form a saturated or unsaturated 3 to 7 membered ring;

admixed with a carrier, diluent, or excipient.

-381-


12. A method for preventing or treating acute, subacute, and chronic thrombotic disorder in a mammal comprising administering to a mammal in need thereof a therapeutically effective amount of a compound of claim 1.

5 13. The method of claim 10, wherein the acute, subacute, and chronic thrombotic disorder is venous thrombosis, arterial thrombosis, pulmonary embolism, myocardial infarction, cerebral infarction, restenosis, cancer, angina, diabetes, atrial fibrillation, or heart failure.

10 14. A method for inhibiting Factor Xa in a mammal, comprising administering to a mammal in need of Factor Xa inhibition a Factor Xa inhibition amount of a compound of claim 1.

15. A process for preparing a compound of Formula I:

15



or a pharmaceutically acceptable salt thereof wherein:

20 A is aryl or substituted aryl or monocyclic heteroaryl or substituted monocyclic heteroaryl;

B is $\text{-NH-C(=O)-(C}_1\text{-C}_6\text{)alkyl}$, $\text{-NH-C(=O)-(C}_3\text{-C}_7\text{)cycloalkyl}$, $\text{-NH(C}_1\text{-C}_6\text{)alkanoyl}$, -C(=O)R^1 , $\text{(C}_3\text{-C}_7\text{)cycloalkyl}$, $\text{(C}_3\text{-C}_7\text{)heterocycloalkyl}$, $\text{(C}_4\text{-C}_7\text{)cycloalkenyl}$, $\text{(C}_4\text{-C}_7\text{)heterocycloalkenyl}$, aryl, or heteroaryl, any of which may

25 be optionally substituted by halo, $\text{(C}_1\text{-C}_6\text{)alkyl}$, or $\text{halo(C}_1\text{-C}_6\text{)alkyl}$, $\text{(C}_1\text{-C}_6\text{)alkenyl}$, $\text{(C}_3\text{-C}_7\text{)cycloalkenyl}$, $\text{(C}_3\text{-C}_7\text{)heterocycloalkenyl}$, $\text{(C}_4\text{-C}_7\text{)cycloalkenyl}$, $\text{(C}_4\text{-C}_7\text{)heterocycloalkenyl}$, aryl, or heteroaryl, any of which may

-382-

C_6)alkoxy, -CN, haloalkyl, amino, alkylamino, amidino, amido, or sulfonamido;

C is phenyl or heteroaryl, wherein phenyl or heteroaryl is optionally substituted with one or more substituents selected from halogen, hydroxy, - CO_2R^2 , - COR^2 , - $CONR^2R^{2'}$, alkoxy, alkyl, -CN, haloalkyl, amino, alkylamino,

5 amidino, amido, or sulfonamido;

G is H, halo, (C_1-C_6) alkyl, halo (C_1-C_6) alkyl, hydroxy (C_1-C_6) alkyl, - $CH_2O-(C_1-C_6)$ alkyl, - $CH_2CO_2(C_1-C_6)$ alkyl, - $CH_2NR^2R^{2'}$, or - $CH_2CONH(C_1-C_6)$ alkyl;

10 W^1 is a saturated or unsaturated, substituted or unsubstituted hydrocarbon chain or hydrocarbon-heteroatom chain having from 2 to 6 atoms, wherein W^1 connects the nitrogen atom at position 1 to the carbon atom at position 2 to form a four to eight membered ring;


R^1 is (C_1-C_6) alkoxy, (C_3-C_7) cycloalkyl, (C_3-C_7) heterocycloalkyl, (C_4-C_7) cycloalkenyl, (C_4-C_7) heterocycloalkenyl, aryl, monocyclic heteroaryl, or -
15 NR^3R^4 ;

R^2 and $R^{2'}$ are each independently H or (C_1-C_6) alkyl; and

R^3 and R^4 are each independently H, (C_1-C_6) alkyl, aralkyl, aryl, monocyclic heteroaryl, alkoxycarbonyl, aralkoxycarbonyl, - SO_2 alkyl, or joined together to form a saturated or unsaturated 3 to 7 membered ring;

20 comprising:


(a) reacting an amino acid of formula IA



25


with a reagent capable of placing a protecting group P^1 on the amino group of an amino acid to form a protected amino acid given by formula IB:

-383-



IB

- 5 wherein P¹ is a protecting group and W¹ is the same as defined above;
- (b) converting the acid moiety in the compound of formula IB to a acid halide;
- 10 (c) reacting the acid halide with a haloaniline or a haloamino-heterocycle to form a compound having formula IC:




IC

- 15 wherein Z₁ is a halogen and A is as defined above;
- (d) coupling the compound of formula IC with a compound H-B to give a compound having formula ID:

20

-384-



- 5 (e) removing the protecting group from Compound ID and reacting the resulting acid moiety with a C-isocyanate to form a compound of the invention.

INTERNATIONAL SEARCH REPORT

 ii
 al Application No
 PCT/IB 02/04757

A. CLASSIFICATION OF SUBJECT MATTER

 IPC 7 C07D207/16 C07D207/26 C07D213/64 C07D401/12 C07D401/14
 C07D403/12 A61K31/40 A61P7/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

 Minimum documentation searched (classification system followed by classification symbols)
 IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 01 64642 A (COR THERAPEUTICS, INC) 7 September 2001 (2001-09-07) page 160 -page 296; examples 1-641 ---	1-15
Y	WO 97 46576 A (GYOGYSZERKUTATO INTEZET KFT) 11 December 1997 (1997-12-11) page 65 -page 66; claims 1-8 ---	1-15
Y	WO 97 05161 A (SANDOZ-PATENT-GMBH) 13 February 1997 (1997-02-13) page 11 -page 12; claims 1-7 --- -/-	1-15

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the International filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the International search

26 March 2003

Date of mailing of the International search report

10/04/2003

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Kyriakakou, G

INTERNATIONAL SEARCH REPORT

Inte

Application No

PCT/IB 02/04757

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	F AL-OBEIDI & JAOSTREM: "Factor Xa Inhibitors" EXPERT OPINION ON THERAPEUTIC PATENTS., vol. 9, no. 7, 1999, pages 931-953, XP001000512 ASHLEY PUBLICATIONS., GB ISSN: 1354-3776 page 941 -page 946	1-15
Y	ERIK VERNER ET AL.: "Development of Serine Protease Inhibitors Displaying a Multicenter Short Hydrogen Bond Binding Mode: Inhibitors of Urokinase-Type Plasminogen Activator and Factor Xa" JOURNAL OF MEDICINAL CHEMISTRY., vol. 44, no. 17, 2001, pages 2753-2771, XP002216503 AMERICAN CHEMICAL SOCIETY., US ISSN: 0022-2623 page 2754; example 4	1-15
Y	MICHAEL R. WILEY ET AL.: "Structure-Based Design of Potent, Amidine-Derived Inhibitors of Factor Xa: Evaluation of Selectivity, Anticoagulant Activity, and Antithrombotic Activity" JOURNAL OF MEDICINAL CHEMISTRY., vol. 43, no. 5, 2000, pages 883-899, XP002186966 AMERICAN CHEMICAL SOCIETY., US ISSN: 0022-2623 the whole document	1-15
Y	YING K.YEE ET AL.: "N-Aroylanthranilamide Inhibitors of Human Factor Xa" JOURNAL OF MEDICINAL CHEMISTRY., vol. 43, no. 5, 2000, pages 873-882, XP002186965 AMERICAN CHEMICAL SOCIETY., US ISSN: 0022-2623 the whole document	1-15
Y	WO 95 28420 A (CORVAS INTERNATIONAL, INC) 26 October 1995 (1995-10-26) page 212 -page 239; claims 1-76; figures 1-3	1-15

INTERNATIONAL SEARCH REPORT

International application No.
PCT/IB 02/04757

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 12-14 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

In al Application No
PCT/18 02/04757

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0164642	A	07-09-2001	AU 4535301 A	12-09-2001
			AU 5078301 A	12-09-2001
			AU 7486700 A	17-04-2001
			BR 0014076 A	15-10-2002
			CN 1391555 T	15-01-2003
			CZ 20020961 A3	14-08-2002
			EP 1216228 A2	26-06-2002
			EP 1259485 A2	27-11-2002
			HU 0203289 A2	28-01-2003
			NO 20021229 A	21-05-2002
			WO 0119788 A2	22-03-2001
			WO 0164642 A2	07-09-2001
			WO 0164643 A2	07-09-2001
			US 2002002183 A1	03-01-2002
WO 9746576	A	11-12-1997	HU 9601526 A2	28-10-1998
			AT 222261 T	15-08-2002
			AU 726516 B2	09-11-2000
			AU 3043697 A	05-01-1998
			DE 69714745 D1	19-09-2002
			DK 922054 T3	09-12-2002
			EP 0922054 A1	16-06-1999
			ES 2182086 T3	01-03-2003
			WO 9746576 A1	11-12-1997
			JP 2000512278 T	19-09-2000
			KR 2000016348 A	25-03-2000
			PT 922054 T	31-12-2002
			US 6235707 B1	22-05-2001
WO 9705161	A	13-02-1997	AU 6737996 A	26-02-1997
			WO 9705161 A1	13-02-1997
WO 9528420	A	26-10-1995	US 5776927 A	07-07-1998
			US 5681844 A	28-10-1997
			AT 195947 T	15-09-2000
			AU 2360995 A	10-11-1995
			DE 69518648 D1	05-10-2000
			EP 0765340 A1	02-04-1997
			JP 9512022 T	02-12-1997
			WO 9528420 A1	26-10-1995
			US 5658939 A	19-08-1997
			US 5770600 A	23-06-1998
			US 5646165 A	08-07-1997

